Climate Change and Veterinary Parasitology: Emerging Risks to Livestock Health and Sustainable Disease Management.

Abstract:

Parasitic diseases affect livestock health, productivity, animal welfare, and farm profitability. Climate conditions can influence the development, survival, and distribution of many parasites and the vectors or intermediate hosts involved in their transmission. Changes in temperature, rainfall, humidity, and seasonal patterns may therefore alter disease risks in some regions. However, these effects vary according to parasite species, host animals, local ecology, management practices, and access to veterinary services. This paper examines the relationship between climate change and veterinary parasitology, with particular attention to livestock gastrointestinal parasites, ticks, and vector-borne diseases. A narrative literature review approach is used to synthesize established knowledge on parasite ecology, livestock management, surveillance, and disease prevention. The paper discusses the importance of veterinary diagnosis, strategic parasite control, pasture management, biosecurity, and responsible use of antiparasitic medicines. It also considers the challenges of drug resistance, limited diagnostic capacity, and unequal access to veterinary care. The review concludes that climate-informed surveillance and integrated parasite management can support livestock health, provided that interventions are based on local evidence and professional veterinary guidance.

Keywords: Veterinary parasitology, climate change, livestock health, parasites, ticks, disease surveillance, anthelmintic resistance, sustainable animal production.

1. Introduction

Livestock contributes to food security, household income, and agricultural livelihoods worldwide. Maintaining animal health is essential for sustainable livestock production, and parasitic diseases remain an important challenge in many production systems.

Parasites may affect animals internally or externally. Internal parasites include gastrointestinal worms and other organisms that can interfere with digestion, nutrient absorption, and general health. External parasites, including ticks and mites, may cause irritation, skin damage, blood loss, or transmit infectious agents.

The development and survival of many parasites are influenced by environmental conditions. Temperature and moisture can affect parasite eggs, larvae, vectors, and intermediate hosts. Consequently, changes in climate may alter the timing, geographical distribution, or intensity of some parasitic diseases.

UNDER PEER REVIEW IN JAVES

These changes are not uniform. Some parasites may become more prevalent in certain locations, while others may decline because of unsuitable environmental conditions. Host immunity, animal movement, grazing practices, veterinary access, and land use also influence disease patterns.

Understanding these relationships is important for developing effective veterinary surveillance and sustainable control strategies.

2. Objectives of the Study

1. To explain how climatic conditions influence parasites and their transmission.
2. To review important parasitic challenges affecting livestock.
3. To examine the role of veterinary surveillance and diagnosis.
4. To discuss integrated and responsible parasite-control strategies.
5. To identify research priorities for climate-informed veterinary healthcare.

3. Review of Literature

Veterinary research has established that parasite development and transmission depend on interactions between the parasite, host, and environment. Temperature and moisture influence many free-living parasite stages, while climatic conditions can affect the distribution and activity of ticks and other vectors.

The World Organisation for Animal Health emphasizes surveillance, prevention, and coordinated approaches to animal disease management. The Intergovernmental Panel on Climate Change has also discussed how climate change can affect animal health and agricultural systems, although outcomes vary by region and disease.

Antiparasitic drug resistance is another major concern. Repeated use of the same medicines without suitable diagnosis or management can contribute to reduced treatment effectiveness in some parasite populations.

The literature supports integrated approaches that combine diagnosis, appropriate treatment, pasture management, animal nutrition, and monitoring rather than relying exclusively on routine medication.

4. Research Methodology

This paper uses a narrative literature review design. It synthesizes established information on climate-sensitive parasite ecology, livestock disease management, veterinary surveillance, and antiparasitic resistance.

The discussion is organized around climatic influences, livestock health risks, diagnosis, prevention, and future research. No original veterinary examinations, animal trials, or field surveillance data were collected.

5. Climate and Parasite Development

UNDER PEER REVIEW IN JNAVES

5.1 Temperature

Temperature can influence the development rate and survival of parasite eggs, larvae, and vectors. Within suitable limits, warmer conditions may accelerate the development of some parasites. Temperatures beyond their tolerance ranges, however, may reduce survival.

These responses vary considerably among species. It is therefore inaccurate to assume that all parasitic diseases will increase as temperatures rise.

5.2 Rainfall and Humidity

Moisture is important for the survival of many free-living parasite stages. Rainfall may create conditions that support development and transmission, while prolonged dryness may reduce survival for moisture-sensitive organisms.

Heavy rainfall can also change grazing patterns, animal movement, and contact with contaminated environments. The resulting disease effects depend on local conditions.

5.3 Seasonal Changes

Changes in seasonal temperature and rainfall may alter the timing of parasite exposure. Traditional treatment schedules may become less effective if parasite activity changes, making local monitoring increasingly important.

Veterinary professionals can help farmers adjust surveillance and control plans to reflect observed seasonal patterns.

5.4 Host and Environmental Interactions

Parasite transmission depends on more than climate. Animal density, grazing management, wildlife interactions, host immunity, and veterinary care can strongly influence the outcome.

Climate-informed management should therefore consider the entire livestock system rather than attributing every change in disease occurrence to climate alone.

6. Important Parasitic Challenges in Livestock

6.1 Gastrointestinal Parasites

Gastrointestinal worms can affect cattle, sheep, goats, and other animals. Depending on the parasite and severity of infection, affected animals may experience poor growth, reduced productivity, digestive disturbances, or anemia.

Young animals and poorly nourished animals may be especially vulnerable to certain infections. Diagnosis and treatment should be based on veterinary assessment and, where appropriate, fecal testing.

6.2 Tick Infestations

UNDER PEER REVIEW IN JNAVS

Ticks can cause irritation, skin damage, blood loss, and reduced productivity. Some species also transmit pathogens responsible for important livestock diseases.

Temperature and moisture can affect tick development and seasonal activity, but the response differs by species and region. Surveillance should identify the ticks present and assess local disease risks.

6.3 Vector-Borne Parasitic Diseases

Some parasitic diseases depend on insect or tick vectors. Changes in temperature, rainfall, and habitat may influence vector populations and their contact with livestock.

Disease risk also depends on the presence of the parasite, host susceptibility, animal movement, and vector-control measures. Veterinary authorities should monitor changes in disease patterns rather than relying on climate assumptions alone.

6.4 Effects on Productivity

Parasitic infections may reduce feed efficiency, milk production, growth, fertility, and work capacity in affected animals. Severe infections can result in substantial animal-health and economic consequences.

Preventive health management can reduce avoidable losses and improve animal welfare.

7. Veterinary Diagnosis and Surveillance

7.1 Clinical Examination

Veterinarians assess clinical signs, body condition, production records, grazing history, and local disease risks. These observations help identify animals that require further investigation.

Clinical signs alone may not reliably identify a particular parasite, so laboratory confirmation may be necessary.

7.2 Laboratory Testing

Fecal examinations, blood tests, microscopic identification, and other diagnostic methods can help detect parasites or evidence of infection. The appropriate test depends on the suspected parasite and the stage of disease.

Laboratory testing also supports more informed decisions about treatment and control.

7.3 Monitoring Disease Trends

Farmers and veterinary services can maintain records of parasite findings, treatment outcomes, seasonal conditions, and animal performance. Reviewing these records may reveal changes in local disease patterns.

UNDER PEER REVIEW IN IJNAVES

Surveillance systems are particularly useful when combined with timely communication between farms, laboratories, and veterinary authorities.

8. Integrated Parasite Management

8.1 Targeted Treatment

Antiparasitic medicines can be important for treating confirmed or strongly suspected infections. Their selection and use should follow veterinary guidance, product instructions, and relevant regulations.

Unnecessary or poorly planned treatment can increase costs and contribute to drug resistance. Treatment effectiveness should be monitored where possible.

8.2 Pasture Management

Appropriate grazing management may reduce exposure to certain parasites. Depending on the parasite and livestock system, strategies can include avoiding excessive stocking density, managing grazing periods, and maintaining suitable pasture conditions.

No pasture method eliminates every parasite. Recommendations should reflect the biology of the parasite and local management conditions.

8.3 Nutrition and Animal Resilience

Adequate nutrition supports immune function and general health. Animals receiving balanced feed and clean water may be better able to withstand some disease challenges, although nutrition cannot replace effective diagnosis and treatment.

Feed plans should account for species, age, production stage, and local resource availability.

8.4 Biosecurity and Animal Movement

Newly introduced animals may carry parasites into a herd or flock. Veterinary assessment and suitable introduction procedures can help reduce this risk.

Animal movement should be managed in accordance with local disease-control requirements and farm biosecurity plans.

8.5 Resistance Management

Resistance can develop when parasites survive treatments and reproduce despite exposure to medicines. Its occurrence varies by parasite species, medicine, and region.

Veterinarians can help design control programs that use diagnostic information, appropriate drug selection, and monitoring. Farmers should not independently change medicine doses or combine treatments without professional guidance.

UNDER PEER REVIEW IN INAVES

9. Technology and Climate-Informed Veterinary Services

Weather information and geographic data may help veterinary authorities identify periods or locations requiring increased surveillance. Digital reporting can improve communication between farmers and veterinary services.

Diagnostic laboratories and mobile veterinary services may be especially useful in remote areas. However, digital systems are only effective when the underlying data are reliable and farmers can access and understand the information.

Future monitoring systems should combine climatic information with confirmed disease observations, animal movement records, and local veterinary expertise.

10. Economic and Social Considerations

Parasite control can involve expenditure on veterinary visits, diagnostic testing, medicines, labor, and pasture management. These costs may be difficult for small-scale farmers, particularly when veterinary services are distant.

Cooperative veterinary programs, farmer education, and accessible diagnostic services may reduce some barriers. Community-based reporting can also help identify emerging disease patterns.

Interventions should be practical for the local production system and should account for farmers' financial circumstances.

11. Challenges and Research Gaps

Major challenges include limited diagnostic facilities, inadequate disease reporting, antiparasitic resistance, and insufficient information about local climate-parasite relationships.

Research priorities include:

- Monitoring parasite distributions over time.
- Evaluating climate-related changes in disease seasonality.
- Improving accessible diagnostic methods.
- Assessing drug resistance in different livestock systems.
- Studying the combined influence of climate, nutrition, and management.
- Developing locally appropriate parasite-control plans.
- Strengthening collaboration among veterinarians, researchers, and farmers.

Long-term research should distinguish observed associations from demonstrated causal relationships.

12. Recommendations

- Strengthen veterinary surveillance and laboratory diagnostic capacity.

UNDER PEER REVIEW IN JNAVES

- Use antiparasitic medicines under appropriate professional guidance.
- Maintain accurate animal-health and treatment records.
- Improve nutrition, water access, and general husbandry.
- Develop parasite-control plans based on local disease patterns.
- Monitor treatment effectiveness and suspected resistance.
- Improve communication between farmers and veterinary authorities.
- Expand farmer education about parasites and prevention.
- Integrate climate information with confirmed veterinary observations.

13. Conclusion

Climate change may alter the development, distribution, and seasonality of some livestock parasites, but the effects vary according to parasite biology, host characteristics, local ecology, and management.

Sustainable parasite control requires accurate diagnosis, responsible use of veterinary medicines, suitable grazing management, good nutrition, and reliable surveillance. Climate information can support these activities when interpreted alongside local veterinary evidence.

Strengthening veterinary services and farmer education can improve preparedness for changing disease risks. Integrated, locally appropriate management offers a practical way to protect animal welfare, farm productivity, and rural livelihoods.

References

1. Intergovernmental Panel on Climate Change. (2022). *Climate Change 2022: Impacts, Adaptation and Vulnerability*. Cambridge University Press. https://www.ipcc.ch/report/ar6/wg2/
2. World Organisation for Animal Health. (n.d.). *Animal health and welfare*. https://www.woah.org/
3. Food and Agriculture Organization of the United Nations. (n.d.). *Animal health*. https://www.fao.org/animal-health/en/
4. Kaplan, R. M., & Vidyashankar, A. N. (2012). An inconvenient truth: Global warming and anthelmintic resistance. *Veterinary Parasitology, 186*(1–2), 70–78. https://doi.org/10.1016/j.vetpar.2011.11.048
5. Taylor, M. A., Coop, R. L., & Wall, R. L. (2016). *Veterinary Parasitology* (4th ed.). Wiley-Blackwell.

UNDER PEER REVIEW IN IJAVS