

1 Migration of heavy metals in the upper horizons of mining soil : the case of 2 the former Poura mine (Burkina Faso)

3 4 5 **ABSTRACT**

6 The presence of contaminants in an environment and their mobility depend greatly on the
7 nature of the environment, climatic and geological conditions, and anthropogenic influences.
8 Soil contamination by these heavy metals poses a significant public health risk due to the
9 various diseases they can cause. The objective of this study was to examine the migration of
10 heavy metals through their concentrations in the upper horizons of a mining soil. Soil profiles
11 were taken between the two piles of mining waste known as tailings 1 and tailings 2, as well
12 as between these tailings and a cyanide tank, all located on the mine site. The soil profiles had
13 four layers or horizons identified by color. The dimensions were 1,5 m deep, 0,90 m wide,
14 and 1,5 m high. The results show that profile 1 was generally the least contaminated with
15 metals. Cadmium (Cd) stands out due to its accumulation in profile 2 (74,07 mg/kg). Arsenic
16 (As) has very low levels in all profiles. It is practically zero in profiles 1 and 2, while a low
17 average value of 0,33 mg/kg is observed in profile 3. Chromium (Cr) concentrations are high
18 in all three profiles, with 87,62 mg/kg in profile 1, 134,09 mg/kg in pit 2, and 133,88 mg/kg in
19 profile 3. Zinc (Zn) levels are very low in all profiles, with a measurable value only in profile
20 2 (0,09 mg/kg). Manganese (Mn) is the predominant metal. Its concentrations increase sharply
21 from profile 1 (713,17 mg/kg) to profile 2 (1689,11 mg/kg), reaching a maximum in profile 3
22 (7349,53 mg/kg). Nickel (Ni) follows an increasing gradient, with 21,51 mg/kg in profile 1,
23 26,27 mg/kg in profile 2, and 185,22 mg/kg in profile 3. Lead (Pb) remains low overall, with
24 no presence in profile 1, a concentration of 2,68 mg/kg in profile 2, and 0,8 mg/kg in profile
25 3. These heavy metals (Mn, Ni, Cr, Cu, and As). The contamination hierarchy observed from
26 the heat map was established as follows: profile 3, profile 2, then profile 1. This trend
27 highlights a higher metal accumulation in profile 3, while profile 1 appears to be the least
28 impacted.

29 **Keywords:** Heavy metals, migration, soil profile, horizon, Poura mine, Burkina Faso

30 31 32 33 34 **INTRODUCTION**

35 The recent development of the mining industry in Burkina Faso has led to a considerable
36 increase in gold production, making it the country's leading source of export revenue ahead of
37 cotton. The country's annual gold production rose from 5,8 tons in 2008 to 14 tons in 2009
38 (SCADD, 2012), reaching 62 tons in 2019 and 74 tons in 2020 (Yao, 2021). This sharp
39 increase in production, resulting from the commissioning of 17 industrial mines over the last
40 decade, has helped Burkina Faso maintain its position as Africa's fifth largest gold producer.

41 In addition to these industrial facilities, there are hundreds of artisanal and semi-industrial
42 operators scattered throughout the country. In 2016, there were already 448 operational
43 artisanal gold production sites spread across all regions of the country except the central
44 region (INSD, 2017). Specifically, in certain precious metal deposits, gold or silver are
45 associated with metalloids elements such as arsenic, which is known for its toxicity in the
46 environment. This arsenic can then be exposed during ore excavation and take on a toxic
47 form, posing a danger to the environment and human health (Matschullat, 2000). In other
48 deposits, the natural oxidation of sulfide minerals in rock when exposed to air and water
49 causes acid mine drainage (AMD), which contributes to various types of pollution. All of
50 these forms of pollution result in reduced access to drinking water, loss of soil fertility,
51 reduced vegetation cover, loss of biodiversity, and habitat fragmentation (Schueler et al.,
52 2011).


53 This study therefore aims to examine the vertical migration of metals in the soils of a former
54 mine in Burkina Faso. Our specific objectives include analyzing the distribution of heavy
55 metals in soil profile horizons, as well as comparing concentrations between horizons on the
56 one hand and between layers on the other. We will present the methodology in detail, before
57 presenting and discussing the results obtained, and finally making recommendations.

58

59 **MATERIALS AND METHODS**

60 *Location of the study site*

61 The study site is focused on the former Poura mine. Poura is a rural commune in the Balé
62 province, located southwest of Ouagadougou, 106 miles from the Ouaga-Bobo highway and
63 27,7 miles from Boromo, the provincial capital. It is located between 11°20' and 11°50' north
64 latitude and between 2°40' and 2°55' west longitude. The commune of Poura covers an area of
65 101,81 km². A single road (Regional Road RR11) connects Poura to National Road (RN) N°
66 1.



67

68 **Figure 1:** Map showing the location of the study area in the municipality of Poura

69

70

71 ***Soils at the study site***

72 The soils of Poura are mainly tropical ferruginous soils, which are shallow, not very fertile,
 73 and susceptible to erosion (Yaméogo *et al.*, 2024). These soils are common in the region and
 74 can be classified into several groups.

75 - Tropical ferruginous soils: located near watercourses, these soils are associated with water
 76 conditions and are often used for agriculture.

77 - Ferralitic soils: found in the provinces of Houet, Kéné Dougou, Comoé, and Mouhoun, these
 78 soils contain clays rich in iron and aluminum.

79 - Brown soils: located in the western, southwestern, north-central, and northwestern parts of
 80 the country, these soils are often used for various crops.

81 - Raw mineral soils or lithosols: formed directly from bedrock, these soils are often poorly
 82 developed and less suitable for agriculture.

83 - Vertisols: rich in clay, these soils are capable of swelling and contracting, and are often used
 84 for growing cotton and cereals.


85 - Sodic soils: found in the south-central and north-central regions, these sodium-rich soils can
 86 cause salinity problems.

87 - Hydromorphic soils: located in the far north of Burkina Faso, these soils are well drained but
 88 can become acidic over time.

89

90 **Soil profile sampling**

91 In order to monitor the different levels or layers of soil through which pollutants migrate,
92 three (03) soil profiles were created between the tailings and between each tailing and the
93 cyanide tank.



94 **Figure 2.** Map showing the location of pits or soil profiles.
95

96 Soil pit 3 is located halfway between the cyanide tank and tailings 1, in an area where gold
97 panning is actively taking place. Soil pit 1 is also located halfway between tailings 2 and the
98 cyanide tank, in the middle of a cornfield that is being invaded by gold panning activities. Soil
99 pit 3 could not be placed halfway between tailings 1 and tailings 2. This is because the point
100 where the pit should be located is occupied by waste rock from the old mine. It has been
101 repositioned a little closer to tailings 1. It is located in an area of intense gold panning activity.

102 In the construction of the soil pits, four layer levels were identified based on color and coarse
103 elements for each pit considered (RECORD, 2006). Pits 1.5 meters deep, 0.90 meters wide,
104 and 1.5 meters long were dug. These different layers were sampled according to the model
105 developed by Laperche, V. and Eisenlohr, L. (2001). A total of twelve (12) layers were
106 identified, at a rate of four (4) layers per pit.

107 **Statistical data analysis**

108 The data were entered using Excel 2016 spreadsheet software. R software version 4.3.1 (R
109 Core Team, 2023) was used for data processing. The distribution of heavy metal values in the
110 different samples was checked for normality and homogeneity using Shapiro-Wilk tests.
111 Student's t-tests were used as appropriate to compare the content per horizon; ANOVA and
112 Kruskal-Wallis tests were used as appropriate to compare the variation in parameter content
113 according to horizon. Tukey's post-hoc test was used to compare the means. The results were
114 interpreted at the 5% significance level.

115


116 **RESULTS**

117 **Heavy metal content in soil profile horizons**

118 *Arsenic levels*

119 The arsenic contents in the horizons of the soil profiles are shown in Figure 3. Except for the
120 first horizon of the third soil profile, which has an arsenic content of 1,3 mg/kg, none of the
121 horizons have an arsenic content. However, this content is below the threshold level for the
122 Earth's crust, which is 1,5 mg/kg.

123



124

125 **Figure 3.** Arsenic content in the horizons (layers) of soil profiles (pits)

126

127 ***Cadmium levels***

128 Figure 4 shows the cadmium content of soil samples in the different horizons of the soil
 129 profiles. Apart from the second horizon (layer) of soil profile 2, which recorded a high Cd
 130 content (296,3 mg/kg), no other horizon showed any Cd content. This cadmium content of
 131 296,3 mg/kg obtained in layer C2 is well above that of the Earth's crust, which is 0,2 mg/kg.

132

133




Figure 4. Cadmium content in soil profile layers

Chromium levels


The results of the analysis of soil samples from the different layers of the soil profiles are shown in Figure 5.

In profile 1, there is a variation in content between the different layers. The highest content is observed in C2 and the lowest content is observed in C4, with respective values of 128.6 mg/kg and 1.6 mg/kg. Of the four horizons in profile 1, only horizon C4 has levels above the Earth's crust threshold of 70 mg/kg.

145 At profile 2, an upward trend in content is observed. All horizons of this profile have
146 chromium contents that are above those of the Earth's crust.

147 In pit 3, an increase in chromium content is noted from layer C2 to layer C4, with respective
148 values of 66,9 mg/kg (C2), 110,4 mg/kg (C3), and 276,8 mg/kg (C4). Layer C1 has a
149 concentration of 81,4 mg/kg. Apart from layer C2, the other layers have concentrations above
150 that of the Earth's crust.

151



152

153 **Figure 5.** Chromium content in soil profile horizons

154


155

156 **Copper levels**

157 The results of the analysis of samples from the layers of the copper soil pits are shown in
158 Figure 6.

159 Regardless of the soil profile, copper content increased with depth. However, in profile 1, a
160 notable decrease is observed between C3 and C4. The highest content is recorded at horizon 4
161 (C4) of profile 3 (pit 3) with a content of 121 mg/kg Cu.

162



163

164 **Figure 6.** Copper content in soil profile horizons

165


166

167 **Manganese levels**

168 The various results of the analysis of manganese content in soil samples from different layers
169 of the soil pits are shown in Figure 7.

170 Manganese is present in all four horizons of the three soil profiles. The highest value is found
171 in horizon 3 (C3) of profile 3, with 13827,7 mg/kg, while the lowest content is recorded in
172 layer C4 of profile 1, with 3,8 mg/kg. Apart from layer C4 of profile 1, which has the lowest
173 value (3,8 mg/kg), all other horizons have manganese contents higher than that of the Earth's
174 crust (600 mg/kg).

175




176

177 **Figure 7.** Manganese content in soil profile horizons

178 *Nickel levels*

179 Figure 8 shows the nickel content in soil samples from different layers of the soil profiles.

180 Analysis of the results shows an absence of nickel in layer C3 of profile 1 and layers C3 and
181 C4 of profile 2. The highest Ni content is found in layer C3 of profile 3, with 300,9 mg/kg of
182 Ni. It should also be noted that the different layers of profile 3 have higher Ni contents than
183 the other two soil profiles.



184

185 **Figure 8.** Nickel content in soil profile horizons

186


187

188 **Lead levels**

189 The various results of the analysis of the lead content of soil samples from different layers of
190 the soil profiles are shown in Figure 9.

191 The results show that lead is present in only two layers out of the twelve horizons. These are
192 layer C4 of soil profiles 2 and 3, with 10,7 mg/kg and 3,2 mg/kg of Pb, respectively.
193 However, none of these levels exceed that of the Earth's crust, which is 20 mg/kg of Pb.

194



195

196 **Figure 9.** Lead content in soil profile layers

197


198

199 **Zinc levels**

200 Figure 10 shows the zinc content results for soil samples from different layers of the soil
201 profiles.

202 Zinc is present only in layer C2 of the second soil profile. The other two profiles show no zinc
203 content in any horizon. Furthermore, the recorded zinc content of 0,3 mg/kg remains lower
204 than that of the Earth's crust, which is 71 mg/kg Zn.

205



206

207 **Figure 10.** Zinc contents in soil profiles

208

209

210 **Comparison of heavy metal content in soils in soil profiles**

211 Normalizing the values of As, Cd, Cr, Cu, Mn, Ni, Pb, and Zn concentrations on a scale of 0
212 to 1 allowed for a comparison of contamination levels between soil profiles 1, 2, and 3
213 (Figure 11).


214 Profile 1 was found to be the least contaminated with metals overall. Most elements showed
215 values close to 0 on the normalized scale, particularly As, Cd, Pb, and Zn. Concentrations of
216 Cr, Cu, Mn, and Ni were present but remained relatively low compared to the other profiles.

217 Profile 2 showed an intermediate level of contamination. Several metals reached high values
218 on the standardized scale, particularly Cd and Pb, which were close to 1. Cr and Zn also
219 showed significant intensities compared to other sources.

220 Profile 3 stood out with the highest levels for several metals. These heavy metals (Mn, Ni, Cr,
221 Cu, and As) reached values close to 1 on the standardized scale, indicating that it concentrates
222 the relative maxima for these elements. Although some metals such as Cd and Zn are low in
223 this profile, the high intensity of several major elements reflects a higher metal load in the
224 other two pits.

225 Ultimately, the contamination hierarchy observed from the heat map was established as
226 follows: profile 3, profile 2, then profile 1. This trend highlights a higher metal accumulation
227 in profile 3, while profile 1 appears to be the least affected.

228



229

230 **Figure 11.** Comparison of contamination intensity between soil profiles 1, 2, and 3

231


232 Figure 12 shows a comparison of heavy metals between the three soil profiles (pit). Arsenic
 233 (As) has very low levels in all profiles. It is practically absent in profiles 1 and 2, while a low
 234 average value of 0,33 mg/kg is observed in profile 3.

235 Cadmium (Cd) stands out due to its accumulation in profile 2 (74,07 mg/kg), while it is absent
 236 in profiles 1 and 3, suggesting sporadic contamination. As for chromium (Cr), concentrations
 237 are high in all three profiles, with 87,62 mg/kg in profile 1, 1134,09 mg/kg in pit 2, and 133,88
 238 mg/kg in profile 3. Profiles 2 and 3 show comparable levels that are higher than those in
 239 profile 1.

240 Copper (Cu) concentrations increase gradually from 37,48 mg/kg in profile 1 to 61,08 mg/kg
 241 in profile 2 and 82,64 mg/kg in profile 3. Zinc (Zn) levels are very low in all profiles, with a
 242 measurable value only in profile 2 (0,09 mg/kg), indicating a marginal presence.

243 Manganese (Mn) is the predominant metal. Its concentrations increase sharply from profile 1
 244 (713,17 mg/kg) to profile 2 (1689,11 mg/kg), reaching a maximum in profile 3 (7349,53
 245 mg/kg). This trend is accompanied by a wide dispersion of values, especially in profile 3.

246 Nickel (Ni) follows an increasing gradient, with 21,51 mg/kg in profile 1, 26,27 mg/kg in
 247 profile 2, and 185,22 mg/kg in profile 3. The high dispersion observed in profile 3 reflects
 248 significant accumulation. Lead (Pb) remains low overall, with no presence in profile 1, a
 249 concentration of 2,68 mg/kg in profile 2, and 0,8 mg/kg in profile 3.



250

251 **Figure 12.** Comparison by heavy metal in each soil pit

252 **DISCUSSION**

253 The study of soil profiles made it possible to examine the vertical migration of the various
254 heavy metals studied. Most of the pollutants are concentrated in the surface layer (0-30 cm) in
255 the immediate vicinity of the sources. Our results on the soil horizons affected by this
256 pollution are consistent with those of Traore *et al.* (2014). However, there is one exception:
257 manganese in profile 3, located between tailings 1 and the cyanide tank. In this profile,
258 manganese contamination increases with depth. The vertical migration of this contaminant
259 indicates a potential risk of groundwater contamination (Bouzahzahet *al.*, 2014). This result
260 highlights a dynamic dispersion of pollutants, characterized by areas of intense pollution and
261 selective mobility of certain metals in specific locations. Targeted remediation measures and
262 continuous environmental monitoring are necessary to mitigate the level of contamination
263 from these identified heavy metals.

264

265 **CONCLUSION**

266 The soil at abandoned former mining sites, if not properly maintained or rehabilitated,
267 becomes a dangerous source of environmental pollution. The chemicals used in mining
268 (cyanide, acids, peroxide, etc.), and especially the various mining waste products, can be
269 washed away by climatic (air and rain) and anthropogenic phenomena, leading to the leaching
270 of various mining pollutants (cyanide and heavy metals). This leaching can lead to the
271 generation of acid mine drainage (AMD) and contaminate soil, water, vegetation, and the
272 ecosystem in general.

273 Analysis of the results obtained by analyzing the ETM content in the different horizons of the
274 three soil profiles reveals that the highest concentrations are located in the 0 to 30 cm
275 horizons. This finding reveals surface migration of metal pollutants through the topsoil
276 horizons of the former mining site of Poura in Burkina Faso (West Africa).

277

278

279 **REFERENCES**

- 280 1. Bouzahzah, H., Benzaazoua, M., Bussière, B., Plante, B.,(2014). Revue de littérature
281 détaillée sur les tests statiques et les essais cinétiques comme outils de prédiction du
282 drainage minier acide. Environnement, Ingénierie & Développement 66,14-31.

283 <https://doi.org/10.4267/dechets-sciences-techniques.340>

- 284 2. INSD. (2017). Enquête nationale sur le secteur de l'orpaillage (ENSO). Principaux
285 résultats. DSSE/SCEAM 2017-04.
- 286 3. Laperche, V., Eisenlohr, L. (2001) - Echantillonnage d'un site contaminé par l'arsenic
287 - Comparaison de différentes stratégies et représentativité de l'échantillonnage des
288 sols. Rapport CNRSSP/01/11, 33 p.
- 289 4. Matschullat, J. (2000). Arsenic in the Geosphere. A Review. The Science of the Total
290 Environment, 249, 297-312. [http://dx.doi.org/10.1016/S0048-9697\(99\)00524-0](http://dx.doi.org/10.1016/S0048-9697(99)00524-0)
- 291 5. RECORD, 2006. Stratégie et technique d'échantillonnage des sols pour l'évaluation
292 des pollutions. 321 p, n° 04-0510/1A.
- 293 6. SCADD. (2012). Stratégie de croissance accélérée et de développement durable,
294 Burkina Faso 2011 -2015. 108p.
- 295 7. Schueler, V., Kuemmerle, T., Schröder, H. (2011). Impacts of surface gold mining on
296 land use systems in Western Ghana. *Ambio*, 40, 528-539.
- 297 8. Traore A., Soro G., Kouadio E. K, Bamba B.S., Oga MS, et Soro N. et Biemi J.
298 (2014). Niveau de contamination en métaux lourds des sédiments d'une lagune
299 tropicale : la lagune ghien (Sud-Est de la Côte d'Ivoire. *Afrique Sc. Revue*
300 *Internationale et Technologie*, 10(3), 73-
301 88, <http://www.afriquescience.info/document.php?id=3720>
- 302 9. Yameogo A., Kagambega N., Bado I.F., Barry A.A., 2024. Trace Metal Elements
303 (TMEs) in Groundwater Around the Former Industrial Gold Mine of Poura in Burkina
304 Faso –West Africa. *International Journal of Environmental Monitoring and Analysis*
305 2024, Vol. 12, No. 5, pp. 111-124
- 306 10. Yao, O. (2021). La référence économique en Afrique de l'Ouest. Sika Finance.
307 [https://www.sikafinance.com/marches/burkina-faso-la-production-industrielle-d-or-](https://www.sikafinance.com/marches/burkina-faso-la-production-industrielle-d-or-atteint-plus-de-27-tonnes-a-fin-mai-2021_29785)
308 [atteint-plus-de-27-tonnes-a-fin-mai-2021_29785](https://www.sikafinance.com/marches/burkina-faso-la-production-industrielle-d-or-atteint-plus-de-27-tonnes-a-fin-mai-2021_29785), Consulté le 19/07/2022 à 20h04
309