

# 1 Incidence and Risk Factors of Perioperative Respiratory Incidents 2 in Pediatric Anesthesia: A Prospective Observational Single-Center 3 Study 4

## 5 ABSTRACT

---

6 **Introduction/Objective:** Perioperative respiratory incidents are a leading cause of morbidity in  
7 pediatric anesthesia. Few prospective studies from low- and middle-income countries have  
8 simultaneously evaluated patient-, surgery-, and anesthesia-related determinants of these events. This  
9 study aimed to determine their incidence and identify independent risk factors in a Moroccan pediatric  
10 surgical cohort.

11 **Methods:** A six-month prospective observational single-center study (April–September 2025) was  
12 conducted at Abderrahim El Harrouchi Children’s Hospital, CHU Ibn Rochd, Casablanca. A total of  
13 470 children undergoing surgery under general anesthesia were enrolled. Risk factors were identified  
14 by univariate analysis followed by binary logistic regression.

15 **Results:** The incidence of perioperative respiratory incidents was 26.8% (126/470). Desaturation  
16 ( $SpO_2 < 90\%$ ) was the most common event (92.9%). On multivariate analysis, three independent risk  
17 factors were identified: upper airway obstruction (adjusted OR = 3.47; 95% CI [1.9–6.3];  $p < 0.001$ ),  
18 multiple intubation attempts (adjusted OR = 5.94; 95% CI [2.6–13.7];  $p < 0.001$ ), and agitation at  
19 induction (adjusted OR = 2.05; 95% CI [1.1–3.8];  $p = 0.04$ ).

20 **Discussion:** The incidence observed is consistent with comparable African settings and substantially  
21 higher than high-income country data, reflecting emergency case burden, comorbidities, and resource  
22 constraints. The three identified factors are all clinically modifiable and amenable to targeted  
23 prevention strategies.

24 **Conclusion:** Perioperative respiratory incidents are frequent in our setting. Systematic preoperative  
25 airway screening, structured difficult intubation protocols, and anxiolytic premedication are priority  
26 interventions for improving pediatric anesthetic safety.

27 **Keywords:** pediatric anesthesia; respiratory incidents, bronchospasm, intubation  
28

## 29 1. INTRODUCTION

---

30 Pediatric anesthetic safety is a major concern in anesthesiology. Children have unique anatomical and  
31 physiological characteristics — narrow airways, reduced thoracic compliance, elevated oxygen  
32 consumption, and rapid metabolism of anesthetic agents — that expose them to greater risk of  
33 perioperative respiratory incidents than adults [1,2].

34 Perioperative respiratory incidents (PRIs) include a spectrum of events: oxygen desaturation ( $SpO_2 < 90\%$ ),  
35 laryngospasm, bronchospasm, upper airway obstruction (UAO), apnea, and mask ventilation  
36 difficulty. Although usually reversible, these events may be life-threatening when not promptly  
37 recognized and managed [3,4].

38 The APRICOT study, conducted across more than 31,000 children in 261 European hospitals,  
39 reported a 5.2% incidence of severe critical respiratory events [4]. The rates seen in Europe cannot be  
40 projected onto middle income countries where the rates tend to be higher because there is a larger  
41 number of emergency procedures, and limited availability of specialized airway equipment, and limited  
42 available personnel [5,6].

43 In Morocco, epidemiological data specific to pediatric anesthesia remain scarce. Understanding the  
44 incidence and determinants of PRIs is therefore essential to guide local clinical practice and resource  
45 allocation.

46 Therefore, the main objective of this study was to determine the incidence of PRIs in pediatric  
47 anesthesia at our institution. Secondary objectives were to identify independent risk factors and  
48 provide recommendations to prevent such occurrences.

49

## 50 2. MATERIALS AND METHODS

---

### 51 2.1 Study design and setting

52 We conducted a prospective observational single-center study over six consecutive months, from April  
53 to September 2025, in the pediatric surgical operating rooms of Abderrahim El Harrouchi Children's  
54 Hospital, CHU Ibn Rochd, Casablanca, Morocco.

### 55 2.2 Study population

56 **Inclusion criteria:** all patients under 14 years of age undergoing surgery under general anesthesia  
57 during the study period.

58 **Exclusion criteria:** age  $\geq 14$  years; refusal of parental or legal guardian informed consent; incomplete  
59 anesthetic records.

### 60 2.3 Data collection

61 Data were collected prospectively using a standardized case report form (CRF) completed in the  
62 operating room by the attending anesthesiologist. Variables recorded included: (i) demographic data  
63 (age in months, sex); (ii) preoperative data (surgical specialty, elective vs. emergency status, ASA  
64 physical status score, history of allergy, Mallampati score, mouth opening, cervical spine mobility,  
65 signs of upper airway obstruction, simplified pediatric difficult intubation score, preoperative anxiety  
66 level); and (iii) intraoperative data (type of anesthetic induction, anesthetic agents, Cormack–Lehane  
67 grade, number of intubation attempts, agitation at induction, occurrence and type of respiratory  
68 incident, and surgical duration in hours).

### 69 2.4 Definitions

70 **Perioperative respiratory incident** was defined as the occurrence of at least one of the following  
71 events from induction to end of surgery: oxygen desaturation ( $SpO_2 < 90\%$ ), laryngospasm,  
72 bronchospasm, mask ventilation difficulty, multiple or failed intubation attempts, apnea, perioperative  
73 bradycardia, or hypotension associated with a respiratory event [3,4]. A patient could present with more  
74 than one type of incident simultaneously.

75 **Upper airway obstruction (UAO):** defined by preoperative clinical signs: snoring, stridor, mouth  
76 breathing, or documented adenotonsillar hypertrophy.

77 **Agitation at induction:** any oppositional or agitated behavior preventing adequate peripheral venous  
78 access or face-mask placement.

79 **Multiple intubation attempts:** two or more laryngoscopy attempts required for tracheal intubation.

## 80 **2.5 Bias recognition**

81 To minimize selection bias, standardized inclusion and exclusion criteria were applied prospectively.  
82 Information bias was reduced by using a structured, pre-defined case report form (CRF) completed at  
83 the bedside by the attending anesthesiologist throughout each procedure. Confounding was  
84 addressed through multivariate logistic regression, adjusting for all variables with  $p < 0.20$  on  
85 univariate analysis. Observer bias was minimized by providing a written operational definition of each  
86 outcome variable to all data collectors prior to data collection.

## 87 **2.6 Statistical analysis**

88 Statistical analysis was performed using **Jamovi version 2.6.44** (The Jamovi Project, Sydney,  
89 Australia). Continuous variables are expressed as median and interquartile range (IQR); categorical  
90 variables as absolute numbers and percentages. Univariate analysis was conducted for each  
91 candidate risk factor to preliminarily identify associations. Then variables with a  $p$ -value  $< 0.20$  in the  
92 univariate analysis were entered into a multivariable logistic regression model to control for  
93 confounding and identify independent predictors of complications. Adjusted odds ratios (aORs) with  
94 95% confidence intervals (CIs) were calculated to quantify the strength of association between risk  
95 factors and complications. Statistical significance was set at a  $p$ -value  $< 0.05$ .

## 96 **3. RESULTS**

### 97 **3.1 Study population**

98 A total of 470 patients were enrolled. The most represented age group was 5–12 years (46.8%). The  
99 male-to-female ratio was 1.94. Orthopedic and trauma surgery was the leading surgical specialty  
100 (45.1%), followed by visceral surgery (28.9%) and urological and neurosurgical procedures (26.0%).  
101 Emergency procedures accounted for 44.0% of cases. ASA class I–II was recorded in 75.1% of  
102 patients and Mallampati score I–II in 63.0%. Signs of upper airway obstruction were present in 18.1%  
103 of patients. Demographic and intraoperative characteristics of the entire cohort are summarized in  
104 **Table 1**.

105 **Table 1.** Demographic and Intraoperative Characteristics of Patients (n = 470).

Variables	n (%)
<b>Patient characteristics</b>	
Gender M / F	310 (66.0%) / 160 (34.0%)
Age group	
• < 12 months	48 (10.2%)
• 1–5 years	117 (24.9%)
• 5–12 years	220 (46.8%)
• > 12 years	85 (18.1%)
ASA physical status I–II	353 (75.1%)
ASA physical status III–IV	117 (24.9%)
Mallampati score I–II	296 (63.0%)
Mallampati score III–IV	89 (18.9%)
Mallampati non-assessable	85 (18.1%)
Signs of upper airway obstruction	85 (18.1%)
<b>Surgical factors</b>	
Surgical specialty	


• Orthopedic / trauma	212 (45.1%)
• Visceralsurgery	136 (28.9%)
• Urological / neurosurgery	122 (26.0%)
Emergency procedure	207 (44.0%)
Surgical duration > 2 hours	98 (20.9%)
<b>Anestheticfactors</b>	
Type of anesthetic induction	
• Intravenous	308 (65.5%)
• Combined (IV + inhalational)	162 (34.5%)
Multiple intubation attempts ( $\geq 2$ )	70 (14.9%)
Agitation at induction	90 (19.1%)
Ketamine use	65 (13.8%)

106 Values are presented as n (%). ASA: American Society of Anesthesiologists. Mallampati score could  
 107 not be assessed in uncooperative patients (n = 85).

### 108 3.2 Incidence and distribution of perioperative respiratory incidents

109 One hundred and twenty-six patients (26.8%) experienced at least one perioperative respiratory  
 110 incident. A single patient could present with more than one type of incident simultaneously. Oxygen  
 111 desaturation ( $SpO_2 < 90\%$ ) was the most frequent event, occurring in 92.9% of affected patients  
 112 (117/126), most commonly in association with bronchospasm. The distribution of incident types is  
 113 illustrated in Fig. 1.

114



115

116 **Fig. 1.** Distribution of perioperative respiratory incident types among affected patients (n = 126).

117 A patient may present with more than one type of incident simultaneously.

118

### 119 3.3 Factors associated with perioperative respiratory incidents — Univariate analysis

120 Univariate logistic regression identified nine significant risk factors for PRI (**Table 2**): age < 12 months  
 121 (crude OR = 6.28; p < 0.001), upper airway obstruction (OR = 5.11; p < 0.001), multiple intubation  
 122 attempts  $\geq 2$  (OR = 16.99; p < 0.001), Mallampati score III–IV (OR = 3.83; p < 0.001), combined  
 123 induction (OR = 3.42; p < 0.001), agitation at induction (OR = 3.08; p < 0.001), ASA class III–IV (OR =

124 2.72;  $p < 0.001$ ), surgical duration  $> 2$  hours (OR = 2.04;  $p = 0.004$ ), and emergency surgery (OR =  
 125 1.65;  $p = 0.004$ ). Male sex (OR = 1.15;  $p = 0.52$ ) and ketamine use (OR = 1.25;  $p = 0.45$ ) were not  
 126 significantly associated with PRI. All variables with  $p < 0.20$  were retained for multivariate analysis.

127  
 128  
 129  
 130

131 **Table 2.** Univariate logistic regression analysis of factors associated with perioperative respiratory  
 132 incidents ( $n = 470$ ).

Variable	Crude OR	95% CI	p-value
<b>Patient characteristics</b>			
Male sex	1.15	[0.74–1.78]	0.52
Age $< 12$ months	6.28	[3.33–11.83]	$< 0.001$
ASA class III–IV	2.72	[1.74–4.25]	$< 0.001$
Mallampati score III–IV †	3.83	[2.30–6.38]	$< 0.001$
Upper airway obstruction	5.11	[3.11–8.38]	$< 0.001$
<b>Surgical and anesthetic factors</b>			
Emergency surgery	1.65	[1.10–2.50]	0.004
Surgical duration $> 2$ hours	2.04	[1.28–3.28]	0.004
Combined induction (IV + inhalational)	3.42	[2.24–5.24]	$< 0.001$
Multiple intubation attempts ( $\geq 2$ )	16.99	[9.09–31.77]	$< 0.001$
Agitation at induction	3.08	[1.91–4.98]	$< 0.001$
Ketamine use	1.25	[0.71–2.22]	0.45

133 OR: Crude Odds Ratio; CI: 95% Confidence Interval. Estimated by univariate binary logistic  
 134 regression.  $p$ -values from Wald test. † Mallampati score assessed in 385/470 patients; uncooperative  
 135 patients ( $n = 85$ ) excluded from this analysis.

### 136 3.4 Independent risk factors — Multivariate logistic regression

137 All variables with  $p < 0.20$  on univariate analysis were entered into a binary logistic regression model  
 138 using a stepwise backward elimination strategy. After adjustment, three independent risk factors were  
 139 identified (**Table 3**): upper airway obstruction (adjusted OR = 3.47; 95% CI [1.9–6.3];  $p < 0.001$ ),  
 140 multiple intubation attempts  $\geq 2$  (adjusted OR = 5.94; 95% CI [2.6–13.7];  $p < 0.001$ ), and agitation at  
 141 induction (adjusted OR = 2.05; 95% CI [1.1–3.8];  $p = 0.04$ ). Model goodness-of-fit was satisfactory  
 142 (Hosmer–Lemeshow  $p = 0.42$ ).

143 **Table 3.** Multivariate logistic regression — Independent predictors of perioperative respiratory  
 144 incidents.

Clinical risk factor	Adjusted OR	95% CI	p-value
Upper airway obstruction	3.47	[1.9–6.3]	$< 0.001$
Multiple intubation attempts ( $\geq 2$ )	5.94	[2.6–13.7]	$< 0.001$
Agitation at induction	2.05	[1.1–3.8]	0.04

145 OR: Adjusted Odds Ratio; CI: 95% Confidence Interval. Hosmer–Lemeshow goodness-of-fit test:  $p =$   
 146 0.42.

## 147 4. DISCUSSION

### 148 4.1 Incidence of perioperative respiratory incidents

149 The incidence of PRIs in our series (26.8%) is consistent with prospective data from comparable  
150 African settings. Wudineh et al. reported a prevalence of 26.2% in a prospective study of 210 pediatric  
151 surgical patients in Northwest Ethiopia [7], and Hordofa et al. similarly identified an incidence of 29.8%  
152 in a 2024 prospective study of 205 pediatric patients in southern Ethiopia [18], further confirming the  
153 elevated burden of perioperative respiratory events in sub-Saharan African contexts. In contrast, Tay  
154 et al. found a 2.97% critical incident rate (77.4% of which were respiratory events) in Singapore [8].  
155 These figures contrast markedly with high-income country data: the APRICOT study reported a 5.2%  
156 incidence of severe critical respiratory events across 261 European hospitals [4], while the  
157 NECTARINE study documented that one third of neonates and young infants undergoing anaesthesia  
158 across 31 European countries required an intervention for a severe critical event [19]. In the ambulatory  
159 setting, Subramanyam et al. reported 3.1% in a large US cohort [9]. These disparities are largely  
160 attributable to the high proportion of emergency surgeries in our cohort (44%), the greater prevalence  
161 of nutritional and respiratory comorbidities, and resource constraints limiting access to advanced  
162 airway management equipment.

#### 163 **4.2 Upper airway obstruction**

164 UAO was the most prevalent independent risk factor in our model (adjusted OR = 3.47).  
165 Adenotonsillar hypertrophy is the leading etiology of UAO in children, generating bronchial  
166 hyperreactivity and airway lability that predispose to laryngospasm and bronchospasm under general  
167 anaesthesia [10,11]. These findings are aligned with the established literature demonstrating that children  
168 with UAO have a threefold to fivefold elevated risk of perioperative respiratory complications. Notably,  
169 a pre-existing pulmonary disorder appears as one of the 11 predictors of the recently validated  
170 SPORC-C score, reinforcing the importance of systematic preoperative airway assessment [20].  
171 Combined (IV + inhalational) induction was also significantly associated with PRIs on univariate  
172 analysis, consistent with data showing that inhalational agents increase airway reactivity in children  
173 with underlying bronchial hyperresponsiveness. Ramgolam et al. demonstrated in a randomized  
174 controlled trial that intravenous induction significantly reduced the risk of respiratory adverse events  
175 compared to inhalational or combined induction in high-risk children [10].

#### 176 **4.3 Multiple intubation attempts**

177 Multiple intubation attempts were the strongest independent predictor in our model (adjusted OR =  
178 5.94). Each additional laryngoscopy attempt causes mucosal trauma, laryngeal edema, and increased  
179 airway reactivity, cumulatively raising the risk of desaturation, bronchospasm, and laryngospasm [12,13].  
180 The rate of multiple intubation attempts in our series (14.9%) likely reflects limited access to pediatric  
181 videolaryngoscopy and flexible fiberoptic equipment, and the absence of formally implemented  
182 difficult airway management algorithms. Konrad et al. demonstrated that structured training protocols  
183 significantly reduced the number of intubation attempts and associated complications [14].  
184 Implementation of such protocols, combined with appropriate equipment availability, constitutes a  
185 high-priority safety intervention in our setting.

#### 186 **4.4 Agitation at induction**

187 Agitation at induction was an independent risk factor in our series (adjusted OR = 2.05), with 19.1% of  
188 children affected. Agitation impairs adequate face-mask placement and preoxygenation, significantly  
189 increasing the risk of early desaturation. Preoperative anxiety is a well-established precipitant,  
190 particularly in children under

191 3 years and in those undergoing inhalational induction [15]. Oral midazolam premedication and  
192 intramuscular ketamine are validated strategies to reduce induction agitation [16,17]. A recent systematic  
193 review and meta-analysis of 7 categories of prophylactic pharmacological interventions in children  
194 undergoing noncardiac surgery confirmed the efficacy of several agents in reducing overall PRAEs  
195 and their subtypes including laryngospasm, bronchospasm, and oxygen desaturation [21]. Of interest,  
196 ketamine showed no significant association with PRI in univariate analysis (crude OR = 1.25; p =  
197 0.45), though this does not exclude a clinically meaningful effect given the limited number of exposed  
198 patients (n = 65). Its bronchodilatory properties and capacity to preserve upper airway reflexes remain  
199 a plausible mechanistic rationale [16]. The limited number of exposed patients (n = 65) precluded a  
200 definitive conclusion, and this hypothesis warrants investigation in a dedicated randomized controlled  
201 trial.

202 **4.5 Other associated factors**

203 Young age, elevated ASA score, and emergency surgery were strongly associated with PRIs on  
204 univariate analysis, reflecting findings from prior literature. Infants under 12 months had a particularly  
205 high incidence (64.6%), attributable to anatomophysiological specificities including a relatively large  
206 occiput, high laryngeal position, reduced thoracic compliance, and predisposition to obstructive  
207 apneas [1,2]. These factors were not retained in the final multivariate model, likely due to collinearity  
208 with the variables that were retained.

209 **4.6 Limitations**

210 This study has several limitations. Its single-center monocentric design limits generalizability to the  
211 national level. Prospective observational data collection may have been subject to variable detection  
212 bias across operators. Missing Mallampati scores in uncooperative patients (18.1%) may have  
213 influenced the analyses. The composite endpoint groups events of variable clinical severity, which  
214 limits the interpretation of certain associations. Finally, potential confounders such as recent upper  
215 respiratory tract infection and passive tobacco smoke exposure were not systematically recorded and  
216 may have contributed to residual confounding.

217

218 **5. CONCLUSION**

---

219 This prospective observational study demonstrates that perioperative respiratory incidents remain  
220 frequent in pediatric anesthesia at a Moroccan referral center, with an incidence of 26.8%. Upper  
221 airway obstruction, multiple intubation attempts, and agitation at induction were identified as the three  
222 independent risk factors.

223 These findings support systematic preoperative airway assessment including active UAO screening,  
224 implementation of structured difficult airway algorithms with appropriate equipment availability, and  
225 targeted anxiolytic premedication strategies as priority measures to improve pediatric anesthetic  
226 safety.

227 Multicenter national studies are needed to confirm these results, identify regional disparities, and  
228 evaluate the impact of educational and organizational interventions on perioperative respiratory safety  
229 in pediatric anesthesia.

230

231 **List of Abbreviations**

232 ASA: American Society of Anesthesiologists; CI: Confidence Interval; IQR: Interquartile Range; OR:  
233 Odds Ratio; PRI: Perioperative Respiratory Incident; SpO<sub>2</sub>: Peripheral Oxygen Saturation; STROBE:  
234 Strengthening the Reporting of Observational Studies in Epidemiology; UAO: Upper Airway  
235 Obstruction.

236 **Ethics Approval and Consent to Participate**

237 This study was conducted in accordance with the ethical principles of the Declaration of Helsinki.  
238 Informed consent was obtained from the parents or legal guardians of each patient prior to inclusion.  
239 Patient data were anonymized prior to analysis.

240 **Funding**

241 This study received no external funding. It was conducted as part of the routine clinical activities of the  
242 Department of Anesthesiology and Critical Care, CHU Ibn Rochd, Casablanca

243 **Conflict of Interest**

244 The author(s) confirm(s) that there is no conflict of interest related to this manuscript.

245

246 **. REFERENCES**

---

247 [1] Bhananker SM, Ramamoorthy C, Geiduschek JM, Posner KL, Domino KB, Haberkern CM, et al. Anesthesia-  
248 related cardiac arrest in children: update from the pediatric perioperative cardiac arrest registry. *Anesth*  
249 *Analg.* 2007;105(2):344–350.

- 250 [2] von Ungern-Sternberg BS, Boda K, Chambers NA, Rebmann C, Johnson C, Sly PD, et al. Risk assessment  
251 for respiratory complications in paediatric anaesthesia: a prospective cohort study. *Lancet*.  
252 2010;376(9743):773–783.
- 253 [3] Mamie C, Habre W, Delhumeau C, Argiroffo CB, Morabia A. Incidence and risk factors of perioperative  
254 respiratory adverse events in children undergoing elective surgery. *Paediatr Anaesth*. 2004;14(3):218–224.
- 255 [4] Habre W, Disma N, Virag K, Habre W, et al.; APRICOT Group of the European Society of Anaesthesiology  
256 Clinical Trial Network. Incidence of severe critical events in paediatric anaesthesia (APRICOT): a prospective  
257 multicentre observational study in 261 hospitals in Europe. *Lancet Respir Med*. 2017;5(5):412–425.
- 258 [5] Murat I, Constant I, Maud’huy H. Perioperative anaesthetic morbidity in children: a database of 24,165  
259 anaesthetics over a 30-month period. *Paediatr Anaesth*. 2004;14(2):158–166.
- 260 [6] Enright A, Sadler LC, Cole JW, Engelhardt T, von Ungern-Sternberg BS. Recent trends in perioperative  
261 serious adverse events (SAE) and associated risk factors for incidents in pediatric anesthesia: a systematic  
262 review. *Paediatr Anaesth*. 2019;29(2):99–113.
- 263 [7] Wudineh DM, Berhe YW, Chekol WB, Adane H, Workie MM. Perioperative respiratory adverse events among  
264 pediatric surgical patients in university hospitals in northwest Ethiopia; a prospective observational study.  
265 *Front Pediatr*. 2022;10:827663.
- 266 [8] Tay CL, Tan GM, Ng SB. Critical incidents in paediatric anaesthesia: an audit of 10,000 anaesthetics in  
267 Singapore. *Paediatr Anaesth*. 2001;11(6):711–718.
- 268 [9] Subramanyam R, Yeramani S, Hossain MM, Anneken AM, Varughese AM. Perioperative respiratory  
269 adverse events in pediatric ambulatory anesthesia: development and validation of a risk prediction tool.  
270 *Anesth Analg*. 2016;122(5):1578–1585.
- 271 [10] Ramgolam A, Hall GL, Zhang G, Hegarty M, von Ungern-Sternberg BS. Inhalational versus intravenous  
272 induction of anesthesia in children with a high risk of perioperative respiratory adverse events: a randomized  
273 controlled trial. *Anesthesiology*. 2018;128(6):1065–1074.
- 274 [11] Parnis SJ, Barker DS, van der Walt JH. Clinical predictors of anaesthetic complications in children with  
275 respiratory tract infections. *Paediatr Anaesth*. 2001;11(1):29–40.
- 276 [12] Mort TC. The incidence and risk factors for cardiac arrest during emergency tracheal intubation: a justification  
277 for incorporating the ASA Guidelines in the remote location. *J Clin Anesth*. 2004;16(7):508–516.
- 278 [13] Peterson GN, Domino KB, Caplan RA, Posner KL, Lee LA, Cheney FW. Management of the difficult airway:  
279 a closed claims analysis. *Anesthesiology*. 2005;103(1):33–39.
- 280 [14] Konrad C, Schüpfer G, Wietlisbach M, Gerber H. Learning manual skills in anesthesiology: is there a  
281 recommended number of cases for anesthetic procedures? *Anesth Analg*. 1998;86(3):635–639.
- 282 [15] Kain ZN, Mayes LC, O’Connor TZ, Cicchetti DV. Preoperative anxiety in children: predictors and outcomes.  
283 *Arch Pediatr Adolesc Med*. 1996;150(12):1238–1245.
- 284 [16] Funk W, Jakob W, Riedl T, Taeger K. Oral preanaesthetic medication for children: double-blind randomized  
285 study of a combination of midazolam and ketamine vs midazolam or ketamine alone. *Br J Anaesth*.  
286 2000;84(3):335–340.
- 287 [17] Kain ZN, Caldwell-Andrews AA, Mayes LC, Weinberg ME, Wang SM, MacLaren JE, et al. Family-centered  
288 preparation for surgery improves perioperative outcomes in children: a randomized controlled trial.  
289 *Anesthesiology*. 2007;106(1):65–74.
- 290 [18] Hordofa TA, Tuna AN, Bussa MB, Waqayo KT, Bariso WT, Ayano GT, et al. Incidence of postoperative  
291 respiratory adverse events and its predictors among pediatric surgical patients at Hawassa University  
292 Comprehensive Specialized Hospital: a prospective follow-up study. *Front Anesth*. 2025;4:1550316. doi:  
293 10.3389/fanes.2025.1550316.
- 294 [19] Disma N, Veyckemans F, Virag K, Hansen TG, Becke K, Harlet P, et al.; NECTARINE Group of the  
295 European Society of Anaesthesiology Clinical Trial Network. Morbidity and mortality after anaesthesia in  
296 early life: results of the European prospective multicentre observational study, neonate and children audit of  
297 anaesthesia practice in Europe (NECTARINE). *Br J Anaesth*. 2021;126(6):1157–1172. doi:  
298 10.1016/j.bja.2021.02.016.
- 299 [20] Luedeke CM, Rudolph MI, Pulverenti TS, Azimaraghi O, Grimm AM, Jackson WM, et al. Development and  
300 validation of a score for prediction of postoperative respiratory complications in infants and children (SPORC-C).  
301 *Br J Anaesth*. 2025;134(1):212–220. doi: 10.1016/j.bja.2024.07.011.
- 302 [21] Deng QW, Tan WC, Zhan YQ, Wang XW, Lai HJ, Wen SH. Prophylactic pharmacological interventions  
303 against perioperative respiratory adverse events in children undergoing noncardiac surgery: a systematic  
304 review and meta-analysis. *J Anesth*. 2025;39(2):248–264. doi: 10.1007/s00540-024-03453-y.
- 305
- 306