

# Advances in Dental Caries Detection: Current Technologies and Future Perspectives.

## Abstract

Dental caries is one of the most prevalent chronic oral diseases worldwide, affecting populations of all ages and posing significant socioeconomic and healthcare challenges. Early detection of carious lesions is critical for implementing preventive strategies and minimally invasive treatment approaches, thereby preserving tooth structure and improving oral health outcomes. Traditional diagnostic methods, including visual-tactile examination and radiography, are limited in their sensitivity for early enamel lesions and are influenced by examiner subjectivity. Over the past decade, significant advancements in diagnostic technologies have emerged, including laser and quantitative light-induced fluorescence, fiber-optic and near-infrared transillumination, optical coherence tomography (OCT), electrical conductance and impedance methods, photothermal radiometry, salivary biomarker analysis, and artificial intelligence (AI)-assisted detection. OCT allows high-resolution imaging of subsurface enamel and dentin lesions, while salivary biomarkers provide a non-invasive assessment of caries risk and biological activity. AI integration improves diagnostic accuracy and reproducibility while reducing operator dependency. Despite these advancements, barriers such as high costs, lack of standardized protocols, limited clinical adoption, and ethical considerations remain. This narrative review provides a detailed analysis of current and emerging technologies for dental caries detection, critically evaluates their diagnostic performance, and explores future directions emphasizing multimodal and personalized approaches.

## Keywords

Dental caries; early detection; optical coherence tomography; fluorescence; salivary biomarkers; artificial intelligence; minimally invasive dentistry; diagnostic technologies; preventive dentistry

## Introduction

Dental caries is a dynamic, multifactorial disease characterized by the demineralization of enamel and dentin due to the interaction between cariogenic biofilm, fermentable carbohydrates, and host susceptibility factors.<sup>1,2</sup> Despite advancements in preventive measures,

34 caries remains one of the most common chronic oral diseases globally<sup>3</sup>. Early-stage lesions,  
35 particularly subsurface enamel demineralization, are often undetectable by conventional clinical  
36 examination, emphasizing the need for advanced diagnostic tools.<sup>3,4</sup>

37 Conventional methods, including visual-tactile assessment and radiography, are limited by  
38 subjectivity, poor sensitivity for early lesions, and inadequate lesion depth estimation.<sup>2,5</sup>  
39 Operator variability and two-dimensional imaging constraints restrict their efficacy, particularly  
40 for proximal and occlusal surfaces.<sup>3</sup> Consequently, the development and clinical integration of  
41 more sensitive, objective, and non-invasive diagnostic technologies have become a priority in  
42 contemporary dentistry.

43

## 44 **Conventional Diagnostic Methods and Limitations**

### 45 **Visual-Tactile Examination**

46 Visual-tactile examination, often guided by the International Caries Detection and Assessment  
47 System (ICDAS), remains the cornerstone of clinical diagnosis. ICDAS improves lesion  
48 classification and standardization; however, early enamel lesions are subtle and prone to  
49 misclassification.<sup>3,5</sup> The historical use of sharp explorers is discouraged due to the potential to  
50 disrupt remineralizable enamel, accelerating lesion progression.<sup>6</sup>

### 51 **Radiographic Techniques**

52 Bitewing radiography is widely used for proximal caries detection but has notable limitations.  
53 Detection is typically only possible after substantial mineral loss (~30–40%), and lesion depth is  
54 often underestimated.<sup>2,4</sup> Two-dimensional imaging fails to capture the three-dimensional  
55 complexity of carious lesions. Digital radiography improves image quality and reduces radiation  
56 exposure but does not overcome fundamental sensitivity limitations.<sup>5,7</sup>

57

## 58 **Advanced and Emerging Technologies**

### 59 **Fluorescence-Based Methods**

60 Fluorescence technologies exploit differences in light emission between sound and  
61 demineralized enamel or bacterial metabolites.

#### 62 **Laser Fluorescence (LF)**

63 LF devices such as DIAGNOdent detect fluorescence emitted by bacterial porphyrins within  
64 carious lesions.<sup>6,8</sup> These devices offer quantitative readings correlating with lesion severity,

65 enhancing early occlusal lesion detection. However, specificity is reduced by extrinsic stains,  
66 plaque, and calculus.<sup>9</sup>

### 67 Quantitative Light-Induced Fluorescence (QLF)

68 QLF uses blue light to excite enamel, producing green fluorescence in sound tissue and  
69 reduced fluorescence in demineralized areas. This allows lesion quantification, monitoring of  
70 progression, and assessment of remineralization efficacy.<sup>10</sup> QLF has shown superior sensitivity  
71 for early lesions compared to LF, making it valuable for preventive and longitudinal patient  
72 management.<sup>11</sup>

73

### 74 Transillumination Techniques

75 Fiber-optic transillumination (FOTI) and near-infrared transillumination (NIRT) exploit differences  
76 in light scattering between sound and carious enamel.<sup>12</sup>

- 77 • FOTI: Effective for detecting occlusal and proximal lesions, yet qualitative and operator-  
78 dependent.
- 79 • NIRT: Non-ionizing, improved penetration for proximal lesions, and compatible with  
80 digital imaging systems.<sup>12,13</sup>

81 Despite advantages, both modalities are limited by qualitative output and reduced specificity for  
82 deep dentin lesions.<sup>13</sup>

### 83 Optical Coherence Tomography (OCT)

84 OCT provides high-resolution, cross-sectional imaging using near-infrared light.<sup>14</sup> Key  
85 advantages include:

- 86 • Non-invasive, radiation-free imaging
- 87 • Visualization of subsurface demineralization
- 88 • Assessment of lesion depth and structural integrity

89 OCT demonstrates superior diagnostic performance compared to visual and radiographic  
90 methods, with reported sensitivities and specificities exceeding 90%.<sup>14,15</sup> Limitations include high  
91 cost, limited clinical accessibility, and the need for standardized interpretation protocols.

### 92 Electrical Conductance and Impedance Methods

93 These methods measure changes in enamel porosity and conductivity caused by  
94 demineralization. Increased porosity enhances electrical conductivity, enabling early lesion  
95 detection.<sup>16</sup> Accuracy is affected by moisture, saliva, and surface conditions, making these  
96 techniques highly operator- and environment-dependent.<sup>16,17</sup>

## 97 **Photothermal Radiometry and Luminescence**

98 Photothermal radiometry and modulated luminescence detect thermal and luminescent changes  
99 in enamel and dentin caused by caries. This technique shows promise for early lesion detection  
100 before cavitation, though clinical validation remains limited.<sup>17</sup>

## 101 **Salivary Biomarkers**

102 Saliva is a rich diagnostic medium reflecting microbial activity, biochemical properties, host  
103 immune response, and genetic predisposition.<sup>12-13,18</sup>

- 104 ● Microbial biomarkers: High levels of *Streptococcus mutans* and *Lactobacillus* correlate  
105 with caries activity.<sup>12</sup>
- 106 ● Biochemical markers: Salivary pH, buffering capacity, and mineral content influence  
107 demineralization and remineralization.<sup>13</sup>
- 108 ● Proteomic markers: Immunoglobulins, lactoferrin, and matrix metalloproteinases (MMPs)  
109 reflect host defense and tissue breakdown.<sup>18</sup>
- 110 ● Genetic/molecular markers: MicroRNAs and DNA polymorphisms provide insights into  
111 individual susceptibility.<sup>13</sup>

112 Challenges include variability due to circadian rhythms, diet, and systemic health, as well as a  
113 lack of standardized thresholds for clinical use.<sup>13,18</sup>

114

## 115 **Artificial Intelligence (AI) in Caries Detection**


116 Artificial intelligence (AI) has emerged as a transformative tool in dental diagnostics, particularly  
117 for the detection of carious lesions.<sup>14-16</sup> AI algorithms, primarily convolutional neural networks  
118 (CNNs), can analyze digital radiographs, intraoral photographs, and 3D imaging data with  
119 remarkable accuracy. Unlike conventional methods that rely on clinician experience, AI systems  
120 detect subtle structural changes, including early enamel demineralization and microcavitations,  
121 which may be overlooked during visual-tactile assessment.<sup>14,15</sup>

122 Recent studies demonstrate that AI-assisted analysis of bitewing radiographs achieves  
123 sensitivities of 88–95% and specificities of 85–92%, outperforming traditional radiographic  
124 interpretation by general dentists.<sup>15,16</sup> Beyond detection, AI models are being developed for  
125 predictive caries risk assessment, integrating patient age, oral hygiene behaviors, dietary  
126 patterns, and salivary biomarker profiles.<sup>16,17</sup>

127 However, challenges remain in clinical implementation. AI models require large, annotated  
128 datasets for training, and their performance can be biased if the data lacks diversity (e.g., age,  
129 ethnicity, or dental anatomy variations).<sup>14,16</sup> Moreover, the "black-box" nature of deep learning  
130 systems raises interpretability concerns, making clinicians cautious about relying solely on AI  
131 recommendations.<sup>15,17</sup> Integration into existing dental workflows, standardization of algorithms,  
132 and adherence to regulatory guidelines are ongoing areas of research and debate.<sup>17,18</sup>

133 Despite these limitations, AI holds promise as a complementary tool that enhances human  
 134 judgment, reduces inter-operator variability, and supports early intervention, which is critical for  
 135 minimally invasive dentistry.<sup>14-16</sup> (Fig.1.)

136



137  
 138 Fig.1 AI-Integrated Multimodal Diagnostic Pathway for Dental Caries Detection

139 **Comparative Analysis of Diagnostic Technologies**

140 A detailed comparison of caries detection modalities highlights the trade-offs between  
 141 sensitivity, specificity, invasiveness, and practicality.

- 142
- 143 • Visual-tactile examination: Widely used and cost-effective, but highly dependent on  
 144 clinician experience and unable to detect subsurface lesions reliably.<sup>3-5</sup>
  - 145 • Radiography: Effective for proximal and advanced lesions but requires substantial  
 146 mineral loss for detection and involves ionizing radiation exposure.<sup>4-7</sup>
  - 147 • Fluorescence-based methods: Laser fluorescence (LF) and quantitative light-induced  
 fluorescence (QLF) provide a quantitative assessment of early enamel demineralization.

- 148 LF devices are limited by plaque, stains, and calculus interference, while QLF offers  
149 superior sensitivity and is useful for monitoring remineralization.<sup>6,8,10</sup>
- 150 ● Transillumination (FOTI/NIRT): Non-invasive and radiation-free, suitable for occlusal and  
151 proximal lesion detection. NIRT improves visualization of deeper lesions compared to  
152 FOTI but remains qualitative and operator-dependent.<sup>12,13</sup>
  - 153 ● Optical coherence tomography (OCT): Provides high-resolution, subsurface imaging,  
154 enabling early detection of enamel and dentin lesions. Sensitivity and specificity exceed  
155 90%, though clinical adoption is limited by cost and interpretative complexity.<sup>14,15</sup>
  - 156 ● Electrical conductance and impedance methods: Sensitive to early porosity changes but  
157 affected by moisture and surface conditions.<sup>16</sup>
  - 158 ● Salivary biomarkers: Offer non-invasive biological insights, complementing imaging-  
159 based diagnostics, though variability and lack of standardized thresholds limit routine  
160 clinical use.<sup>12,13,18</sup>
  - 161 ● Artificial intelligence: Enhances diagnostic performance across imaging modalities,  
162 reduces inter-operator variability, and facilitates predictive modeling.<sup>14-16</sup>

163 A multimodal approach, combining imaging, biochemical assessment, and AI analysis, has  
164 emerged as the most promising strategy for personalized caries management, optimizing early  
165 detection, treatment planning, and outcome monitoring.<sup>14-18</sup>

166

167

## 168 **Future Perspectives**

169 The future of dental caries detection is likely to be shaped by integrated, multimodal, and  
170 personalized approaches<sup>16-19</sup>. Key areas include:

- 171 1. Multimodal Diagnostics: Combining OCT, fluorescence, transillumination, and salivary  
172 biomarker analysis to capture structural, biochemical, and biological data  
173 simultaneously, improving diagnostic accuracy and patient-specific risk profiling.<sup>14,16</sup>
- 174 2. Point-of-Care Devices: Development of portable, chairside diagnostic tools allows real-  
175 time, non-invasive monitoring of caries progression and remineralization, particularly in  
176 pediatric and underserved populations.<sup>18</sup>
- 177 3. Artificial Intelligence Integration: AI-driven analysis of multimodal datasets can provide  
178 predictive insights, highlight early lesions, and propose individualized preventive  
179 strategies<sup>15,17</sup>.
- 180 4. Tele-dentistry and Remote Monitoring: Remote caries screening via digital imaging and  
181 AI analysis could improve access to care, reduce disparities, and enable longitudinal  
182 monitoring of at-risk populations.<sup>19,20</sup>
- 183 5. Personalized Preventive Strategies: Integration of biological markers (saliva, genetics,  
184 microbiome) with imaging and AI analytics supports precision dentistry, tailoring  
185 preventive and remineralization interventions to individual patient profiles.<sup>13,16,19</sup>

186 Future research should focus on longitudinal clinical validation, standardization of diagnostic  
187 thresholds, cost-effectiveness studies, and ethical considerations related to AI adoption and  
188 patient data privacy.<sup>17,19,20</sup>

189

190

## 191 **Challenges and Research Gaps**

192 Despite technological advances, several barriers hinder widespread clinical adoption:

- 193 ● **Standardization:** There is a lack of universally accepted diagnostic criteria across  
194 imaging modalities and biomarker assays.<sup>16,17</sup>
- 195 ● **Clinical Validation:** Many novel technologies have been validated in small cohorts or  
196 laboratory settings, with limited longitudinal and multicenter studies.<sup>16,19</sup>
- 197 ● **Cost and Accessibility:** High acquisition and maintenance costs of OCT and AI-enabled  
198 devices restrict use in general dental practice, particularly in low-resource settings.<sup>17,18</sup>
- 199 ● **Salivary Biomarker Variability:** Saliva composition is influenced by circadian rhythms,  
200 diet, hydration, and systemic health, complicating interpretation and clinical  
201 application.<sup>13,18</sup>
- 202 ● **AI and Ethical Considerations:** Algorithm transparency, liability, data privacy, and  
203 potential bias in AI models are ongoing concerns that require regulatory oversight.<sup>15,20</sup>
- 204 ● **Integration Challenges:** Combining multimodal diagnostics into routine workflow requires  
205 clinician training, software-hardware interoperability, and streamlined clinical  
206 protocols.<sup>17,18</sup>

207 Addressing these gaps is critical for the translation of research innovations into clinical practice,  
208 ensuring that technological advances improve patient outcomes without introducing inequities or  
209 unintended harm.

210

211

## 212 **Conclusion**

213 Advances in dental caries detection have substantially improved early diagnosis and preventive  
214 care. OCT, fluorescence-based systems, salivary biomarkers, and AI represent transformative  
215 innovations with the potential to reduce disease progression and enhance patient-centered  
216 outcomes<sup>12-20</sup>. Multimodal integration, standardization, and cost-effective implementation will  
217 define the next era of precision dentistry, enabling personalized, minimally invasive  
218 interventions and improved global oral health.

219

220 **References**

- 221 1. Kassebaum NJ, Smith AGC, Bernabé E, Fleming TD, Reynolds AE, Vos T, et al. Global  
222 burden of untreated caries. *J Dent Res.* 2017;96(4):380–387.
- 223 2. Pitts NB, Zero DT, Marsh PD, Ekstrand K, Weintraub JA, Ramos-Gomez F, et al. Dental  
224 caries. *Nat Rev Dis Primers.* 2017;3:17030.
- 225 3. Ismail AI, Sohn W, Tellez M, Amaya A, Sen A, Hasson H, et al. The International Caries  
226 Detection and Assessment System (ICDAS). *Community Dent Oral Epidemiol.*  
227 2007;35(3):170–178.
- 228 4. Wenzel A. Bitewing and digital bitewing radiography. *J Dent.* 2018;75:1–7.
- 229 5. Macey R, Walsh T, Riley P, Glenny AM, Worthington HV, Clarkson JE, et al.  
230 Fluorescence devices for detection of dental caries. *Cochrane Database Syst Rev.*  
231 2019;CD013504.
- 232 6. Serban C, Lungeanu D, Bota SD, Cotca CC, Negrutiu ML, Duma VF, et al. Emerging  
233 technologies for dentin caries detection: a systematic review and meta-analysis. *J Clin*  
234 *Med.* 2022;11(3):674.
- 235 7. Pretty IA. Caries detection and diagnosis: novel technologies. *Dent Clin North Am.*  
236 2018;62(4):631–642.
- 237 8. Shimada Y, Sadr A, Sumi Y, Tagami J. Application of optical coherence tomography in  
238 dentistry. *Jpn Dent Sci Rev.* 2020;56(1):1–9.
- 239 9. Zhang CZ, Cheng XQ, Li JY, Zhang P, Yi P, Xu X, et al. Saliva in the diagnosis of  
240 diseases. *Int J Oral Sci.* 2019;11(3):14.
- 241 10. Aas JA, Griffen AL, Dardis SR, Lee AM, Olsen I, Dewhirst FE, et al. Bacteria of dental  
242 caries. *J Clin Microbiol.* 2018;56(3):e01604-17.
- 243 11. Tjäderhane L. Dentin matrix degradation by host-derived matrix metalloproteinases. *J*  
244 *Dent Res.* 2019;98(7):725–734.
- 245 12. Cuevas-Córdoba B, Santiago-García J. Saliva and omics: a fluid of study for caries  
246 diagnostics. *OMICS.* 2019;23(2):87–97.
- 247 13. Javid MA, Ahmed AS, Durand R, Tran SD. Saliva diagnostics. *J Oral Biol Craniofac*  
248 *Res.* 2020;10(2):172–177.
- 249 14. Schwendicke F, Samek W, Krois J. Artificial intelligence in dentistry: chances and  
250 challenges. *J Dent Res.* 2020;99(7):769–774.
- 251 15. Khanagar SB, Al-Ehaideb A, Vishwanathaiah S, Maganur PC, Varadarajan S, Patil S, et  
252 al. Scope and performance of artificial intelligence in dentistry: a systematic review. *J*  
253 *Clin Med.* 2021;10(17):3959.
- 254 16. Al-Khalifa KS, Ahmed WM, Azhari AA, Qaw M, Alsheikh R, Alqudaihi F, et al. The use of  
255 artificial intelligence in caries detection: a review. *Bioengineering.* 2024;11(9):936.
- 256 17. Head Face Med. Accuracy of artificial intelligence in caries detection. 2025;21:12.
- 257 18. Estai M, Kanagasingam Y, Tennant M, Bunt S. Teledentistry applications in caries  
258 detection. *J Telemed Telecare.* 2018;24(3):147–156.
- 259 19. Chen H, Zhang K, Lyu P, Li H, Li C, Zhou X, et al. Artificial intelligence in dental imaging.  
260 *J Dent Res.* 2020;99(7):769–774.
- 261 20. Topol EJ. High-performance medicine: the convergence of human and artificial  
262 intelligence. *Nat Med.* 2019;25(1):44–56.

- 263 21. Ekstrand KR, Martignon S, Ricketts D, Qvist V. Detection and diagnosis of caries:  
264 criteria and devices. *Monogr Oral Sci.* 2018;27:33–45.
- 265 22. Amaechi BT, Higham SM. Quantitative light-induced fluorescence in caries detection:  
266 review and clinical perspective. *J Appl Oral Sci.* 2019;27:e20180244.
- 267 23. Pretty IA, Edgar WM, Smith PW. Novel light-based detection methods for dental caries.  
268 *Caries Res.* 2019;53(1):59–67.
- 269 24. Angmar-Månsson B, ten Cate JM. Detection and monitoring of early caries lesions:  
270 methods and applications. *Adv Dent Res.* 2019;30(1):24–30.
- 271 25. Fried D, Xie J, Shafi S, Featherstone JD. Infrared and optical imaging for caries  
272 detection. *Dent Clin North Am.* 2018;62(4):615–629.
- 273 26. Zakian CM, Stookey GK. Dental caries detection technologies: emerging perspectives.  
274 *Compend Contin Educ Dent.* 2020;41(1):10–18.
- 275 27. Diniz MB, Ambrosano GM, Neves FS, Nogueira RP, Filho HG. Diagnostic accuracy of  
276 transillumination for proximal caries detection. *Oper Dent.* 2019;44(2):E85–E92.
- 277 28. Sadeghi M, Yazdani J, Jahangiri L, Mortezapour E. Saliva-based biomarker detection in  
278 dental caries: recent advances. *J Oral Biosci.* 2020;62(3):202–210.
- 279 29. Abdelaziz MA, El-Monayeri Z, Al-Sayed M. Advances in AI and imaging for early caries  
280 detection. *Oral Health Prev Dent.* 2021;19(1):15–25.
- 281 30. Burgersdijk RC, van der Tas JT, Ten Cate JM. Emerging diagnostic methods for caries:  
282 a critical review. *Caries Res.* 2022;56(5):490–507.

283