

Primary Squamous Cell Carcinoma of the Breast: A Diagnostic Enigma.

Abstract

Primary squamous cell carcinoma of the breast is an uncommon malignancy and represents a very small proportion of breast cancers. Because of its rarity and non-specific clinical presentation, diagnosis is challenging and often requires correlation of imaging, cytology, and histopathology. Pure squamous cell carcinoma of the breast can deviate from epidermis, the nipple or the epithelium of deep-seated epidermoid cyst or squamous metaplasia on chronic inflammatory background. We report the case of a 45-year-old female who presented with a painful lump in the left breast for three years. Radiological imaging revealed a predominantly cystic lesion and initial cytology suggested fibrocystic disease. However, histopathological and IHC examination of the lesion demonstrated features of primary squamous cell carcinoma. This case emphasizes the importance of histopathological confirmation in breast lesions that clinically and radiologically resemble benign conditions.


Keywords : PSCC (Primary squamous cell carcinoma), BIRADS, HER-2

Introduction

Squamous cell carcinoma (SCC) of the breast is an extremely rare malignant neoplasm, accounting for less than 0.1% (approximately 0.04–0.075%) of all invasive breast cancers[1,2]. The diagnosis of primary SCC of the breast is based on strict criteria, which includes: (1) more than 90% of the malignant cells being of squamous cell origin, (2) the tumor being independent of the overlying skin and nipple, and (3) exclusion of a primary squamous cell carcinoma at any other site in the body. [14] Clinically and radiologically, these tumors have nonspecific findings and may mimic benign or inflammatory breast conditions. They are commonly hormone receptor and HER2 negative[3], limiting treatment options and contributing to a poor prognosis. Due to its rarity, diagnostic challenges, and aggressive behavior, we are presenting this case to contribute to the existing literature and improve understanding of this uncommon entity.

Case Presentation

A 45-year-old female presented with a history of a lump in the left breast since approximately three years which gradually increased in size over time and was associated with pain. There was no history of nipple discharge, bleeding from the nipple, ulceration of the skin, or trauma to the breast. The patient did not report any significant past medical illness or family history of breast malignancy. On clinical examination, a palpable mass was identified in the upper inner quadrant of the left breast. The swelling was localized to the breast tissue and the overlying skin appeared normal without signs of ulceration or inflammatory changes. The nipple-areolar complex appeared normal. Radiological evaluation of the left breast revealed a large lobulated cystic lesion measuring approximately $5.1 \times 4.8 \times 3.5$ cm which was wider than tall with no internal calcifications. The lesion was reported as complex cystic lesion of the breast with BIRADS score of III. (Fig 1)



36
37

Fig1 Cystic lesion




38
39

Fig 2 Cystic cavity with partial septations

40 On further evaluation by fine needle aspiration, approximately 20 ml of pale yellow fluid was aspirated.
41 Microscopic examination showed numerous foamy macrophages, few ductal epithelial cells along with
42 many neutrophils in a background of red blood cells. No atypical cell was seen. Cytological features
43 suggested Benign cystic lesion consistent with fibrocystic change.


44 Left breast lumpectomy was done and we received a globular mass measuring 5.6 x 4 x 3 cm, on cut
45 section a cystic cavity was identified with partial septations measuring 4.6 x 3 cm (Fig 2). No lymph
46 nodes were retrieved. On microscopic examination a cystic lesion was identified, which was partially
47 lined by atypical squamous epithelium. In addition, several irregular nests and islands of tumour cells
48 were seen infiltrating the surrounding breast stroma. The tumour cells exhibited moderate nuclear
49 pleomorphism, hyperchromatic nuclei with abundant eosinophilic cytoplasm, prominent keratin pearl
50 formation and individual cell keratinization. Features were consistent with squamous differentiation. The
51 adjacent breast tissue revealed duct ectasia accompanied by dense chronic inflammatory cell infiltrate,
52 and preserved normal terminal ductal units. No atypical glandular elements were identified in the sections
53 submitted.(Fig 3).

54 IHC for CK5/6, Pan-Cytokeratin and p40 was done and the case was diagnosed as primary squamous
55 cell carcinoma (PSCC).



56
57

Fig 3 Cavity lined by atypical squamous cells with island of malignant cell in surrounding area.



58
59

Fig 4 Squamous cell carcinoma with normal ductal lobular unit (H&E 100x)

60 Discussion

61 Primary squamous cell carcinoma of the breast is a rare entity. It is termed “pure” when the tumor is
62 composed entirely of malignant squamous cells with no connection with the overlying skin or nipple, and
63 when a primary tumor at another site has been excluded [8,9]. It is important to differentiate it from mixed
64 tumors with squamous components and from metastatic squamous cell carcinoma arising elsewhere.
65 The exact etiology and pathogenesis of squamous cell carcinoma of the breast remain unclear. Malignant
66 transformation following squamous metaplasia in benign conditions such as cysts, chronic inflammation,
67 abscesses, and fibroadenomas has been proposed. This is supported by reports of SCC developing
68 following long-standing benign conditions, breast implants, or prior radiation therapy.[9,1,11] In our case
69 also, the patient had a multiloculated cystic lesion for 3 years, which may support this proposed
70 pathway. The age of our case was 45 years which is similar to the median age of patients presenting with
71 squamous cell carcinoma that is 52 years. It is reported that squamous cell carcinoma occurs in slightly
72 younger females than that of infiltrating duct carcinoma. [12] There are no typical findings on the

73 mammogram, ultrasound. In this case USG had reported it as complicated cystic lesion. Cystic changes
74 maybe because of central degeneration or associated necrosis. As per literature they are known to mimic
75 benign conditions radiologically[13]. Approximately 70% of patients with SCC of the breast have no
76 axillary node involvement in axillary dissection.[1] Menes et al.[3] reported that they are associated with a
77 lower rate (22%) of lymph node metastasis and a significant rate of distant metastasis without lymph node
78 involvement, in this study axillary lymph nodes were not involved. Suresh et al [14]. reported malignant
79 cells on FNAC but in our case FNAC showed features consistent with benign cystic lesion. Squamous cell
80 carcinomas are generally hormonereceptor-negative [8,9,1,10,5]similar to our study. It is recommended to
81 give patients adjuvant therapy but the radiosensitivity of squamous cell carcinomas is uncertain. The 5-year
82 survival is 67% [1].

83 **Conclusion**

84 Primary squamous cell carcinoma of the breast is a rare malignant tumor that may clinically,
85 radiologically and on cytology resemble benign breast lesions. This case illustrates the importance of
86 thorough clinical workup, histopathological evaluation in patients presenting with long-standing breast
87 lumps, abscess and cyst particularly when imaging and cytology suggest benign conditions.
88 Accurate diagnosis is essential to guide appropriate management. The prognosis of this type of breast
89 cancer is still regarded as somewhat controversial, though many studies suggest that it is an aggressive
90 disease that may behave like poorly differentiated breast carcinoma [7]

91 **References**

- 92 1. Behranwala KA, Nasiri N, Abdullah N, Trott PA, Gui GP. Squamous cell carcinoma of the breast: clinicopathologic implications and
93 outcome. *Eur J Surg Oncol.* 2003;29:386–389.
- 94 2. Aparicio I, Martínez A, Hernández G, Hardisson D, De Santiago J. Squamous cell carcinoma of the breast. *Eur J Obstet Gynecol Reprod*
95 *Biol.* 2008;137:222–226.
- 96 3. Menes T, Schachter J, Morgenstern S, Fenig E, Lurie H, Gutman H, et al. Primary squamous cell carcinoma (SqCC) of the breast. *Am J*
97 *Clin Oncol.* 2003;26:571–573.
- 98 4. Karoui A, Chaffai O, Ghorbel A, et al. Primary squamous cell carcinoma of the breast: a case report. *Indian J Gynecol Oncol.*
99 2022;20(3):31.
- 100 5. Gupta G, Malani AK, Weigand RT, Rangenini G. Pure primary squamous cell carcinoma of the breast: a rare presentation and
101 clinicopathologic comparison with usual ductal carcinoma of the breast. *Pathol Res Pract.* 2006;202(6):465–469.
- 102 6. Damin AP, Nascimento FC, Andreola JB, Cerutti TH, Roehe A, Damin DC. Primary epidermoid carcinoma of the breast presenting as a
103 breast abscess and sepsis. *Sao Paulo Med J.* 2011;129(6):424–427.
- 104 7. Moisisidis E, Ahmed S, Carmalt H, Gillett D. Primary squamous cell carcinoma of the breast. *ANZ J Surg.* 2002;72:65–67.
- 105 8. Siegelmann-Danieli N, Murphy TJ, Meschter SC, et al. Primary pure squamous cell carcinoma of the breast. *Clin Breast Cancer.*
106 2005;3:270–272.
- 107 9. Zoltan TB, Konick L, Coleman RJ. Pure squamous cell carcinoma of the breast in a patient with previous adenocarcinoma of the breast: a
108 case report and review of the literature. *Am Surg.* 2001;67(7):671–673.
- 109 10. Wrightson WR, Edwards MJ, McMasters KM. Primary squamous cell carcinoma of the breast presenting as a breast abscess. *Am Surg.*
110 1999;65(12):1153–1155.
- 111 11. Singh H, Williams SP, Kinsella V, Lynch GR. Postradiation squamous cell cancer of the breast. *Cancer Invest.* 2000;18:343–346.
- 112 12. Weigel RJ, Ikeda DM, Nowels KW. Primary squamous cell carcinoma of the breast. *South Med J.* 1996;89:511–515.
- 113 13. Flikweert ER, Hofstee M, Liem MSL. Squamous cell carcinoma of the breast: a case report. *World J Surg Oncol.* 2008;6:135.
- 114 14. Bhosale SJ, Kshirsagar AY, Deshmukh SJ, Jagtap SV, Langade
115 YB. Squamous cell carcinoma of the breast. *Am J Case Rep.* 2013;14:188.