

1 **A FATAL PERIOPERATIVE STORM: SYNERGISTIC STATIN–**
2 **HALOGENATED MYOTOXICITY WITH REFRACTORY**
3 **HYPERKALAEMIA AND ACUTE KIDNEY INJURY — A CASE**
4 **REPORT AND PROPOSED CLINICAL REASONING FRAMEWORK.**

5 **ABSTRACT**

6 **Background:** Statins and halogenated volatile anaesthetics are independently myotoxic, but their
7 concurrent perioperative interaction remains poorly characterised. Current guidelines recommend
8 perioperative statin continuation in coronary patients, yet the resulting pharmacological intersection
9 with inhalational anaesthesia has not been integrated into routine risk stratification.

10 **Case Presentation:** A 66-year-old woman receiving long-term high-intensity atorvastatin (40
11 mg/day), with poorly controlled type 2 diabetes (HbA1c 9.0%), recent inferior ST-elevation
12 myocardial infarction with drug-eluting stent, and ischaemic cardiomyopathy (LVEF 35%),
13 underwent uneventful 180-minute open reduction–internal fixation of an AO/OTA 13-C2 distal
14 humeral fracture under sevoflurane in left lateral decubitus. Twenty-two hours postoperatively, after
15 an asymptomatic interval, she abruptly developed circulatory collapse, diffuse myalgia, anuria and
16 obtundation. Investigations revealed massive rhabdomyolysis (creatinine phosphokinase 31,010 IU/L),
17 severe hyperkalaemia (8.0 mmol/L), KDIGO stage 3 acute kidney injury, severe metabolic acidosis
18 (pH 7.20), and high-sensitivity troponin I 13,000 pg/mL with unchanged left-ventricular function.
19 Despite intensive multimodal therapy and emergent intermittent haemodialysis, rebound
20 hyperkalaemia and refractory cardiac arrest occurred 35 hours after surgery.

21 **Conclusion:** This case integrates six convergent injurious pathways — chronic high-intensity statin
22 therapy, halogenated anaesthetic exposure, uncontrolled diabetes, advanced ischaemic
23 cardiomyopathy, prolonged lateral decubitus and hyperkalaemia-promoting polypharmacy — to
24 explain a fulminant fatal outcome. We propose a structured perioperative reasoning framework (the
25 STAT-OP concept) and argue for individualised consideration of total intravenous anaesthesia and
26 continuous high cut-off renal replacement therapy in highest-risk statin-treated patients.

27 **Keywords:** Anaesthetics, inhalation; Atorvastatin; Hydroxymethylglutaryl-CoA Reductase Inhibitors;
28 Hyperkalaemia; Perioperative Care; Rhabdomyolysis.

29 **INTRODUCTION**

30 Rhabdomyolysis is a syndromic disintegration of skeletal muscle that releases intracellular
31 constituents into the systemic circulation, with creatine phosphokinase (CPK) values exceeding five
32 times the upper limit of normal as its diagnostic biological cornerstone, and acute kidney injury
33 (AKI), life-threatening hyperkalaemia, severe metabolic acidosis and cardiac arrest as its principal
34 lethal complications.¹ In the perioperative setting, two pharmacological classes deserve particular
35 attention. Statins, the cornerstone of secondary cardiovascular prevention, exert dose-dependent
36 myotoxicity through inhibition of HMG-CoA reductase, depletion of mevalonate-derived isoprenoids
37 that anchor small GTPases (Rab, Rho), reduction of mitochondrial coenzyme Q10 and consecutive
38 impairment of cellular bioenergetics and sarcolemmal integrity.^{2,3} Halogenated volatile anaesthetics
39 — sevoflurane, isoflurane, desflurane — activate the type 1 ryanodine receptor (RyR1) on the

40 sarcoplasmic reticulum, producing cytosolic Ca²⁺ overload that, in genetically predisposed patients,
41 may manifest as classical malignant hyperthermia (MH) or as atypical, subclinical, anaesthetic-
42 induced rhabdomyolysis (AIR) without overt hyperthermia or rigidity.^{4,5}

43 A 2021 meta-analysis of 14 studies confirmed that the *SLCO1B1* c.521T>C (rs4149056)
44 polymorphism, which reduces hepatic OATP1B1-mediated atorvastatin uptake, is robustly associated
45 with statin-induced myopathy (pooled OR 2.9 [95% CI 1.59–5.34] for CC homozygotes; OR 4.0
46 [1.23–12.63] for atorvastatin users with CC genotype).⁶ Concurrently, *RYR1* and *CACNA1S* variants
47 underlie the entire continuum from clinical MH to delayed AIR.^{5,7} A perioperative pharmacological
48 intersection between these two pathways is therefore biologically inevitable in a substantial subset of
49 statin-treated surgical patients, but remains unaccounted for in current risk-stratification practice. The
50 2024 AHA/ACC perioperative guideline for noncardiac surgery recommends continuation of
51 established statin therapy in coronary patients,⁸ consistent with meta-analytic evidence of reduced
52 myocardial infarction (OR 0.44 [0.30–0.64]) and 30-day mortality with perioperative statin
53 continuation.⁹ Yet a recent case of severe rhabdomyolysis in a statin user undergoing
54 parathyroidectomy under general anaesthesia explicitly raised the hypothesis of a synergistic statin–
55 anaesthetic toxicity.¹⁰

56 We report a fatal postoperative rhabdomyolysis in which the convergence of chronic high-intensity
57 statin therapy, sevoflurane-based anaesthesia and a heavy comorbidity load produced a fulminant,
58 refractory and lethal trajectory. Beyond the descriptive interest of the case, we propose an integrative
59 pathophysiological model and a structured perioperative reasoning framework (the **STAT-OP**
60 concept — *Statin–Anaesthetic Toxicity in the Operative Period*) intended to support individualised
61 risk stratification in routine practice.

62 **CASE REPORT**

63 A 66-year-old woman (60 kg, 162 cm, body mass index 22.9 kg/m²) was admitted following a
64 domestic fall with a closed comminuted distal humeral palette fracture (AO/OTA 13-C2). Her
65 medical history included a 15-year type 2 diabetes mellitus with poor control (glycated haemoglobin
66 9.0%), persistent moderate asthma, and a recent inferior ST-elevation myocardial infarction (two
67 months earlier) treated by drug-eluting stent in the right coronary artery, complicated by ischaemic
68 dilated cardiomyopathy (left ventricular ejection fraction [LVEF] 35%; hypokinesia of the inferior,
69 basal-to-mid lateral and basal septal segments).

70 Chronic medications comprised **atorvastatin 40 mg/day**, dual antiplatelet therapy (clopidogrel 75
71 mg, aspirin 75 mg), sacubitril/valsartan 49/51 mg twice daily, spironolactone 25 mg/day, furosemide
72 40 mg/day, bisoprolol 2.5 mg/day, empagliflozin 10 mg/day, and inhaled bronchodilators.

73 Preanaesthetic evaluation classified her as American Society of Anesthesiologists (ASA) physical
74 status III. Baseline laboratory values were within normal limits, including serum creatinine (56.3
75 µmol/L), potassium (4.3 mmol/L), and complete blood count. According to current consensus,
76 clopidogrel was withheld for 5 days and empagliflozin for 3 days; atorvastatin and other
77 cardiovascular drugs were continued in line with the 2024 AHA/ACC perioperative guideline.⁸
78 Surgery was performed on the fifth day after admission.

79 After standard monitoring, anaesthesia was induced with intravenous fentanyl, propofol, lidocaine and
80 rocuronium, and **maintained with sevoflurane** (end-tidal concentration 1.0–1.4%) in left lateral

81 decubitus for 180 minutes. Open reduction and internal fixation by dual anatomic plating with ulnar
82 neurolysis was performed. The intraoperative course was uneventful: stable haemodynamics,
83 normothermia (36.4–36.8 °C), normocapnia (end-tidal CO² 32–36 mmHg), no muscular rigidity, no
84 metabolic acidosis on intraoperative blood gas, glycaemia 1.4–1.8 g/L. Extubation was uneventful and
85 the patient was transferred to the orthopaedic ward in stable condition.

86 **Twenty-two hours after surgery**, the patient developed acute hypotension (64/32 mmHg), oligo-
87 anuria, polypnoea, diffuse myalgia and obtundation. She was urgently transferred to the intensive care
88 unit on norepinephrine support. Investigations revealed massive rhabdomyolysis (CPK 31,010 IU/L;
89 lactate dehydrogenase [LDH] 1,995 IU/L), severe hyperkalaemia (potassium 8.0 mmol/L), KDIGO
90 stage 3 AKI (creatinine 130 µmol/L), severe metabolic acidosis (pH 7.20; bicarbonate 12 mmol/L),
91 severe hyperglycaemia (5.0 g/L), and high-sensitivity troponin I 13,000 pg/mL. Bedside transthoracic
92 echocardiography showed unchanged LVEF (35%) without new regional wall motion abnormalities.
93 Twelve-lead electrocardiography demonstrated peaked T waves typical of hyperkalaemia, without
94 new ischaemic changes or stent-territory ST-segment elevation.

95 The differential diagnosis of acute coronary stent thrombosis was considered but judged unlikely
96 given the absence of characteristic angina, the lack of new ischaemic ECG changes beyond
97 hyperkalaemia signatures, the unchanged echocardiographic pattern, and the recognition that troponin
98 elevation in massive rhabdomyolysis is non-specific (myocardial injury secondary to circulatory
99 collapse, hyperkalaemia and acidosis — type 2 MINS rather than type 1 acute myocardial
100 infarction).¹¹ After multidisciplinary discussion (anaesthesia, intensive care, cardiology, nephrology),
101 management focused on the dominant rhabdomyolysis–AKI–hyperkalaemia syndrome while
102 reinstating dual antiplatelet therapy.

103 Treatment included intravenous calcium gluconate, glucose-insulin, isotonic crystalloid resuscitation
104 calibrated to a urine output target of ≥ 1 mL/kg/h, vasopressors, and emergent intermittent
105 haemodialysis with ultrafiltration. Sodium bicarbonate and mannitol were not used as standard, in line
106 with current evidence.^{12,13} After two hours of dialysis, partial correction was achieved (potassium 4.1
107 mmol/L; creatinine 90 µmol/L; CPK 27,710 IU/L). Three hours after dialysis cessation, the patient
108 deteriorated with extreme bradycardia (32 bpm) and **rebound hyperkalaemia** (6.5 mmol/L),
109 followed by refractory cardiac arrest 35 hours after the index procedure. Resuscitation was
110 unsuccessful. The chronological sequence is summarised in **Figure 1**, and the proposed integrative
111 pathophysiological model in **Figure 2**. Written informed consent for publication was obtained from
112 the patient's next of kin.

113 **DISCUSSION**

114 This observation describes a fulminant fatal postoperative rhabdomyolysis whose temporal pattern —
115 uneventful intraoperative course, asymptomatic 22-hour interval, abrupt biological storm, rebound
116 hyperkalaemia despite renal replacement therapy and refractory arrest — is not explained by any
117 single established mechanism. We argue that it represents a paradigmatic instance of synergistic
118 statin–halogenated myotoxicity amplified by convergent comorbid stressors, and propose that this
119 perioperative phenotype deserves explicit recognition.

120 ***Mechanistic synthesis***

121 Statins inhibit HMG-CoA reductase, depleting mevalonate-derived isoprenoids (geranylgeranyl-PP,
122 farnesyl-PP) required for prenylation of small GTPases (Rab, Rho, Rac), causing impaired vesicular
123 trafficking, sarcolemmal destabilisation, and CoQ10-dependent mitochondrial dysfunction with
124 reduced ATP availability and increased oxidative stress.^{2,3} At the genetic level, the *SLCO1B1*
125 c.521T>C polymorphism reduces hepatic OATP1B1-mediated statin uptake, increasing systemic
126 exposure and myopathy risk roughly 3- to 4-fold for atorvastatin users carrying the CC genotype.⁶ All
127 halogenated agents activate skeletal-muscle RyR1, producing pathological sarcoplasmic Ca²⁺ leak; in
128 *RYR1/CACNAIS* variant carriers (1:200–1:2,000 prevalence), this leak may exceed cellular buffering
129 capacity, manifesting as classical MH or, increasingly, as **delayed atypical anaesthetic-induced**
130 **rhabdomyolysis** without hyperthermia or rigidity — a phenotype consistent with our patient's
131 presentation.^{4,5,7,14} The mechanistic intersection is biologically transparent: a statin-treated myocyte
132 operates with diminished energetic reserves, weakened membrane integrity and impaired Ca²⁺
133 handling, so that a superimposed RyR1-mediated Ca²⁺ overload may overwhelm an already
134 isoprenoid-depleted, CoQ10-deficient mitochondrial network and precipitate irreversible necrosis. A
135 2025 case of severe rhabdomyolysis in a statin user undergoing parathyroidectomy under general
136 anaesthesia advanced the same hypothesis.¹⁰

137 In our patient, four additional cofactors converged: **uncontrolled diabetes** (hyperosmolarity,
138 microvascular dysfunction, electrolyte and phosphate disturbance); **severe ischaemic**
139 **cardiomyopathy** (low cardiac output reducing skeletal-muscle perfusion and metabolic reserve);
140 **prolonged left lateral decubitus** (~3 h, generating direct mechanical compression and ischaemia–
141 reperfusion injury of dependent muscle masses); and **polypharmacy**, with sacubitril/valsartan and
142 spironolactone amplifying hyperkalaemia risk and surgical stress potentially impairing hepatic
143 CYP3A4 clearance of atorvastatin. The marked troponin elevation was attributed to type 2 myocardial
144 injury secondary to circulatory collapse, hyperkalaemia and acidosis, rather than to a primary type 1
145 ischaemic event. The complete six-pathway model is presented in **Figure 2**.

146 ***Lessons for management***

147 Three management lessons emerge. First, postoperative myalgia was masked by surgical pain until
148 frank circulatory deterioration; in highest-risk patients, **systematic biochemical surveillance** (CPK,
149 potassium, creatinine, urine output, serial ECG) during the first 24 hours may allow earlier detection.
150 Second, regarding renal replacement, intermittent haemodialysis with standard high-flux membranes
151 has limited capacity to clear myoglobin (17.8 kDa). Heyne et al. demonstrated up to 20-fold higher
152 myoglobin clearances using high cut-off (HCO) membranes (median clearance 44.2 vs 3.7 mL/min
153 for conventional HD).¹⁵ The 2025 *ReplaceRhabdo* pilot randomised trial confirmed that continuous
154 veno-venous haemodialysis with HCO dialyzers and haemoadsorption-augmented strategies offer
155 superior myoglobin elimination versus conventional CVVH.¹⁶ These findings support a low threshold
156 for **continuous high cut-off RRT** in massive ongoing rhabdomyolysis. Third, regarding adjunctive
157 therapies, the EAST (Eastern Association for the Surgery of Trauma) guideline conditionally
158 recommends *against* bicarbonate or mannitol, with isotonic crystalloid resuscitation as the
159 cornerstone.^{12,13}

160 ***Proposed STAT-OP framework***

161 We propose that statin-treated patients undergoing inhalational anaesthesia warrant a structured
162 perioperative approach, the **STAT-OP** framework (Statin–Anaesthetic Toxicity in the Operative
163 Period), articulated across four phases.

164 **Preoperative.** Identify the high-risk profile: high-intensity statin (atorvastatin \geq 40 mg, rosuvastatin \geq
165 20 mg), uncontrolled diabetes (HbA1c \geq 8%), advanced heart failure (LVEF \leq 40%), chronic kidney
166 disease (eGFR \leq 60 mL/min/1.73 m²), polypharmacy with CYP3A4 modulators, and surgery
167 requiring prolonged dependent positioning. Continue statin per cardiovascular guidelines but
168 document the risk in the anaesthetic record.⁸ Optimise glycaemic control. Withhold clopidogrel 5 days
169 and SGLT2 inhibitors 3 days. Measure baseline CPK, creatinine, electrolytes. Where
170 pharmacogenomic testing is available, *SLCO1B1* c.521T>C and *RYR1/CACNA1S* genotyping may
171 further refine risk in patients with previous unexplained myalgia or family history.^{6,7}

172 **Intraoperative.** In the highest-risk subset, prefer **total intravenous anaesthesia (TIVA)** with
173 propofol-based maintenance over halogenated agents. TIVA avoids RyR1 activation entirely and has
174 been associated with reduced inflammatory and oxidative-stress responses in observational and
175 randomised work.^{17,19} Avoid succinylcholine. Apply strict positioning protocols with pressure-
176 distribution devices. Maintain normothermia, normocapnia and normovolaemia; monitor temperature,
177 ETCO₂, arterial pressure and neuromuscular blockade.

178 **Early postoperative (0–24 h).** Active inquiry for myalgia, weakness and dark urine. Serial CPK at
179 H+6, H+12 and H+24 in selected patients. Continuous ECG and electrolyte surveillance. Maintain
180 isotonic crystalloid infusion with urine output \geq 1 mL/kg/h.

181 **Established rhabdomyolysis.** Discontinue statin. Aggressive isotonic crystalloid resuscitation
182 calibrated to a urine-output target of \geq 1 mL/kg/h, while avoiding both under-resuscitation and
183 excessive volumes that may precipitate fluid overload.^{12,18,20} Treat hyperkalaemia urgently. Avoid
184 bicarbonate and mannitol unless individualised indication exists.^{12,13} Low threshold for ICU transfer
185 and **continuous high cut-off renal replacement therapy** when ongoing necrosis is suspected, with
186 consideration of haemoadsorption-augmented strategies in selected refractory cases.^{15,16,18}

187 ***Strengths and limitations***

188 The principal limitation is the single-patient nature of the observation: causality cannot be
189 quantitatively apportioned among the convergent mechanisms. Muscle biopsy, in vitro contracture
190 testing, *RYR1/CACNA1S* and *SLCO1B1* genotyping, atorvastatin plasma levels and autopsy were not
191 available. The proposed synergy and the STAT-OP framework therefore remain hypothesis-
192 generating, and confirmation requires translational studies, pharmacovigilance signal analyses and
193 ideally a prospective registry of perioperative rhabdomyolysis in statin users. The principal strength is
194 the integration of mechanistic, genetic, and management dimensions into a coherent model anchored
195 in 2020–2025 high-quality evidence.

196 **CONCLUSION**

197 We report a fatal postoperative rhabdomyolysis emerging within 22 hours of elective orthopaedic
198 surgery under sevoflurane-based anaesthesia in a high-risk diabetic patient on chronic high-intensity
199 atorvastatin. The clinical pattern is most consistent with a multifactorial process in which a probable
200 statin–halogenated synergy was amplified by uncontrolled diabetes, advanced cardiomyopathy, lateral

201 decubitus positioning and hyperkalaemia-promoting polypharmacy. While guidelines correctly
202 recommend perioperative statin continuation, this case highlights three actionable points:
203 individualised myotoxic risk stratification (potentially augmented by *SLCO1B1* and *RYR1*
204 pharmacogenomics), a low threshold for total intravenous anaesthesia in the highest-risk subset, and
205 prompt deployment of continuous high cut-off renal replacement therapy when massive necrosis is
206 suspected. The proposed **STAT-OP** framework provides a structured starting point for prospective
207 evaluation, and we encourage anaesthesiologists, intensivists, cardiologists and pharmacologists to
208 recognise and report this clinically meaningful, potentially preventable, and currently under-reported
209 perioperative phenotype.

210 **ACKNOWLEDGEMENTS**

211 [Detailed acknowledgements are provided in the separate Title Page in accordance with the IJAR
212 double-blind peer-review policy.] The authors thank the multidisciplinary clinical team involved in
213 the care of the patient.

214 **ETHICAL CONSIDERATIONS AND CONSENT**

215 Written informed consent for the publication of this anonymised case report and accompanying data
216 was obtained from the patient's next of kin in accordance with the principles of the Declaration of
217 Helsinki (revised 2013). Institutional review board approval was waived in accordance with local
218 regulations governing single anonymised case reports. The signed consent form is archived under the
219 authors' responsibility and is available upon request from the journal editorial office. This case report
220 was prepared in accordance with the CARE (CAse REport) reporting guideline.

221 **CONFLICTS OF INTEREST**

222 The authors declare that no conflict of interest, financial or otherwise, exists in relation to this work.

223 **FUNDING**

224 This work received no specific grant from any funding agency in the public, commercial, or not-for-
225 profit sectors.

226 **USE OF ARTIFICIAL INTELLIGENCE TOOLS**

227 Artificial intelligence tools were used solely for English-language editing and clarity refinement. No
228 AI tools were used for data collection, analysis, interpretation, or image generation. The authors retain
229 full responsibility for the content of the manuscript.

230 **REFERENCES**

- 231 1. Chavez LO, Leon M, Einav S, Varon J. Beyond muscle destruction: a systematic review of
232 rhabdomyolysis for clinical practice. *Crit Care*. 2016;20(1):135.
- 233 2. Sakamoto K, Kimura J. Mechanism of statin-induced rhabdomyolysis. *J Pharmacol Sci*.
234 2013;123(4):289–294.

- 235 3. du Souich P, Roederer G, Dufour R. Myotoxicity of statins: mechanism of action. *Pharmacol Ther.*
236 2017;175:1–16.
- 237 4. Riazi S, Kraeva N, Hopkins PM. Malignant hyperthermia in the post-genomics era: new
238 perspectives on an old concept. *Anesthesiology.* 2018;128(1):168–180.
- 239 5. Hopkins PM, Gupta PK, Bilmen JG. Malignant hyperthermia. *Handb Clin Neurol.* 2018;157:645–
240 661.
- 241 6. Turongkaravee S, Jittikoon J, Lukkunaprasit T, Sangroongruangsri S, Chaikledkaew U,
242 Thakkinstian A. A systematic review and meta-analysis of genotype-based and individualized
243 data analysis of SLCO1B1 gene and statin-induced myopathy. *Pharmacogenomics J.*
244 2021;21(3):296–307.
- 245 7. Carsana A. Exercise-induced rhabdomyolysis and stress-induced malignant hyperthermia events,
246 association with malignant hyperthermia susceptibility, and RYR1 gene sequence variations.
247 *Sci World J.* 2013;2013:531465.
- 248 8. Thompson A, Fleischmann KE, Smilowitz NR, et al. 2024
249 AHA/ACC/ACS/ASNC/HRS/SCA/SCCT/SCMR/SVM guideline for perioperative
250 cardiovascular management for noncardiac surgery: a report of the American College of
251 Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines.
252 *Circulation.* 2024;150(19):e351–e442.
- 253 9. Putzu A, de Carvalho e Silva CMPD, de Almeida JP, Belletti A, Cassina T, Landoni G, et al.
254 Perioperative statin therapy in cardiac and non-cardiac surgery: a systematic review and meta-
255 analysis of randomized controlled trials. *Ann Intensive Care.* 2018;8(1):95.
- 256 10. Adiat M, Ravikumar A, Yadav S, Chandrasekaran V, Kannan M, Rajan N, et al. A case report of
257 severe rhabdomyolysis following parathyroidectomy: a likely synergistic effect of statins and
258 general anaesthetics. *Cureus.* 2025;17(7):e89047.
- 259 11. Devereaux PJ, Szczeklik W. Myocardial injury after non-cardiac surgery: diagnosis and
260 management. *Eur Heart J.* 2020;41(32):3083–3091.
- 261 12. Cabrera JL, Auerbach JS, Merola J, Carlin BW, Brown CVR, Tominaga GT, et al. Management
262 of rhabdomyolysis: a practice management guideline from the Eastern Association for the
263 Surgery of Trauma. *Am J Surg.* 2022;224(1 Pt B):196–204.
- 264 13. Somagutta MR, Pagad S, Sridharan S, Nanthakumaran S, Arnold AA, May V, et al. Role of
265 bicarbonates and mannitol in rhabdomyolysis: a comprehensive review. *Cureus.*
266 2020;12(8):e9742.
- 267 14. Belitova M, Nikolova GG, Usheva S, Mladenova MT, Marinov T, Kaneva R, et al. Diagnostic
268 challenges in malignant hyperthermia and anesthesia-induced rhabdomyolysis: a case study.
269 *Am J Case Rep.* 2024;25:e946306.
- 270 15. Heyne N, Guthoff M, Krieger J, Haap M, Häring HU. High cut-off renal replacement therapy for
271 removal of myoglobin in severe rhabdomyolysis and acute kidney injury: a case series. *Nephron*
272 *Clin Pract.* 2012;121(3–4):c159–c164.

- 273 16. Weidhase L, Borrmann A, Willenberg A, Mende M, Scharf-Janßen C, Petros S, et al. Kidney
274 REPLACEMENT therapies in patients with acute kidney injury and RHABDOMYOLYSIS
275 (ReplaceRhabdo): a pilot trial. BMC Nephrol. 2025;26(1):23.
- 276 17. Yao J, Gao Z, Qu W, Li J. Propofol total intravenous anesthesia vs. sevoflurane inhalation
277 anesthesia: effects on post-operative cognitive dysfunction and inflammation in geriatric
278 patients undergoing laparoscopic surgery. Exp Ther Med. 2024;28(3):343.
- 279 18. Forni L, Aucella F, Bottari G, Büttner S, Cantaluppi V, Fries D, et al. Hemoadsorption therapy for
280 myoglobin removal in rhabdomyolysis: consensus of the hemoadsorption in rhabdomyolysis
281 task force. BMC Nephrol. 2024;25(1):247.
- 282 19. Nance JR, Mammen AL. Diagnostic evaluation of rhabdomyolysis. Muscle Nerve.
283 2015;51(6):793–810.
- 284 20. Bosch X, Poch E, Grau JM. Rhabdomyolysis and acute kidney injury. N Engl J Med.
285 2009;361(1):62–72.

286 **FIGURE LEGENDS**

287 **Figure 1.** Chronological timeline of perioperative events. From the index orthopaedic surgery (T0)
288 through an asymptomatic interval to abrupt clinical deterioration at T+22 h, ICU admission with
289 massive rhabdomyolysis and life-threatening hyperkalaemia, partial correction after haemodialysis,
290 rebound hyperkalaemia at T+33 h, and refractory cardiac arrest at T+35 h. CPK = creatine
291 phosphokinase; LVEF = left ventricular ejection fraction; HD = haemodialysis.

292 **Figure 2.** Proposed integrative pathophysiological model. Six convergent injurious pathways acting
293 on skeletal muscle generate massive necrosis (CPK 31,010 IU/L), triggering myoglobinuric AKI, life-
294 threatening hyperkalaemia, metabolic acidosis, cardiac irritability and systemic inflammation,
295 culminating in refractory cardiac arrest. RyR1 = ryanodine receptor type 1; CYP3A4 = cytochrome
296 P450 3A4; AKI = acute kidney injury; SIRS = systemic inflammatory response syndrome; SR =
297 sarcoplasmic reticulum; MH = malignant hyperthermia.

298 *[Figures 1 and 2 are submitted as separate high-resolution JPEG files, in accordance with the IJAR submission*
299 *requirements: > 1800 × 1200 px, < 4 MB.]*