

1
2
3
4 **Histopathological Toxicity Evaluation of Deltamethrin on the Integrity of Ovarian Follicles**
5 **on Mice (*Mus musculus*).**
6
7
8
9
10

11 Abstract— Deltamethrin is a widely used pyrethroid insecticide. Exposure to deltamethrin can
12 have negative effects on non-target organisms, such as mammals. Metabolites of deltamethrin
13 can trigger the formation of reactive oxygen species (ROS) and further cause oxidative stress.
14 Oxidative stress can cause cellular damage to lipids, DNA, and proteins, and may further lead to
15 cell death and apoptosis. One of the target organs of deltamethrin is the ovary.

16 This study aims to investigate the effects of deltamethrin insecticide exposure on the ovarian
17 structure of mice (*Mus musculus* L.). The study subjects consisted of 15 female mice, divided
18 into three treatment groups. The first group, the control group, received corn oil without
19 deltamethrin insecticide. The second and third groups (D1 and D2) received deltamethrin
20 insecticide at doses of 18 and 36 mg/kg body weight, respectively. The treatment was
21 administered intraperitoneally for 14 days, with a daily dose of 1 ml.

22 Data on the number of primordial, primary, secondary, Graafian, and atretic follicles were
23 statistically analyzed using one-way analysis of variance (ANOVA) and post hoc Duncan's
24 Multiple Range Test (DMRT). The results of this study indicate that the administration of
25 deltamethrin insecticide at doses of 18 and 36 mg/kg BW in mice (*Mus musculus* L.) caused
26 damage to ovarian structure, including a decrease in the number of secondary and Graafian
27 follicles, and an increase in the number of atretic follicles.

28 **Key words:** *deltamethrin, insecticide, follicle, ovary, mice.*
29

30 **I. INTRODUCTION**

31 Deltamethrin is a widely used pyrethroid insecticide. Exposure to deltamethrin not only affects insects but can also
32 negatively impact non-target organisms, such as mammals. Toxic effects caused by deltamethrin exposure on non-
33 target organisms include disruption of the nervous system and other organs. Direct contact with the skin can cause
34 irritation, itching, rashes, and inflammation. Contact with the eyes can cause irritation, inflammation, and redness.
35 Inhalation of deltamethrin can irritate the respiratory tract, causing coughing and difficulty breathing. Ingestion can
36 cause digestive problems, nausea, vomiting, and diarrhea (Tomassoni et al., 2015).

37 Deltamethrin, once ingested, is metabolized by microsomal enzymes in the liver and broken down into metabolites.
38 Deltamethrin metabolites can trigger the formation of reactive oxygen species (ROS). Continuous exposure to
39 deltamethrin, when the amount of ROS in the body is too high, can lead to oxidative stress. This oxidative stress can
40 cause damage to lipids, DNA, and proteins, and can further lead to cell death and apoptosis (Sharma et al., 2014).
41

42 Rehman et al. (2006) stated that oral administration of deltamethrin at doses of 5, 6, and 18 mg/kgBW for 15 days in
43 Swiss strain mice significantly induced lipid peroxidation (LPO) in the liver. In line with this, there was a decrease
44 in the activity of antioxidant enzymes such as glutathione peroxidase (GPx), glutathione S-transferase (GST), and

catalase. Research by Otçu et al. (2023) showed that intraperitoneal administration of deltamethrin at a dose of 30 mg/kgBW in Wistar rats during the 6-21 day gestation period caused follicular degeneration, pyknotic nuclei, and hyperplasia cells in the mother rats. Another study conducted by Ali and Farzaneh (2014) showed the effect of deltamethrin at doses of 2.5, 5, and 10 mg/kgBW, causing a decrease in the number of primary and secondary follicles, an increase in the number of atretic follicles, and a significant decrease in the corpus luteum. Previous research used pure deltamethrin, so further research is needed on the effect of administering the insecticide deltamethrin on the ovarian structure of mice (*Mus musculus L.*).

II. MATERIALS AND METHODS

Research materials

The test animals used in this study were 15 female mice, aged 12 weeks with an average body weight 26.7 grams. The mice were acclimatized for 7 days (1 week) and given food and water ad libitum. The test material used was the insecticide deltamethrin® in packaging with a deltamethrin content of 25 grams/liter.

Research procedures

Mice were treated with deltamethrin at doses of 18 mg/kgBW, 36 mg/kgBW, and controls were given corn oil or no deltamethrin. Treatment was carried out daily for 14 days intraperitoneally with a dose of 1 ml/day. On the 15th day, the mice were anesthetized and dissected, then the ovaries were removed for histological preparations using the paraffin method and Hematoxylin-Eosin staining. The parameters observed in this study included the number of primordial, primary, secondary, Graafian, and atresi follicles. Ovarian histology slides were observed using a binocular microscope at 400x magnification.

Statistical Analysis

The research data was analyzed quantitatively using a One-Way ANOVA test with a 95% significance level ($\alpha = 0.05$) to evaluate the effect of the dose treatment. Duncan's test was performed to identify significant differences between treatment groups.

III. RESULTS AND DISCUSSION

In this study, we observed the number of follicles in mice (*Mus musculus L.*) after administration of the insecticide deltamethrin. The observed follicles included primordial, primary, secondary, Graafian, and atresi follicles. The results of the calculation of the number of follicles in each treatment group are shown in Table 1. Based on one-way ANOVA analysis, the p value > 0.05 was shown for the number of primordial and primary follicles, but the p value < 0.05 was shown for the number of secondary, Graafian, and atresi follicles. This indicates that the treatment of deltamethrin insecticide did not significantly affect the average number of primordial and primary follicles but significantly affected the average number of secondary, Graafian, and atresi follicles. The results of further tests using DMRT (Table 1) showed that the number of secondary and atresi follicles in treatment K was not significantly different from treatment D1, but significantly different from treatment D2 and the number of Graafian follicles in treatment K was significantly different from treatments D1 and D2.

Table 1 Average number of primordial, primary, secondary, Graafian and atresi follicles in mice after administration of deltamethrin insecticide

Follicle type	Average number of follicles ($\bar{x} \pm SD$)		
primordial	20 ± 17,25 ^a	6 ± 12,89 ^a	15,77 ^a
primary	8 ± 28,39 ^a	3 ± 26,86 ^a	17,32 ^a

secondary	$10 \pm 2,30^a$	$\pm 2,00^a$	$\pm 1,83^b$	93
antral	$10 \pm 1,30^a$	$\pm 1,00^b$	$\pm 0,82^b$	94
antral	$10 \pm 1,48^a$	$\pm 8,89^a$	$\pm 11,09^b$	95
				96

Description: K (control), D1 (dose 1: 18 mg/kgBW), D2 (dose 2: 36 mg/kgBW), \bar{x} (mean), SD (standard deviation). The same letter notation behind the numbers in the same row indicates no significant difference between treatment groups.

Deltamethrin that enters the body of an organism will be metabolized by the body. The main metabolic results of deltamethrin are oxidative metabolites (2', 4'- and 5-OH-deltamethrin) and metabolites of trans-methyl and ester groups (3-phenoxybenzoic acid (3-PBA) and 4'- and 2'-OH-PBA) (Lu et al., 2018). Previous research in rodents showed that deltamethrin is rapidly metabolized by esterase, which is then widely distributed by liver microsomal enzymes (Rehman et al., 2014). Deltamethrin metabolism in mice produces the main metabolites in the form of 3-PBA, 4'- and 5'-OH-deltamethrin (Shono et al., 1979). Other research added that 4-OH-deltamethrin is more toxic than deltamethrin in SH-SY5Y cells. The results of the study by Romero et al. (2012) showed that deltamethrin and its metabolites cause oxidative stress.

Oxidative stress is a condition of imbalance between oxidants and antioxidants in the body, through the excessive formation of free radicals and/or ROS, such as hydroxyl (OH), superoxide (O₂⁻), and hydrogen peroxide (H₂O₂) (Lu et al., 2018). Research by Rehman et al. (2006) states that deltamethrin causes oxidative stress in the body through the formation of ROS. The results of other studies support this statement, which show that deltamethrin significantly increases ROS production in PC12 cells (Li et al., 2007).

The presence of ROS in the ovaries is found in follicular fluid. Follicular fluid contains macrophages, leukocytes, and cytokines, which are known sources of ROS. ROS present in this follicular fluid plays a role in oocyte maturation and development. This suggests that ROS at certain levels are necessary for ovarian physiological processes. However, when ROS accumulates beyond a certain level or even too high in the ovaries, it can cause a decrease in follicular fluid, inhibit meiosis, disrupt oocyte maturation, cause direct damage to the oocyte, and even lead to infertility (Ali and Farzaneh, 2014).


Several types of pyrethroids and their metabolites can disrupt the function of several hormone receptors, further affecting the endocrine system and reproductive organs. Several studies have identified deltamethrin and its metabolites as endocrine disruptors (Marettova et al., 2017). Deltamethrin is considered a compound similar to estrogen. Consequently, deltamethrin can inhibit estrogen action through direct interaction with estrogen receptors and may function as an endocrine modulator (Kim et al., 2004).

Estrogen is a steroid hormone that plays a crucial role in regulating the female reproductive system. In the ovaries, estrogen is produced by follicle cells or granulosa cells. Estrogen production in granulosa cells occurs when Luteinizing Hormone (LH) stimulates theca cells, resulting in the production of androgens. The aromatase enzyme found in granulosa cells, along with Follicle Stimulating Hormone (FSH) stimulation, converts androgens into estrogen. Estrogen binds to estrogen receptors in target cells, such as granulosa cells and the endometrium. Activation of these receptors regulates the production of gonadotropin hormones, FSH and LH, by the anterior pituitary, and induces follicle proliferation and maturation. When deltamethrin binds to estrogen receptors, it can disrupt the regulation of FSH and LH production, as well as disrupt follicle proliferation and maturation (Marettova et al., 2017).


Based on research by Ali and Farzaneh (2014), it was shown that administration of deltamethrin at doses of 2.5, 5, and 10 mg/kgBW to mice resulted in a significant decrease in the number of primary and secondary follicles, and a significant increase in the number of atresia follicles. In this study, it was found that administration of deltamethrin at doses of 18 and 36 mg/kgBW to mice resulted in a significant decrease in the number of secondary and Graafian

149 follicles, and a significant increase in the number of atresia follicles. This indicates that administration of
150 deltamethrin to rodents (in this case, rats and mice) can cause a decrease in the number of follicles at several stages
151 of development and an increase in the number of atresia follicles. The results of observations of mouse ovarian
152 follicles after administration of deltamethrin can be seen in Figure 1.

153
154 folikel ovarium menciit setelah pemberian deltametrin dapat dilihat pada Gambar 1.



155
156
157 (A) (B) (C)




158
159
160
161
162
163
164
165
166
167
168
169

170 Description: (A) Primordial follicle, (B) Primary follicle, (C) Secondary follicle, (D) Graafian follicle, (E) Atresia
171 follicle; (a) Oocyte; (b) Follicular cell; (c) Antrum; (d) Connective tissue

172
173 Figure 1 Mouse ovarian follicles at various stages of development (Magnification 400x)

174
175 The results showed that the number of secondary and Graafian follicles decreased significantly. Furthermore,
176 follicles in the mouse ovaries experienced degeneration (atresia) at various stages of development, as indicated by a
177 significant increase in their number. The higher the deltamethrin dose given to the mice, the more the number of
178 secondary and Graafian follicles decreased, while conversely, the number of atresia follicles increased. The
179 increasing number of atresia follicles with increasing deltamethrin dose can be seen in Figure 2.



180
181 Description: (A) Control; (B) Dose 1; (C) Dose 2; arrows indicate atretic follicles

182
183 Figure 2 Atretic follicles in each treatment group (400x magnification)

184
185
186
187 **IV. CONCLUSION**

188 Administration of the insecticide deltamethrin at doses of 18 and 36 mg/kgBW to mice (*Mus musculus* L.) caused
189 damage to the ovarian structure, including a decrease in the number of secondary and Graafian follicles, as well as
190 an increase in the number of atresia follicles.

191

192 REFERENCES

193 Ali, P.M., & Farzaneh, R. (2014). The effect of vitamin C as antioxidant on the toxic effects due to deltamethrin
194 pesticide on morphometric and histopathological parameters of ovary in the rats. *Biosci. Biotechnol. Res.*
195 *Asia*, 11, 959–964.

196 Kim, S.S., Kwack, S.J., Lee, R.D., Lim, K.J., Rhee, G.S., Seok, J.H., Kim, B.H.,
197 Won, Y.H., Kojima, H., Katsura, E., Takeuchi, S., Niiyama, K., & Kobayashi,
198 K. (2004). Screening for estrogen and androgen receptor activities in 200 pesticides by in vitro reporter gene
199 assays using Chinese hamster ovary cells. *Environ. Health Perspect*, 112, 524–531.

200 Li, H.Y., Zhong, Y.F., Wu, S.Y., Shi, N. (2007). NF-E2 related factor 2 activation and heme oxygenase-1 induction
201 by tert-butylhydroquinone protect against deltamethrin- mediated oxidative stress in PC12 cells. *Chem. Res.*
202 *Toxicol.* 20, 1242-1251.

203 Lu, Q., Sun, Y., Ares, I., Anadón, A., Martínez, M., Martínez-Larrañaga, M. R., Yuan, Z., Wang, X., & Martínez,
204 M. A. (2018). Deltamethrin toxicity: a review of oxidative stress and metabolism. *Environmental Research*,
205 1–86. <https://doi.org/10.1016/j.envres.2018.12.045>

206 Marettova, E., Maretta, M., & Legáth, J. (2017). Effect of pyrethroids on female genital system: Review. *Animal*
207 *Reproduction Science*, 2017, 184, 132-138.

208 Otçu, S., Deveci, E., Özgökçe, Ç., Gürsoy, G. T., & Tuncer, M. C. (2023). Protective effect of nebulolol on rat ovary
209 exposed to deltamethrin toxicity. *Acta Cirurgica Brasileira*, 38, 1–10. <https://doi.org/10.1590/acb385423>

210 Rehman, H., Ali, M., Atif, F., Kaur, M., Bhatia, K., & Raisuddin, S. (2006). The modulatory effect of deltamethrin
211 on antioxidants in mice. *Clinica Chimica Acta*, 369, 61–65. <https://doi.org/10.1016/j.cca.2006.01.010>

212 Rehman, H., Aziz, A. T., Saggu, S., Abbas, Z. K., Mohan, A., & Ansari, A. A. (2014). Systematic review on
213 pyrethroid toxicity with special reference to deltamethrin. *Journal of Entomology and Zoology Studies*, 2(6),
214 60–70.

215 Rehman, H., Aziz, A. T., Saggu, S., Vanwert, A. L., Zidan, N., & Saggu, S. (2017). Additive toxic effect of
216 deltamethrin and cadmium on hepatic, hematological, and immunological parameters in mice. *Toxicology*
217 *and Industrial Health*, 1–8. <https://doi.org/10.1177/0748233716684710>

218 Romero, A., Ramos, E., Castellano, V., Martinez, M.A., Ares, I., Martinez, M., Martinez- Larranaga, M.R., &
219 Anadón, A. (2012). Cytotoxicity induced by deltamethrin and its metabolites in SH-SY5Y cells can be
220 differentially prevented by selected antioxidants. *Toxicol. In Vitro*, 26, 823-830.

221 Sharma, P., R.Singh dan M.Jan. 2014. Dose-dependent effect of deltamethrin in testis, liver and kidney of wistar rat.
222 *Toxicology International*. 21(2), 131-139.

223 Shono, T., Ohsawa, K., Casida, J.E.. (1979). Metabolism of trans-permethrin and cis- permethrin, trans-
224 cypermethrin, and cis-cypermethrin, and decamethrin by microsomal-enzymes. *J. Agric. Food Chem*, 27,
225 316-325.

226 Tomassoni, A.J., French, R.N., & Walter, F.G. (2015). Toxic industrial chemicals and chemical weapons: exposure,
227 identification, and management by syndrome. *Emergency Medicine Clinics*, 33(1), 13-36.

228