

# 1 **Congenital Lobar Emphysema Mimicking Pneumothorax in a 6-Month-Old Infant** 2 **With Acute Respiratory Distress: A Case Report.**

## 3 **Introduction**

4 Congenital lobar emphysema (CLE), increasingly referred to as congenital lobar overinflation, is a rare congenital  
5 lung malformation characterized by abnormal hyperinflation of a pulmonary lobe. Despite the term emphysema,  
6 the condition is not primarily a destructive alveolar disease. Instead, the affected lobe becomes progressively  
7 overdistended, compressing adjacent normal lung parenchyma and potentially causing mediastinal shift,  
8 respiratory distress, and impaired ventilation [1,2].

9 The mechanism is most often related to a ball-valve effect caused by intrinsic bronchial abnormality, such as  
10 deficient bronchial cartilage, bronchomalacia, mucosal folds, or bronchial stenosis. Extrinsic bronchial  
11 compression by vascular structures or mediastinal lesions may also occur, although in many patients no precise  
12 cause is identified [1-3]. CLE usually presents in the first months of life with tachypnea, retractions, wheezing,  
13 cyanosis, poor feeding, or recurrent respiratory infections. Severe cases can present as acute respiratory  
14 emergencies [1,4].

15 The diagnosis can be challenging because the radiographic appearance may resemble pneumothorax. Both  
16 conditions may present with unilateral hyperlucency and contralateral mediastinal displacement. However,  
17 treatment is different: pneumothorax may require urgent pleural drainage, whereas chest tube insertion is not  
18 therapeutic in CLE and may delay appropriate treatment [5,6]. Computed tomography (CT) is essential when the  
19 diagnosis is uncertain because it confirms the intrapulmonary nature of the hyperinflation, identifies the affected  
20 lobe, excludes pleural air, and assesses associated lung disease [2,7].

21 We report a case of congenital lobar emphysema in a 6-month-old infant presenting with acute respiratory  
22 distress, in whom the initial chest radiograph suggested pneumothorax, while CT confirmed CLE and  
23 demonstrated associated contralateral pulmonary involvement.

## 24 **Case Presentation**

25 A 6-month-old male infant was admitted to the emergency department for acute respiratory distress. He had  
26 previously required hospitalization in the pediatric intensive care unit during early infancy for severe respiratory  
27 distress in the context of recurrent lower respiratory tract infection. There was no known history of thoracic  
28 trauma, foreign body aspiration, congenital heart disease, or previous thoracic surgery.

29 The current episode began with cough, fever, progressive tachypnea, and feeding difficulty, followed by  
30 worsening respiratory effort. On admission, the infant was conscious but irritable and dyspneic. He had marked  
31 intercostal and subcostal retractions, nasal flaring, and oxygen saturation of 86% on room air, improving to 95%  
32 under supplemental oxygen. Body temperature was 38.3 degrees C, heart rate was 164 beats/min, and respiratory  
33 rate was 62 breaths/min. Chest examination revealed decreased breath sounds over the left hemithorax with  
34 scattered crackles on the right side.

35 Initial laboratory tests showed inflammatory features, with leukocytosis at 18,200/mm<sup>3</sup> and C-reactive protein at  
36 64 mg/L. Blood gas analysis showed mild respiratory acidosis with hypoxemia before oxygen supplementation.  
37 Because of the acute presentation and asymmetric chest findings, an urgent chest radiograph was obtained.

38 Chest radiography demonstrated marked hyperlucency and overexpansion of the left hemithorax, flattening of the  
39 left hemidiaphragm, compression of adjacent lung parenchyma, and mediastinal shift toward the right side.  
40 Pneumothorax was initially suspected because of the hyperlucent hemithorax and mass effect. However, careful  
41 review showed persistence of fine vascular markings within the hyperlucent area and no clear pleural line. In view  
42 of this diagnostic uncertainty, thoracic CT was performed before any pleural drainage.

43 Thoracic CT excluded pneumothorax and showed major overinflation of the left upper lobe, with stretching of  
44 bronchovascular structures, compression atelectasis of the remaining left lung, and rightward mediastinal  
45 displacement. These findings were consistent with congenital lobar emphysema of the left upper lobe. CT also

46 demonstrated contralateral pulmonary abnormalities, including patchy right basal consolidation and peribronchial  
47 thickening, suggesting associated bronchopneumonia. No endobronchial foreign body, pleural air collection,  
48 diaphragmatic hernia, or obvious mediastinal mass was identified. Echocardiography did not show significant  
49 congenital heart disease.

50 The infant was admitted for close monitoring and stabilization. Initial treatment included humidified oxygen  
51 therapy, careful hydration, antipyretics, respiratory physiotherapy when tolerated, and avoidance of unnecessary  
52 pleural drainage. Because of fever, inflammatory markers, and contralateral parenchymal abnormalities on CT,  
53 empirical intravenous antibiotic therapy with cefotaxime and gentamicin was started, then continued according to  
54 clinical response. Respiratory distress improved partially after supportive and antibiotic treatment, but persistent  
55 left upper lobe hyperinflation and mediastinal mass effect remained evident.

56 After multidisciplinary discussion involving pediatrics, pediatric intensive care, radiology, anesthesia, and  
57 pediatric surgery, operative management was indicated because of symptomatic CLE with significant mass effect.  
58 The infant underwent left posterolateral thoracotomy. Intraoperatively, the left upper lobe was markedly  
59 overinflated and compressed the remaining ipsilateral lung. There was no pleural air leak or mechanical  
60 obstruction. Left upper lobectomy was performed. The remaining lung re-expanded progressively after resection.

61 The postoperative course was favorable. The infant was monitored in the pediatric intensive care unit during the  
62 early postoperative period and was extubated without difficulty. Oxygen therapy was progressively weaned.  
63 Follow-up chest radiography showed satisfactory re-expansion of the remaining left lung and improvement of  
64 mediastinal position. Histopathological examination confirmed features compatible with congenital lobar  
65 emphysema, including overdistended alveolar spaces without malignant lesion. The infant was discharged in good  
66 clinical condition after 9 postoperative days. At 1-month follow-up, he had no respiratory distress, good feeding  
67 tolerance, and no clinical evidence of recurrence.

## 68 **Discussion**

69 This case illustrates a major diagnostic pitfall in pediatric respiratory emergencies: congenital lobar emphysema  
70 can closely mimic pneumothorax on initial chest radiography. In both conditions, the radiograph may show a  
71 hyperlucent hemithorax, mediastinal displacement, and apparent compression of the opposite lung. In an unstable  
72 infant, this similarity can lead to urgent chest tube insertion. However, pleural drainage does not treat CLE and  
73 may cause complications or delay definitive management [5,6].

74 Several radiographic signs may suggest CLE rather than pneumothorax. In CLE, the hyperlucency is  
75 intrapulmonary and vascular markings may still be visible within the affected area, although they are stretched  
76 and attenuated. A definite pleural line is absent. The affected lobe is overexpanded and may cause compression of  
77 adjacent lung, diaphragmatic depression, and mediastinal shift. In pneumothorax, peripheral vascular markings  
78 are absent beyond the pleural line. These distinctions are important but may be subtle on emergency radiographs,  
79 particularly in infants with severe respiratory distress [2,7].

80 CT was decisive in our patient. It confirmed that the hyperlucency represented lobar overinflation rather than  
81 pleural air, identified the left upper lobe as the affected lobe, and demonstrated compression of adjacent lung. It  
82 also showed contralateral parenchymal disease, which was clinically relevant because the infant had fever and  
83 inflammatory markers. The contralateral inflammatory changes justified antibiotic therapy and careful  
84 stabilization before surgery. CT also helps exclude other differential diagnoses, including congenital pulmonary  
85 airway malformation, pneumatocele, bronchogenic cyst, diaphragmatic hernia, Swyer-James syndrome, and  
86 foreign body obstruction [1,2,7].

87 The left upper lobe is the most commonly involved lobe in CLE, followed by the right middle lobe and right  
88 upper lobe. Bilateral or multilobar disease is less frequent but has been reported and may complicate management  
89 [2,3,8]. Associated congenital heart disease has also been described, so echocardiographic evaluation is often  
90 useful before surgery, especially when the infant presents with severe symptoms or when anesthesia is planned  
91 [1,3].

92 Treatment depends on clinical severity. Conservative management may be considered in selected mildly  
93 symptomatic or incidentally diagnosed cases under close follow-up [9]. However, severe or persistent respiratory  
94 distress, progressive hyperinflation, recurrent infection, or marked mediastinal shift usually favors surgery.  
95 Lobectomy of the affected lobe remains the standard treatment for symptomatic CLE and generally provides  
96 excellent outcomes, as observed in our patient [2-4,10].

97 The present case has several practical messages. First, the absence of a clear pleural line on chest radiography  
98 should prompt reconsideration of pneumothorax before chest tube insertion. Second, CT should be obtained when  
99 the infant is stable enough and the diagnosis is uncertain. Third, associated infection or contralateral lung disease  
100 may influence timing and perioperative management. Finally, multidisciplinary decision-making is essential  
101 because these patients may require intensive respiratory support, specialized anesthesia, and pediatric thoracic  
102 surgical expertise.

### 103 **Conclusion**

104 Congenital lobar emphysema is a rare but important cause of respiratory distress in infants. It may mimic  
105 pneumothorax on chest radiography, especially when unilateral hyperlucency and mediastinal shift are present.  
106 Thoracic CT is essential to confirm the diagnosis, avoid inappropriate pleural drainage, identify associated  
107 pulmonary lesions, and plan treatment. In symptomatic infants with significant mass effect, surgical lobectomy  
108 after stabilization is an effective treatment with favorable outcome.

### 109 **Learning Points**

- 110 • Congenital lobar emphysema should be considered in any infant with respiratory distress and unilateral  
111 hyperlucency on chest radiography.
- 112 • Persistence of vascular markings and absence of a clear pleural line favor lobar overinflation rather than  
113 pneumothorax.
- 114 • Thoracic CT is the key investigation when pneumothorax and congenital lobar emphysema are difficult to  
115 distinguish.
- 116 • Contralateral pulmonary infection or inflammatory disease may justify antibiotic therapy before definitive  
117 surgery.
- 118 • Symptomatic congenital lobar emphysema with significant mass effect is usually treated by lobectomy, with  
119 good postoperative outcomes.

### 120 **Declarations**

121 Ethical approval: This anonymized single-patient case report does not include experimental intervention. Local  
122 institutional requirements should be followed for case report publication.

123 Consent: Publication of any real patient case and clinical images requires written informed consent from the  
124 parent or legal guardian before journal submission.

125 Conflicts of interest: The author declares no conflicts of interest.

126 Funding: No specific funding was received for this work.

127 Data availability: The clinical data supporting this case report are available from the treating team and are not  
128 publicly shared in order to preserve patient confidentiality.

### 129 **References**

- 130 1. Mukhtar S, Baloch NU, Anjum F. Congenital Lobar Emphysema. In: StatPearls. Treasure Island (FL):  
131 StatPearls Publishing; 2024.
- 132 2. Demir OF, Hangul M, Kose M. Congenital lobar emphysema: diagnosis and treatment options. Int J Chron  
133 Obstruct Pulmon Dis. 2019;14:921-928.

- 134 3. Karnak I, Senocak ME, Ciftci AO, Buyukpamukcu N. Congenital lobar emphysema: diagnostic and therapeutic  
135 considerations. *J Pediatr Surg.* 1999;34(9):1347-1351.
- 136 4. Abdel-Bary MA, Abdel-Gawad M, Abdellatif M, et al. Clinical and surgical aspects of congenital lobar over-  
137 inflation: a single-center experience. *J Cardiothorac Surg.* 2020;15:95.
- 138 5. Prabhu SM, Choudhury SR, Solanki RS, Shetty GS, Agarwala S. Inadvertent chest tube insertion in congenital  
139 cystic lung lesions: highlighting an important problem. *Indian J Radiol Imaging.* 2013;23(1):8-14.
- 140 6. Ulku R, Onat S, Ozcelik C. Congenital lobar emphysema: differential diagnosis and therapeutic approach.  
141 *Pediatr Int.* 2008;50(5):658-661.
- 142 7. Kravitz RM. Congenital malformations of the lung. *Pediatr Clin North Am.* 1994;41(3):453-472.
- 143 8. Perea L, Blinman T, Piccione J, Laje P. Bilateral congenital lobar emphysema: staged management. *J Pediatr*  
144 *Surg.* 2017;52(9):1442-1445.
- 145 9. Mulvany JJ, Weatherall A, Charlton A. Congenital lobar emphysema: diagnostic and therapeutic dilemmas. *J*  
146 *Paediatr Child Health.* 2016;52(8):835-837.
- 147 10. Ibrahim R, Irodi A, Venkatachalapathy E, et al. Congenital lobar emphysema case report. *Radiol Case Rep.*  
148 2022;17(7):2426-2430.