

Stress During Clinical Postings in Anaesthesia and Operation Theatre Technology Education: A Narrative Review.

Abstract

Background: Clinical posting is an integral component of Anaesthesia and Operation Theatre Technology (A&OTT) education, providing students with opportunities to apply theoretical knowledge in real clinical settings. During clinical postings in operation theatres, intensive care units, emergency departments, labour rooms, and post-anaesthesia care units, students acquire essential technical, clinical, and professional competencies required for safe and effective perioperative patient care. However, exposure to demanding clinical environments, academic workload, unfamiliar procedures, fear of making mistakes, communication difficulties, language barriers, and limited clinical experience often results in considerable stress.

Objective: This narrative review aims to summarize the available literature on stress among Anaesthesia and Operation Theatre Technology students during clinical postings, focusing on the major sources of stress, its effects on students, assessment methods, coping strategies, and interventions to reduce stress.

Methods: A narrative review was conducted using published literature retrieved from electronic databases. Original research articles, systematic reviews, and review articles published in English were included. Owing to the limited literature specifically related to A&OTT students, evidence from operating room, anaesthesia, nursing, medical, and allied health students was also reviewed.

Results: The literature indicates that stress during clinical postings is influenced by academic, clinical, personal, and environmental factors. Common stressors include heavy academic demands, fear of clinical errors, inadequate knowledge and practical skills, communication and language barriers, workload, emotional exposure to critically ill patients, and insufficient supervision. These stressors may adversely affect students' confidence, psychological well-being, academic performance, and clinical competence. Supportive faculty mentorship, simulation-based learning, peer support, stress management programmes, and communication skills training were consistently reported as effective strategies for reducing stress.

Conclusion: Stress is a common challenge during clinical postings among A&OTT students. Strengthening clinical supervision, promoting supportive learning environments, and implementing evidence-based stress management interventions may enhance students' well-being, clinical competence, and overall educational experience.

Keywords: Stress; Clinical Posting; Anaesthesia and Operation Theatre Technology; Allied Health Students; Clinical Education; Coping Strategies.

1. Introduction

Clinical education is a fundamental component of healthcare professional training, bridging the gap between theoretical knowledge and practical application. For students pursuing Anaesthesia and Operation Theatre Technology (A&OTT), clinical postings provide valuable opportunities to develop technical skills, clinical reasoning, professional behaviour, and practical competence in real healthcare settings. These postings are undertaken in various clinical areas, including operation theatres, intensive care units (ICUs), emergency departments, labour rooms, and post-anaesthesia care units (PACUs). Under the supervision of anaesthesiologists, surgeons, nurses, and clinical instructors, students gain hands-on experience in perioperative patient care while becoming familiar with anaesthesia equipment, patient monitoring, infection prevention practices, operation theatre protocols, and multidisciplinary teamwork.

Although clinical postings are essential for professional development, they are also recognized as one of the most challenging phases of healthcare education. Students are expected to apply theoretical knowledge in real clinical situations while adapting to unfamiliar hospital environments, mastering new technologies, maintaining patient safety, and fulfilling academic responsibilities. The transition from classroom learning to direct patient care often creates uncertainty, particularly among students with limited clinical exposure. Fear of making mistakes, inadequate knowledge and practical skills, communication difficulties with patients and healthcare professionals, language barriers, long clinical hours, and insufficient supervision have been consistently identified as important sources of stress among healthcare students.

Stress refers to the physical and psychological responses that occur when individuals perceive that academic or environmental demands exceed their ability to cope. While moderate stress may enhance motivation and improve learning, excessive or prolonged stress can adversely affect students' psychological well-being, academic achievement, clinical performance, and professional development. High stress levels have been associated with anxiety, depression, emotional exhaustion, burnout, sleep disturbances, impaired concentration, and reduced self-confidence. These consequences not only influence students' educational experiences but may also affect patient safety and the quality of healthcare delivery.

The operating theatre is a particularly demanding clinical learning environment because it requires strict adherence to aseptic techniques, technical precision, rapid decision-making, effective communication, and close collaboration within multidisciplinary healthcare teams. A&OTT students are expected to become familiar with anaesthesia workstations, monitoring devices, surgical instruments, infection control measures, and emergency protocols while simultaneously fulfilling academic requirements. Exposure to critically ill patients, emergency surgeries, unexpected intraoperative events, and high expectations from faculty

members and clinical staff may further contribute to psychological stress during clinical training.

Numerous studies have investigated stress among nursing, medical, operating room, surgical technology, and other allied health students, identifying academic workload, clinical responsibilities, inadequate preparation, interpersonal relationships, and environmental challenges as major contributors to stress. Persistent stress has been shown to impair learning, reduce clinical performance, increase the risk of burnout, and negatively influence students' overall educational experience. Several interventions, including simulation-based education, faculty mentorship, peer support, stress management programmes, mindfulness training, and communication skills development, have demonstrated promising outcomes in reducing stress and improving students' confidence during clinical education.

Although stress among healthcare students has been widely studied, findings remain scattered across different healthcare disciplines. To date, evidence focusing specifically on Anaesthesia and Operation Theatre Technology students remains limited. Most available studies have been conducted among nursing, medical, and operating room students, whose clinical experiences share similarities with those of A&OTT students. Therefore, synthesizing evidence from these related disciplines provides valuable insights into the stressors experienced by A&OTT students and highlights evidence-based strategies that may improve their clinical learning experience.

This narrative review aims to synthesize the current evidence on stress among Anaesthesia and Operation Theatre Technology students during clinical postings. It discusses the major sources of stress, their effects on academic and clinical performance, commonly used stress assessment tools, coping strategies adopted by students, and evidence-based interventions to reduce stress. By identifying the factors contributing to stress and summarizing the available evidence, this review seeks to support educators, clinical instructors, and educational institutions in developing strategies that promote student well-being, clinical competence, and high-quality clinical education.

2. Methodology

This narrative review was conducted to summarize the available literature on stress experienced by Anaesthesia and Operation Theatre Technology (A&OTT) students during clinical postings. A comprehensive literature search was performed using electronic databases. Search terms included "stress," "clinical stress," "clinical posting," "clinical education," "Anaesthesia Technology students," "Operation Theatre Technology students," "Operating Room students," "Surgical Technology students," "Allied Health students," "Nursing students," "Medical students," "burnout," "anxiety," "communication barriers," and "coping strategies."

Original research articles, systematic reviews, review articles, and qualitative studies published in English were considered for inclusion. As the literature specifically related to

A&OTT students is limited, studies involving operating room, anaesthesia, nursing, medical, midwifery, paramedic, physiotherapy, and other allied health students were also included to provide a broader understanding of stress during clinical education. Editorials, conference abstracts, letters to the editor, case reports, and studies unrelated to clinical training were excluded.

The selected articles were critically reviewed and the findings were synthesized narratively. The evidence was organized into major themes, including sources of stress, effects of stress, assessment tools, coping strategies, interventions to reduce stress, and research gaps, to provide a comprehensive overview of the current evidence relevant to A&OTT students. Narrative reviews synthesize evidence thematically rather than using the structured selection process of systematic reviews.

3. Clinical Education in Anaesthesia and Operation Theatre Technology

Clinical education is a core component of the Anaesthesia and Operation Theatre Technology (A&OTT) curriculum, enabling students to integrate theoretical knowledge with practical skills in real healthcare settings. Clinical postings expose students to operation theatres, intensive care units (ICUs), emergency departments, labour rooms, post-anaesthesia care units (PACUs), and other perioperative areas, where they actively participate in patient care under the supervision of healthcare professionals. These experiences help students develop technical competence, clinical decision-making, communication skills, professionalism, and teamwork, all of which are essential for safe perioperative practice (15,16,19,25).

The operating theatre provides a unique learning environment that requires students to understand aseptic techniques, infection prevention measures, anaesthesia equipment, patient monitoring, surgical instruments, and multidisciplinary collaboration. During clinical postings, students are expected to observe and assist in various perioperative procedures while maintaining patient safety and professional ethics. Such exposure enhances confidence and prepares students for future clinical responsibilities; however, it also presents several academic and clinical challenges that may contribute to stress (15,16,18,19,28).

Previous studies have reported that a supportive clinical learning environment, effective supervision, and opportunities for hands-on practice positively influence students' learning experiences and professional development. In contrast, inadequate supervision, unfamiliar clinical settings, limited practical exposure, and ineffective communication with healthcare professionals may reduce students' confidence and negatively affect their clinical performance (2,15,18,19,25,27). Therefore, creating a positive and supportive clinical learning environment is essential to facilitate skill development while minimizing stress during clinical education (2,10,16,27).

4. Sources of Stress During Clinical Postings

The literature indicates that stress during clinical postings is multifactorial and arises from several academic, clinical, interpersonal, and organizational factors. Although the nature and intensity of these stressors may differ among students, they collectively influence clinical learning, confidence, and psychological well-being. The following sections discuss the major factors contributing to stress during clinical education based on the available evidence.

4.1 Academic Demands

Academic responsibilities are consistently identified as one of the major sources of stress during clinical education. In addition to attending clinical postings, students are required to complete assignments, maintain logbooks, prepare case presentations, and perform well in practical and theoretical examinations. Balancing these academic responsibilities with demanding clinical schedules often creates time constraints and increases psychological pressure. Evidence from studies involving nursing, operating room, and allied health students indicates that continuous assessments, practical examinations, and academic expectations may negatively affect students' confidence, concentration, and learning outcomes. The simultaneous demands of classroom learning and clinical responsibilities have been associated with increased stress and reduced academic satisfaction, highlighting academic workload as a significant contributor to stress during clinical training (19,23,26,27,28).

4.2 Clinical Learning Environment

The clinical learning environment plays a central role in shaping students' experiences during clinical postings. The operating theatre is a highly specialized setting that requires technical precision, strict adherence to aseptic techniques, effective communication, rapid decision-making, and multidisciplinary teamwork. Students entering this environment for the first time frequently experience anxiety because of unfamiliar equipment, complex surgical procedures, and uncertainty regarding their clinical responsibilities. Studies consistently demonstrate that unfamiliar clinical settings, limited practical experience, and inadequate orientation increase stress among healthcare students during clinical training. Conversely, supportive supervision, constructive feedback, and opportunities for supervised hands-on practice improve students' confidence, encourage active participation, and enhance the overall learning experience. These findings emphasize that the quality of the clinical learning environment has a direct influence on students' stress levels and professional development (15,16,18,19,25,28).

4.3 Communication and Interpersonal Challenges

Effective communication is fundamental to successful clinical learning and safe patient care. During clinical postings, students are expected to communicate confidently with anaesthesiologists, surgeons, nurses, patients, and other healthcare professionals while functioning as members of multidisciplinary teams. However, language barriers, fear of asking questions, uncertainty regarding professional roles, and limited confidence in communicating with senior staff frequently contribute to stress. The literature suggests that positive interpersonal relationships and supportive mentorship significantly improve students' confidence and clinical participation. In contrast, ineffective communication, inadequate

supervision, and limited constructive feedback may reduce students' motivation, hinder learning opportunities, and increase psychological stress during clinical education (15,18,19,22,25).

4.4 Emotional and Psychological Factors

Clinical education exposes students to emotionally demanding situations that may affect their psychological well-being. Caring for critically ill patients, participating in emergency procedures, witnessing unexpected intraoperative complications, and occasionally encountering patient death can be emotionally challenging, particularly for students with limited clinical experience. Fear of making mistakes and concerns regarding patient safety further contribute to anxiety and emotional distress. Several studies have reported that prolonged exposure to these stressors may lead to emotional exhaustion, reduced self-confidence, anxiety, and burnout if adequate emotional support is not available. Early identification of psychological stress and the provision of appropriate support are therefore important for maintaining students' well-being throughout clinical training (13,17,18,19,27).

4.5 Institutional and Organizational Factors

Institutional support plays an important role in determining the quality of students' clinical learning experiences. Factors such as inadequate orientation, insufficient supervision, limited opportunities for supervised practice, lack of timely feedback, long clinical hours, and heavy workloads have been identified as important contributors to stress during clinical postings. These organizational challenges may reduce students' confidence and negatively influence both learning and clinical performance. Evidence suggests that educational institutions can minimize stress by providing structured orientation programmes, supportive faculty mentorship, regular feedback, and adequate learning opportunities throughout clinical training. A positive educational environment not only improves students' confidence but also promotes professional development and successful adaptation to clinical practice (2,10,15,18,19,25).

Overall, the available evidence indicates that stress during clinical postings is influenced by multiple interacting factors rather than a single cause. Academic demands, challenging clinical environments, communication difficulties, emotional experiences, and institutional support collectively shape students' learning experiences and psychological well-being. Recognizing these stressors is essential for developing effective interventions that promote resilience, improve clinical competence, and enhance the quality of clinical education (10,18,19,27).

Table 1. Major Sources of Stress During Clinical Postings Reported in the Literature

Source of Stress	Examples Reported in the Literature
Academic factors	Examinations, assignments, logbooks, case presentations, academic workload
Clinical learning	Unfamiliar operating theatre, ICU, emergency department,

environment	complex procedures, advanced equipment
Knowledge and skills	Limited practical experience, inadequate preparation, fear of making mistakes
Communication and interpersonal factors	Language barriers, interaction with faculty, surgeons, nurses, and patients, lack of constructive feedback
Emotional and psychological factors	Caring for critically ill patients, emergency surgeries, patient death, anxiety, emotional exhaustion
Institutional and organizational factors	Inadequate supervision, lack of orientation, insufficient feedback, limited learning opportunities
Workload	Long clinical hours, physical fatigue, balancing academic and clinical responsibilities

225

226 5. Effects of Stress During Clinical Postings

227 Stress experienced during clinical postings can influence multiple aspects of students'
 228 academic and professional development. The available literature indicates that prolonged or
 229 poorly managed stress affects not only learning outcomes but also clinical competence,
 230 psychological well-being, professional behaviour, and patient safety. Although moderate
 231 levels of stress may improve motivation and learning, excessive or persistent stress has
 232 consistently been associated with adverse educational and psychological outcomes among
 233 healthcare students.

234 5.1 Academic Performance

235 Clinical stress can negatively affect students' academic performance by impairing
 236 concentration, memory, critical thinking, and decision-making abilities. Balancing academic
 237 responsibilities with clinical duties often creates psychological pressure that reduces learning
 238 efficiency and academic satisfaction. Evidence from systematic reviews and observational
 239 studies indicates that high levels of clinical stress are associated with poorer academic
 240 performance, reduced confidence, and lower learning satisfaction among healthcare students
 241 (26,27,28).

242 5.2 Clinical Competence

243 Clinical competence develops through repeated exposure to patient care, supervised practice,
 244 and active participation in clinical activities. However, excessive stress may reduce students'
 245 confidence, impair psychomotor performance, and discourage participation because of fear of
 246 making mistakes. Studies have shown that reducing stress through simulation-based learning
 247 and structured clinical preparation improves students' technical performance, confidence, and
 248 readiness for clinical practice (6,8,9,16,21).

249 5.3 Psychological Health

Persistent clinical stress has significant consequences for students' psychological well-being. Anxiety, emotional distress, sleep disturbances, reduced self-esteem, and depressive symptoms are commonly reported among healthcare students during clinical education. The literature indicates that psychological support programmes and educational interventions can effectively reduce stress and improve students' emotional well-being and resilience during clinical training (2,13,17,18).

5.4 Burnout

Burnout is characterized by emotional exhaustion, depersonalization, and reduced personal accomplishment resulting from prolonged exposure to academic and clinical stress. Students experiencing persistent stress are more likely to develop burnout, which may reduce motivation, learning satisfaction, and professional commitment. Early recognition of stress and timely psychological support are therefore essential to promote resilience and prevent burnout among healthcare students (13,18,27).

5.5 Patient Safety

Clinical stress may indirectly compromise patient safety by affecting students' concentration, communication, clinical judgment, and procedural performance. Increased stress may result in hesitation during clinical procedures, reduced situational awareness, and a greater likelihood of errors. The literature suggests that supportive supervision, effective communication, and simulation-based education improve students' confidence while promoting safer clinical practice and better patient care (8,15,16,18,21).

Overall, the available evidence demonstrates that unmanaged stress can adversely affect students' academic achievement, clinical competence, psychological well-being, professional development, and patient safety. Early identification of stress and implementation of supportive educational strategies are therefore essential for optimizing students' learning experiences and preparing competent healthcare professionals (10,18,19,27)

Table 2. Major Effects of Stress During Clinical Postings Reported in the Literature

Effect	Impact Reported in the Literature
Academic performance	Reduced concentration, memory, learning efficiency, examination performance, and academic satisfaction
Clinical competence	Reduced confidence, hesitation during procedures, impaired psychomotor performance, and limited clinical participation
Psychological health	Anxiety, emotional distress, depression, sleep disturbances, reduced self-esteem, and psychological distress
Burnout	Emotional exhaustion, depersonalization, reduced motivation, and decreased professional commitment
Patient safety	Increased risk of procedural errors, impaired clinical decision-making, reduced situational awareness, and communication difficulties

6. Stress Assessment Tools

Accurate assessment of stress is essential for identifying students who may be at risk of psychological distress during clinical postings and for evaluating the effectiveness of stress-reduction interventions. Various validated instruments have been used to measure perceived stress, anxiety, burnout, and general psychological well-being among healthcare students. The selection of an assessment tool depends on the study objectives and the specific psychological outcomes being investigated.

The Perceived Stress Scale (PSS) is one of the most frequently used instruments for measuring perceived stress. It evaluates the extent to which individuals perceive situations in their lives as stressful over the previous month. Owing to its simplicity, good reliability, and ease of administration, the PSS has been widely applied in studies involving nursing, medical, and allied health students during clinical education (21,22).

Another commonly used instrument is the Depression, Anxiety and Stress Scale-21 (DASS-21), which simultaneously assesses symptoms of depression, anxiety, and stress. Compared with the PSS, DASS-21 provides a broader evaluation of psychological health and has been extensively used to assess emotional distress among healthcare students during clinical training (23,24).

Burnout associated with prolonged academic and clinical stress is frequently assessed using the Maslach Burnout Inventory (MBI). This instrument evaluates three dimensions of burnout: emotional exhaustion, depersonalization, and reduced personal accomplishment. It has been widely used to investigate the relationship between stress, emotional well-being, and professional development among healthcare students (25).

General psychological well-being is commonly assessed using the General Health Questionnaire (GHQ), particularly the GHQ-12 and GHQ-28 versions. The GHQ is a validated screening instrument designed to identify symptoms of psychological distress and has been used extensively to evaluate the impact of clinical education on students' mental health (26).

Overall, the literature demonstrates that the PSS, DASS-21, MBI, and GHQ are reliable and validated instruments for assessing stress and related psychological outcomes among healthcare students. Their application enables early identification of students requiring academic or psychological support and facilitates the evaluation of interventions aimed at reducing stress during clinical education.

Table 3. Commonly Used Instruments for Assessing Stress Among Healthcare Students

Assessment Tool	Measures	Common Application
Perceived Stress Scale	Perceived stress	Measures overall stress levels

(PSS)		during clinical postings
Depression, Anxiety and Stress Scale-21 (DASS-21)	Depression, anxiety, and stress	Assesses psychological distress during clinical education
Maslach Burnout Inventory (MBI)	Emotional exhaustion, depersonalization, and personal accomplishment	Evaluates burnout associated with prolonged academic and clinical stress
General Health Questionnaire (GHQ-12/GHQ-28)	General psychological well-being	Screens for psychological distress and mental health problems

311

312 7. Coping Strategies

313 Healthcare students adopt a variety of coping strategies to manage stress during clinical
314 postings. The literature suggests that effective coping not only reduces psychological distress
315 but also improves confidence, resilience, and overall clinical performance. Coping strategies
316 may be problem-focused, such as seeking guidance from faculty members and improving
317 time management, or emotion-focused, including relaxation techniques, physical activity, and
318 seeking emotional support from peers and family.

319 Several studies have reported that peer support and positive interactions with clinical
320 instructors play an important role in reducing stress during clinical education. Students who
321 receive constructive feedback and encouragement from supervisors are more likely to
322 participate actively in clinical learning and develop greater self-confidence (2,18,19,22,25).
323 Similarly, maintaining healthy lifestyle habits, engaging in regular physical exercise, and
324 practising mindfulness or relaxation techniques have been associated with lower stress levels
325 and better psychological well-being among healthcare students (2,17,18).

326 Simulation-based learning has also emerged as an effective coping strategy by allowing
327 students to practise clinical skills in a safe and controlled environment before entering real
328 clinical settings. Evidence indicates that simulation improves clinical confidence, reduces
329 anxiety, and enhances preparedness for clinical practice, thereby minimizing stress during
330 clinical postings (6,8,9,16,21).

331 8. Interventions to Reduce Stress

332 Educational institutions play a crucial role in reducing stress by creating supportive clinical
333 learning environments. The literature highlights that structured orientation programmes,
334 effective mentorship, and regular supervision contribute significantly to students' confidence
335 and adaptation during clinical postings. Early orientation to operation theatre protocols,
336 anaesthesia equipment, and clinical expectations can reduce uncertainty and improve
337 students' readiness for clinical practice (15,19,21,25).

Simulation-based education has consistently demonstrated positive outcomes in reducing clinical stress. Randomized controlled trials have shown that virtual reality-based orientation, scenario-based simulation, and high-fidelity simulation improve students' confidence while decreasing anxiety before clinical exposure (6,8,9,16,21). In addition, stress management programmes incorporating relaxation techniques, mindfulness training, and cognitive behavioural approaches have been found to reduce perceived stress and enhance emotional well-being among healthcare students (2,10,17,18).

Institutional support through accessible counselling services, regular feedback, peer mentoring, and faculty development programmes further contributes to a positive clinical learning environment. Such interventions not only improve students' psychological well-being but also promote professional growth and clinical competence (2,10,15,18,19,25).

Table 4. Interventions to Reduce Stress During Clinical Postings

Intervention	Description	Reported Outcomes
Structured orientation programmes	Orientation to the operating theatre, anaesthesia equipment, clinical protocols, and student responsibilities before clinical postings	Reduced uncertainty, improved preparedness, and facilitated adaptation to the clinical environment
Simulation-based education	High-fidelity simulation, scenario-based learning, and virtual reality training before clinical exposure	Reduced stress and anxiety, improved clinical confidence, technical skills, and decision-making ability
Stress management programmes	Programmes incorporating relaxation techniques, mindfulness, and cognitive behavioural strategies	Decreased perceived stress, enhanced emotional well-being, and improved coping abilities
Faculty mentorship and clinical supervision	Regular supervision, constructive feedback, and guidance from faculty and clinical instructors	Improved confidence, clinical competence, and learning experience
Peer mentoring and support	Peer-assisted learning, emotional support, and collaborative learning activities	Reduced feelings of isolation, promoted resilience, and improved adaptation to clinical training
Counselling and psychological support	Access to counselling services and mental health resources	Improved psychological well-being and reduced symptoms of stress and anxiety

9. Research Gaps

Although stress among healthcare students has been widely investigated, research specifically focusing on **Anaesthesia and Operation Theatre Technology (A&OTT) students** remains limited. Most published studies have been conducted among nursing, medical, or operating room students, and their findings are often extrapolated to A&OTT education. Consequently, evidence specific to the unique clinical responsibilities and learning experiences of A&OTT students is lacking.

10. Future Directions

Future studies should focus on identifying stressors unique to Anaesthesia and Operation Theatre Technology students and evaluating interventions that improve both psychological well-being and clinical competence. Multicentre studies involving larger and more diverse student populations are needed to enhance the generalizability of findings. Longitudinal research would provide valuable insights into changes in stress levels throughout different stages of clinical education.

Educational institutions should also explore innovative teaching methods, including simulation-based learning, virtual reality, structured mentorship programmes, and digital mental health interventions. Integrating stress management training into the A&OTT curriculum may help students develop resilience, improve coping skills, and enhance preparedness for clinical practice. Further research evaluating the effectiveness of these strategies will support the development of evidence-based educational policies.

11. Limitations

This review has some limitations. Because of the limited number of studies specifically investigating stress among Anaesthesia and Operation Theatre Technology (A&OTT) students, evidence from operating room, nursing, medical, midwifery, paramedic, physiotherapy, and other allied health students was included to provide a broader understanding of the topic. Therefore, some findings may not fully represent the unique clinical experiences and stressors of A&OTT students. In addition, the included studies varied in their study designs, sample characteristics, clinical settings, and stress assessment instruments, making direct comparison of findings challenging. As this is a narrative review, a formal assessment of the methodological quality and risk of bias of the included studies was not performed. Consequently, the findings should be interpreted in the context of the available evidence.

12. Summary Points

- Clinical postings are an essential component of Anaesthesia and Operation Theatre Technology (A&OTT) education but are frequently associated with moderate to high levels of stress.

- Stress during clinical training is multifactorial, arising from academic demands, unfamiliar clinical environments, communication challenges, emotional experiences, and institutional factors.
- Excessive stress can negatively affect academic performance, clinical competence, psychological well-being, increase the risk of burnout, and potentially influence patient safety.
- Validated instruments such as the Perceived Stress Scale (PSS), Depression Anxiety Stress Scale-21 (DASS-21), Maslach Burnout Inventory (MBI), and General Health Questionnaire (GHQ) are commonly used to assess stress and related psychological outcomes.
- Individual coping strategies, including time management, self-directed learning, physical activity, mindfulness, and seeking social support, help students manage clinical stress effectively.
- Faculty mentorship, peer support, constructive feedback, and a positive clinical learning environment play a significant role in improving students' confidence and reducing stress.
- Educational interventions such as simulation-based learning, structured orientation programmes, stress management training, and counselling services have demonstrated positive outcomes in reducing stress and enhancing clinical preparedness.
- Current evidence specific to A&OTT students remains limited, with most available studies conducted among nursing and medical students.
- Future research should focus on multicentre, longitudinal studies evaluating stress, coping strategies, and targeted interventions specifically for A&OTT students.
- Developing supportive, student-centred clinical learning environments is essential for improving students' well-being, professional competence, and quality of patient care.

13. Conclusion

Clinical posting is an indispensable component of Anaesthesia and Operation Theatre Technology education, providing students with essential opportunities to develop clinical knowledge, technical skills, and professional competence. However, the available evidence indicates that clinical training is frequently associated with stress arising from academic demands, challenging clinical environments, communication difficulties, emotional experiences, and limited institutional support.

If left unaddressed, excessive stress may adversely affect students' academic performance, clinical competence, psychological well-being, and ultimately patient care. The literature consistently demonstrates that supportive clinical supervision, simulation-based education, effective communication, stress management programmes, and faculty mentorship can reduce stress and improve students' confidence and learning experiences.

Although evidence from nursing, medical, and operating room education provides valuable insights, research specifically involving Anaesthesia and Operation Theatre Technology students remains limited. Future well-designed studies are needed to better understand the unique challenges faced by this group and to develop targeted interventions. Creating a supportive and student-centred clinical learning environment will be essential for promoting resilience, improving educational outcomes, and preparing competent healthcare professionals.

REFERENCES

1. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav.* 1983;24(4):385-396.
2. Emel Gülnar, Aşık E, Özveren H. The effect of a stress management program on first-year nursing students' clinical stress: A randomized controlled experimental study. *Nurse Educ Today.* 2024;136:106131.
3. Goldberg DP, Williams P. *A User's Guide to the General Health Questionnaire.* Windsor: NFER-Nelson; 1988.
4. Hebel K, Steliga A, Lewandowska K, et al. Simulated learning, real emotions: The impact of simulation-based education on nursing students' stress levels during objective structured clinical examination: A longitudinal observational cohort study. *Nurs Rep.* 2025.
5. Hernández Ortega R, González Pascual JL, Fernández Araque AM. Impact of an intervention program with reinforcement on nursing students' stress and anxiety levels in their clinical practices. *Nurse Educ Pract.* 2021;55:103179.
6. Hung CC, Kao HS, Liu HC, Liang HF, Chu TP, Lee BO. Effects of simulation-based learning on nursing students' perceived competence, self-efficacy, and learning satisfaction: A repeat measurement method. *Nurse Educ Today.* 2021;97:104725.
7. Judd BK, Alison JA, Waters D, Gordon CJ. Comparison of psychophysiological stress in physiotherapy students undertaking simulation and hospital-based clinical education. *Simul Healthc.* 2016;11(4):271-277.
8. Kaldheim HKA, Bergland Å, Ølnes MA, Hofso K, Dihle A, Creutzfeldt J, Zhang C, Steindal SA. Use of simulation-based learning among perioperative nurses and students: A scoping review. *Nurse Educ Today.* 2019;73:31-37.
9. Kim HY. Validating the effectiveness of a VR-based and haptic technology-based operating room simulation program among nursing students. *Comput Inform Nurs.* 2025.
10. Ko KY, Montayre JR, Chiu PL, Lai TKH, Chan K, Chan HHT. Effects of interventions to reduce clinical placement-related psychological distress among nursing students: A systematic review with meta-analysis. *Int J Nurs Stud Adv.* 2025.
11. Lee EH. Review of the psychometric evidence of the Perceived Stress Scale. *Asian Nurs Res.* 2012;6(4):121-127.
12. Lovibond SH, Lovibond PF. *Manual for the Depression Anxiety Stress Scales.* 2nd ed. Sydney: Psychology Foundation; 1995.

13. Martín-Parrilla MÁ, Durán-Gómez N, Montanero-Fernández J, Casas-Méndez C, López-Jurado CF, Cáceres MC. Assessing emotional intelligence and reducing burnout in nursing students through simulation-based learning experiences: An experimental design. *Int J Ment Health Nurs*. 2025;34(5):e70151.
14. Maslach C, Jackson SE, Leiter MP. *Maslach Burnout Inventory Manual*. 4th ed. Menlo Park (CA): Mind Garden; 2018.
15. Meyer R, Van Schalkwyk SC, Prakaschandra R. The operating room as a clinical learning environment: An exploratory study. *Nurse Educ Pract*. 2016;18:60-72.
16. Mohammadi G, Tourdeh M, Ebrahimian A. Effect of simulation-based training method on the psychological health promotion in operating room students during the educational internship. *J Educ Health Promot*. 2019;8:172.
17. Nebhinani N, Kuppili PP, Mamta. Feasibility and effectiveness of stress management skill training in medical students. *Med J Armed Forces India*. 2024;80(2):140-144.
18. Ng R, Chahine S, Lanting B, Howard J. Unpacking the literature on stress and resiliency: A narrative review focused on learners in the operating room. *J Surg Educ*. 2019;76(2):343-353.
19. Norouzi N, Imani B. Clinical education stressors in operating room students: A qualitative study. *Invest Educ Enferm*. 2021;39(1):e08.
20. Osman A, Wong JL, Bagge CL, Freedenthal S, Gutierrez PM, Lozano G. The Depression Anxiety Stress Scales-21 (DASS-21): Further examination of dimensions, scale reliability, and correlates. *J Clin Psychol*. 2012;68(12):1322-1338.
21. Parlak EA, Ayhan H, Erer I, Iyigun E. The effect of virtual reality-based operating room tour on nursing students' anxiety, stress and self-confidence levels in an operating room setting: A randomized controlled trial. *Nurse Educ Pract*. 2025;85:104380.
22. Pinks D, Warren-James M, Katsikitis M. Does a peer social support group intervention using the CARES skills framework improve emotional expression and emotion-focused coping in paramedic students? *Australas Emerg Care*. 2021;24(4):308-313.
23. Poorheidari M, Delvarzian ZM, Yahyae S, Montazeri AS. Study of the stressful experiences of midwifery students during clinical education in the labour room. *Res Med Educ*. 2018;9(4):58-66.
24. Ross S, Cleland J, Macleod MJ. Stress, debt and undergraduate medical student performance. *Med Educ*. 2006;40(6):584-589.
25. Saxton R, Nauser J. Students' experiences of clinical immersion in operating room and emergency department. *Nurse Educ Pract*. 2020;43:102709.
26. Sohail N. Stress and academic performance among medical students. *J Coll Physicians Surg Pak*. 2013;23(1):67-71.
27. Welch SR. Clinical stress and clinical performance in prelicensure nursing students: A systematic review. *J Nurs Educ*. 2023;62(1):36-41.
28. Yildiz Findik U, Ozbas A, Cavdar I, Yildizeli Topcu S, Onler E. Assessment of nursing students' stress levels and coping strategies in operating room practice. *Nurse Educ Pract*. 2015;15(3):192-195.