

1
2
3 **QUATERNARY VOLCANIC ERUPTION TOBA ASH FALL ITS IMPACTS ON**
4 **ENVIRONMENT OF LATE PLEISTOCENE HOMININES IN INDIAN**
5 **SUBCONTINENT WITH SPECIAL REFERENCE TO NARMADA VALLEY.**
6 **INDIA.**

7
8 **ABSTRACT :**
9

10 The 74,000 year-old super eruption of the Toba volcano, located in northern Sumatra, is
11 recognized as one of Earth's largest known eruptions and was certainly the largest of the
12 Quaternary period (Smith and Bailey, 1968). It is presumed that it have led to both global
13 climatic and environmental deterioration and had a impact on decimation of modern
14 human populations (Rampino et al., 1988; Rampino and Self, 1992, 1993a) (Rampino
15 and Self, 1993b; Ambrose, 1998, 2003a, 2003b; Rampino and Ambrose, 2000).
16

17 However, the severity of Toba's impact on climate and hominines has been contested
18 and debated by scientist (Oppenheimer, 2002; Gathorne-Hardy and Harcourt-Smith,
19 2003). Geological, paleontological and archaeological evidence from the Indian
20 subcontinent provides an excellent opportunity to address these issues. The intensity and
21 scale of the Toba super eruption was multi dimensional and its column of eruption was
22 so vast that it forma canopy of volcanic matrix led to the deposition of a blanket of
23 volcanic ash across the continents and river valleys over India, Malaysia, the Indian
24 Ocean, and the Arabian and South China Seas. Resulting persistent terrestrial tephra
25 deposits have been documented in a number of river valleys throughout India. (Khan et.al
26 1991 Acharyya and Basu, 1993; Shane et al., 1995; Westgate et al., 1998). The
27 occurrences of volcanic ash have been located in Narmada, Tapti, Purna Son valleys.
28 These occurrences are associated and preserved with archaeological, paleontological sites
29 in these valleys.
30

31 The present work is first attempt to collect condense the data on various aspects of
32 eruption of the Tuba volcano, its impact on environment on population with special
33 reference to Narmada valley. It is also attempted to study the relationship between Tuba
34 ash and archaeology from Narmada valley and adjoining area. Besides attempts were
35 made to study the nature of occurrences of ash bed its composition chemical and physical
36 characteristics to assess paleo- environment. In Narmada valley and Indian sub continent
37 the Environment and climate during late Pleistocene has been significantly affected after
38 Toba eruption, according to Rampino et al. (1988) the size of Toba at 74 ka could have
39 induced a volcanic winter, similar to predicted nuclear winter scenarios, as modeled by
40 Turco et al. (1983, 1990). The injection of vast amounts of gaseous aerosols and volcanic
41 dust and matrix into the atmosphere, which follow large volcanic eruptions, is predicted
42 to have detrimental and decisive consequences for changes of global climate. Past
43 historical eruptions, such as Tambora in 1815 (Stothers, 1984) and Pinatubo in 1991
44 (McCormick et al., 1995), have provided evidence of post-eruption climatic deterioration
45 and climatic. With the eruption of Toba having been far larger than both of these

historical eruptions, its consequences are therefore assumed to have been far more devastating.

The imprints of devastating impacts in the area of study and specific on human population are matter of scientific concern and yet to be investigated. The studies conducted of Ash bed associated with Quaternary deposits of Narmada Valley consist of sediments of two domains viz. deposits of interglacial domain (Boulder conglomerate-Hathnora formation) and fluvial deposit of paleo-domain of Narmada. The boulder conglomerate is fossiliferous horizon of Narmada and has yielded skull cap of *Homo erectus* Sonakia (1984). (Khan & Solaria (1991). It is marker horizon and represents interglacial phase in the history of Quaternary sedimentation in Narmada Valley. The occurrence of ash bed reported from Quaternary sediments of Narmada are associated with two horizons of ash beds of middle and upper Pleistocene age (Khan et.al.1991). These ash beds are designated as NAB-I and NAB-II consisting of five layers designated as (L1 to L5) in ascending antiquity in the valley. The Ash bed NAB-I (L1- to L-3) is associated with upper gritty units of boulder conglomerate (Hathnora formation) and is identified at an elevation of about 290m. above m.s.l. The Ash bed NAB-II (L-4 to L5) is associated with upper units of clay silt deposits of paleo-domain of Narmada (Shahganj formation) and is identified at an elevation of 310m. above m.s.l. The study of grain morphology of glass matrix, their relation with other minerals shape, size, texture of lithic fragments and association of other ashy sediments of pyroclastic origin suggest that sediments were brought from distant source by Aeolian agencies, during the different phases of sedimentation in Narmada Valley. It is observed that in stratigraphic column two horizons of Ash bed occurred at the vertical distance of about 20 m. which indicate that there were two phases of settling volcanic ash in Narmada valley and Indian subcontinent as a whole with significant time break. The element of time break is in terms of sedimentation perceptible appears to be related with global climatic changes induced by the super volcanic eruption. These ash beds are used as tool in correlation of different quaternary deposits and archeological sites in the valley. The occurrences of two horizons of Ash bed and their deposition by settling of volcanic dust with time gaps suggest induced and defused atmospheric conditions in Narmada valley for a very long time after volcanic eruption.

The atmosphere contaminated by volcanic ash volcanic matrix and huge amount of gases and dust particles had definitely affected the atmosphere and climate had have led to temporarily darkened skies in the Narmada valley and Indian subcontinent for very long time.

The study of grain morphology of glass matrix, their relation with other minerals shape, size, texture of lithic fragments and association of other ashy sediments of pyroclastic origin suggest that sediments were brought from distant source by Aeolian agencies, during the different phases of sedimentation in Narmada Valley.

It is commonly argued that if Toba was truly devastating for *H. sapiens* then comparable bottlenecks should also be seen in many other species.

Introduction

The Quaternary tract of Narmada basin covers an area of about 12950 sq. km starting from west of Jabalpur (23°07'790530) to east of Handia (22° 29'; 76° 58') for a distance of about 320 km. It is found to be ideal locus of Quaternary sedimentation in Central India as witness by multi-cyclic sequence of Quaternary terraces in the valley. The total average estimated thickness of Quaternary sediments in the central sector of Narmada is about 325 m. where the level of Ash bed occurrence has been identified between the depths of 75-83 m of Quaternary column in unified Quaternary Platform.

The skull cap of *Homo erectus* (Sonakia1984) and other fauna recorded along with calc-nodules near village Hathnora (22° 52" N; 77° 52" E) in fossiliferous boulder conglomerate; named as Hathnora formation Khan & Sonakia (1992). It is found to be associated with volcanic Ash bed of Quaternary age in the area around Hathnora, and upstream Khan et.al. (1991). The two levels of horizons of Ash bed identified are designated as NAB-I and NAB-II in ascending antiquity in the valley. The Ash bed NAB-1 is associated lower litho units of boulder conglomerate which is well preserved and persistent where as NAB-II is associated with younger deposits. The NAB-1 contains three micro layer (L-1 to -L3) and NB-II two micro layers (L-4 to L-5) in increasing antiquity. The area around Hathnora and in Central Narmada valley possesses rich Paleolithic occurrences about and 300 Lower Paleolithic occurrences have been reported in different contexts. The present work deals with these assemblages and associated raw material sources in valley .These rock types were used as raw material by various hominines populations. The area appears to be thickly inhabited and influence by human activity which was adversely affected by severe Toba volcanic eruption and hominines were forced to be migrated in struggle of survival to the safe places. The nearest safe place was Bhimbetka about twenty kilometers in the north located in obsequent slope of Vindhyan rocks of Bhopal inliers which possess about 700 rock shelters of different size and dimension. The study of human activity in the area around Bhimbetka indicates the hominines populations lived around 100000 years.

The occurrences of volcanic ash have been located in Narmada, Tapi, Purna Son valleys. These occurrences are associated and preserved with archaeological, paleontological sites in these valleys. The present work is first attempt to collect and condense the data on various aspects of eruption of the Toba volcano, its impact on environment on population with special reference to Narmada valley. It is also attempted to study the relationship between Toba ash and archaeology from Narmada valley and adjoining area. Besides attempts were made to study the nature of occurrences of ash bed its composition chemical and physical characteristics to assess pale environment. This data can be combined with additional evidence from a wider context in order to give a broader understanding of the regional impacts of the eruption on Narmada valley and Indian subcontinent as a whole.

The 74,000 year-old super eruption of the Toba volcano, located in northern Sumatra, is recognized as one of Earth's largest known eruptions and was certainly the largest of the

Quaternary period (Smith and Bailey, 1968). It is presumed that it have led to both global climatic and environmental deterioration and had a impact on decimation of modern human populations (Rampino et al., 1988; Rampino and Self, 1992, 1993a) (Rampino and Self, 1993b; Ambrose, 1998, 2003a, 2003b; Rampino and Ambrose, 2000). However, the severity of Toba's impact on climate and hominines has been contested and debated by scientist (Oppenheimer, 2002; Gathorne-Hardy and Harcourt-Smith, 2003). Geological, paleontological and archaeological evidence from the Indian subcontinent provides an excellent opportunity to address these issues. The intensity and scale of the Toba super eruption was multidimensional and its coloumn of eruption was so vast that it forma canopy of volcanic matrix led to the deposition of a blanket of volcanic ash across the continents and river valleys over India, Malaysia, the Indian Ocean, and the Arabian and South China Seas. Resulting persistent terrestrial tephra deposits have been documented in a number of river valleys throughout India. (Khan et.al 1991 Acharyya and Basu, 1993; Shane et al., 1995; Westgate et al., 1998).

4.2.00 Pleistocene Sedimentation in Narmada Rift Valley

The Narmada Rift valley formed a linear trench in the middle of Indian subcontinent was an ideal loci for accumulation of sediments .The rift trench is intruded by the dolerite and other mafic and siliceous dykes and sills along lineaments in different phases of tectonic deformation. The Quaternary sedimentation incepting from glacial activity, followed by fluvio-glacial, lacustrine and fluvial phase within the rinsing and sinking environment, block faulting and segmental and linear displacement and dislocation, uplifting and isolated domal up-lift, Neogene rifting and Quaternary sedimentation and rift-bound Pliocene-Pleistocene rifting and volcanic activity specifically during glacial and fluvio-glacial phase are major component of the Quaternary period and tectonic processes of the Narmada Rift System and form the base of quaternary deposits . The Quaternary sedimentation was triggered by tectonic activities / up lift and climatic changes. The provenance for these sediments is the weathering products of eroding pre-Cambrian, meta-sediments, sedimentary and volcanic rocks along the watershed upland, rift escarpments and shoulders; faulted and uplifted blocks, volcanic fissure zones, and plateaus within and outside the rift. The Narmada Rift System, bounded by adjacent plateaus rising 300–700 m above the rift floor, consists of number symmetrical and symmetrical faulted blocks, escarpment, rock cut terraces, rock floors and segments of micro half grabben. Although rift-related basins started to form during the late Oligocene to early Miocene times, the Narmada Rifts were fully defined by middle to late Miocene time.

The Narmada rift system basin platform provided a unique setting for dynamic ecosystems that were characterized by Rift-related subsidence and coeval sedimentation also created an ideal loci of Quaternary sedimentation and environment for the accumulation of sediments volcanic fabrics sediments, burial,

digenesis, and preservation of organic remains. Because rifts formed after widespread Quaternary sedimentation occurred and voluminous sediments in the rift basins were accumulated by glacial activity consequential upon the lowering of temperature and climatic changes in the region.

The Miocene -Pliocene–Pleistocene lake deposit of Katni on the eastern rift shoulder was created by faulting, topographic control, or isostatic depression similar to that of other Rift system.

Abundant faunal remains in rift settings, including hominids Sonakia (1984), suggest that the Narmada Rift System created productive ecosystems during Pliocene–Pleistocene time. The volcanic rocks within the fossiliferous sediments provide temporal information for calibrating and sequencing hominid and other faunal evolution. The detailed geological sediment logical geochemical of interbedded tephra ,Quartz grain morphology of sediments of quaternary strata and palo-sole of and geochronological studies of from the different localities for establishing accurate biostratigraphic and lithostratigraphic data, sedimentation rates, and paleoenvironmental and tectonic histories of different sediment columns in area along the rift system. Interbedded volcanic rocks allow determination of the time of rifting, the beginning of sedimentation, sedimentation rates, and the oscillation from glacial, fluvio-glacial lacustrine to fluvial environments. The cyclic environmental transitions recorded in the sedimentary sequences of the rift basins are caused by tectonic activities (uplift and subsidence), changes in relief, and climatic variations. The climatic changes in uplift, topographic and landscape features, coupled with block faulting, rinsing and sinking platform, created basins for the accumulations of thick lacustrine and fluvial sediments sequences with terrestrial and aquatic fossils. The sequential change in the sediment facies from finely bedded lacustrine deposits to fluvial sediments are commonly noted in the sedimentary sequences and reflect environmental and tectonic changes that can be temporally determined. Moreover, regional correlation based on the chemistry and geochronology of interbedded tephra has made it possible to establish accurate stratigraphic relations that are useful for paleo- environment reconstruction and evolutionary studies of fossil remains in the Narmada rift valley Khan et.al. (2012). Regional tephra correlation is being used increasingly to link sites together, and has already established that similar tephra layers are known from other parts of rift valley, as well as from other basin and peninsular India Achariya,(1995), Khan (1992) Khan et.al. (2012) Tiwari (1996). There is a great potential for further correlation of tephra in the Rift System and marine sediments in the Arabian Sea. The Arabian Sea has a continuous record of deposition that extends to at least 7 million years. The Quaternary sediments interbedded with tephra with within the age range of the ODP Ocean Drilling Program 721/722 stratigraphic sections of the Arabian Sea are also present within the rift floor and the western rift margin of the region. The chemical and chronological correlations of ash beds within the rift sequences of have been made with ashes described in marine sections. Detailed correlations based on orbitally calibrated time scales of pale magnetic stratigraphy Rao () within Quaternary sediments of rift deposits will provide ties to establish

global climate changes based on the terrestrial and marine sediments of the rift system.

Moreover, because of tephra layers in sedimentary basins of different geologic periods, processes such as faulting, rifting, sedimentation and diagenesis, impact of climatic changes, age of fossils, nature and acquisition of archeological implements, and the origin, distribution, and functional significance of early hominid artifact assemblages can be deciphered. However, evidence of the effects of tectonics on fauna and flora are distinct and its signatures on dislocation and concealing of fossiliferous horizons are uncontrolled and ill defined in the ecosystem in the valley during the Pliocene–Pleistocene periods. The boulder conglomerate which yielded the skull cap of *Homo erectus* in Narmada rift from Hathnora Sonakia (1984) remained only discovery of hominid fossil in last two and half decade due inconsistency concealed nature of fossiliferous horizon due faulting, dislocation and subsidence of Quaternary blanket of Narmada rift system as such researcher and scientist failed to add any further knowledge to hominid discovery any further.

The Narmada Rift System consists of symmetrical basins that have been evolved in different stages of tectonism. The 100–120 km-wide and 1300 km long rift bounded by Satpura in south and Vindhyan in north constitutes conspicuous ENE–WSW to E–W rift basin zone is filled with Pliocene–Pleistocene sediments, whereas some of them contain Miocene sedimentary deposits. Most of the sedimentary sequences contain faunal and floral remains including hominid species. Most of the basin-fill sediments were derived from topographically elevated rocks that are present within and outside the rift basins. Lava flows and tephra are interbedded with the fossiliferous sediments clastic sediments derived from the provinces of sediment mostly from crystalline basement volcanic, sedimentary, meta basic and sedimentary rocks, aided in the cementation and preservation of organic remains by providing secondary minerals released during alteration in a burial environment. Quick burial minimized the effect of reburials taphonomic processes. Moreover, chemical constituents released by the alter have provided critical temporal and spatial information without which the study of hominid evolution and pale-environmental reconstruction in the Rift System would have been impossible.

Moreover, because of tephra layers in sedimentary basins of different geologic periods, processes such as faulting, rifting, sedimentation and diagenesis, impact of climatic changes, age of fossils, nature and acquisition of archeological implements, and the origin, distribution, and functional significance of early hominid artifact assemblages can be deciphered. However, evidence of the effects of tectonics on fauna and flora are distinct and its signatures on dislocation and concealing of fossiliferous horizons are uncontrolled and ill defined and the ecosystem in the rift system during the Pliocene–Pleistocene periods is not clear. The boulder conglomerate which yielded the skull cap of *Homo erectus* in Narmada rift from Hathnora remained only discovery of hominid fossil in last two and half decade due inconsistency concealed nature of fossiliferous horizon in Narmada valley. Historical or modern analogs illustrate the potential of the regional and sometimes

global effects of such major silica eruptions in the geologic past , sedimentation rates, and the oscillation from lacustrine to fluvial environments. The cyclic environmental transitions recorded in the sedimentary sequences of the rift basins are caused by tectonic activities (uplift and subsidence), changes in relief, and climatic variations. Changes in topographic features, coupled with volcanic damming, created basins for the accumulations of thick glacial, fluvio-glacial lacustrine and fluvial sequences with terrestrial and aquatic fossils. Changes from finely bedded lacustrine deposits to fluvial sediments are commonly noted in the sedimentary sequences and reflect environmental and tectonic changes that can be temporally determined. Moreover, regional correlation based on the chemistry and geochronology of interbedded tephra has made it possible to establish accurate stratigraphic relations that are useful for pale environment reconstruction and evolutionary studies of fossil remains in the rift valleys across India sub continent. Regional tephra correlation is being used increasingly to link sites together, and has already established that similar tephra layers are known in the Rift system.

The Boulder Bed and Boulder conglomerate which form the base of Quaternary sediments in Central sector of Narmada Rift Valley are not exposed in the lower Narmada valley. These deposits are concealed under the sediments of lacustrine and fluvial deposits in the valley. The presence of these deposits is confirmed by study of bore hole data and logs of State and Federal agencies drill in lower Narmada valley under various projects.

The glacial and fluvio-glacial deposits of Narmada unconformably overlie the Vindhyan and the basaltic Deccan Trap rocks. The sediments consist of a Hetero-heterogeneous assemblage of sub-angular to angular, sub-rounded, unsorted, stratified rock fragments ranging from boulders to small pebbles, predominantly of quartzite, gneiss, sandstone, basalt, jasper, chert, gneiss, sandstone, basalt, chert, altered feldspar, ferruginous nodules, in a matrix of very coarse to very fine-sand, silt and clay. These sediments are highly angular, generally poorly sorted and isotropically imbricated. Fine sediments comprise of reddish grayish and greenish sand with appreciable amount of mica flakes, altered feldspar, brick-red and buff silt, greenish-brown silt and clay, and greenish, reddish and dark maroon hard and plastic clay. These fine sediments contain fairly good amount of ferruginous material, quartz, mica flakes and altered feldspar grain. Through these sediments are similar in composition to the other deposits of Narmada Valley, exhibit entirely different sedimentary pattern, sediment characters and mineral composition. These rock clastics are largely angular, very poorly sorted and demonstrate isotropic imbrications pattern in the valley. The sediments of glacial domain of Narmada occur between and average elevation of about 20-265 m above m.s.l. and were deposited in glacial environments during Pleistocene time. These deposits are concealed under boulder conglomerate in the valley.

The Vindhyan Group of rocks and Deccan Trap in the Central sector of the Narmada Valley form the basement for the Quaternary deposits. The conglomerate bed that constituted the fossiliferous horizon of Narmada is sandwiched between older Alluvium and the glacio-fluvial boulder bed. This conglomerate bed is a very persistent marked horizon indicating a distance phase of sedimentation in the Narmada Valley. It is exposed in the bluff/scrap of Narmada around Sardar Nager, Hathnora, Surajgarh, Budhni, Hoshangabad, Khoksa, Tigharia, Demawar and Bhariya-Ghat at the base of terraces NT₂ (Khan, 1984). The measured thickness of the exposed boulder conglomerate varies from 1.5 to 6.5m but average 5m.

The boulder conglomerate predominantly consists of sub-rounded to well rounded boulder, cobble and pebble of quartzite, gneiss, sandstone, basalt, agate, jasper, chert, chalcedony tightly cemented in a matrix of sand and silt. The finer sediments include different grade of sand and silt, brown and maroon in color often laminated and cross laminated. The boulder conglomerate is fossil and skull cap of early man *Homo erectus* (Sonakia, 1984)

The boulder conglomerate consists of three sub-litho units; each sub-unit characterized by distinct rock fragment shape, size, lithological abundance and allied sediment characters. The sub-units are composed of variable assemblage of quartzite, gneiss, basalt, sandstone, agate, jasper, chalcedony, chert, sand and silt (Khan1992) These sub-litho units display facies variation in the valley and upper units grades into gritty sandstone upstream of Hathnora

The basal unit of boulder conglomerate and is characterized by sub-rounded to rounded boulders cobbles, pebble of quartzite (38.50%), gneiss (18.50%) sandstone (14.50%), basalt (6.50%), agate and other rock fragments (3.50%). These clastic are tightly cemented in a reddish brown and maroon sand and silt which exhibit lamination, cross-lamination and cut and fill feature. It is capped by a pebbly layer containing vertebrate fossils and a human skull of early man (Sonakia, 1984). The average measured thickness of this layer is about 2.25m.

The middle sub-unit of boulder conglomerate characterized by an abundance of spherical, discoidal and bladed cobbles, pebbles of quartzite (31.50%), gneiss (14.00%), jasper (8.50%), sandstone (15.00%), basalt (10.50%), agate (12.50%), chert (6.50%) and other rock fragments chert (1.50%). These coarse clastic are tightly cemented by a deep brown and maroon sand, silt and clay. The finer clastics display sedimentary feature like lamination, cross-bedding load structure, small ripples and, cut and fill feature. Out of this sedimentary structure the cross bed units of boulder conglomerate vary from 5.15° and average about is 8.5°. The cross-beds average is often truncated by a pebbly layer at the top. This sub-unit contains mammalian fossil and some stone implements. The average measured thickness of these sub-units is about.

The upper sub-units of the boulder conglomerate. It consists of sub-rounded to well-rounded pebble of quartz (23.50%), gneiss (11.50%), sandstone (14.50%), basalt (13.50%), agate (15.50%), jasper (11.50%), chert (8.00%) and other rock fragments

(2.00%). These clastics are commented by grey and brownish sand and silt containing appreciable amount of calcareous matrix.

These deposits identified between an average elevations of 245 to 300 m above m.s.l. the basal unit of boulder conglomerate identified is marked at an elevation of about 268 m above m.s.l, exposed on the northern bank of Narmada around Hathnora (22 ° 52" N - 77 ° 52" E) at the depth of about 83 m in stratigraphic column of Quaternary sediments of Narmada.

The boulder conglomerate is of middle Pleistocene age equivalent to Siwalik boulder conglomerate (India), Trini bed of Java (Indonesia) and boulder conglomerate of Tapti Khan (1982). These deposits have yielded skull cap of early man, *Homo erectus Narmadensis* along with other mammalian fossils Sonakia, (1984). The Ash bed of Quaternary age is recorded associated with these deposits around Timrawan upstream of Hathnora is of aeolian nature and perhaps indicates volcanic activity during middle Pleistocene time.

The sediments of paleo-domain of Narmada conformably overlie the boulder conglomerate and represent the flood-plain fluvial facies of the Narmada. The sediments of the facies predominantly consist of clay silt and sand, discontinuous nodules and plates. The beds are horizontal, exhibit upward fining sequence typical of fluvial deposits. This domain may be divided into three formations based on lithology, sediment assemblage, shape and size of rock clastics, relative disposition and diagnostic sedimentary characteristics. These formations are, viz. (i) Shohagpur, (ii) Shahganj, and (iii) Hoshangabad Formations respectively. These formations represent the sediments the complete sequence of Narmada deposited in channel and flood plain environments during Upper Pleistocene times. The lowest Sohagpur Formation is named after Sohagpur town. The unit occurs along the outer flanks of Narmada Valley bounded by Vindhya range to the north and Satpura to the south. It consists of sediments of paleo-domain of Narmada. It is represented by a thick sequence of clay, silt-sand and rock gravels. The unit is divisible into three sub litho unit. The basal sub-unit is chiefly red and brownish sand, silt, clay containing appreciable amount of cal matrix. The average measured thickness of this sub-unit is about 6.25m. The middle sub-unit consists of yellow and brownish silt, clay with subordinate sand and occasional rock gravel lenses. The average measured thickness of this sub-unit is about 5.50m. The upper sub-units predominantly consist of compact yellow clay, silt and calcareous concretion. The average measured thickness of this sub-unit is about 3.25m.

The Sohagpur formation is often association with discontinues and persistent pebbly horizon containing well sorted polymodal gravel of quartzite, gneiss, basalt, agate, jasper and chart in the matrix of course to fine-sand. The gravel is general discoidal, spherical and exhibits higher indices of sphericity and roundness indicating their derivation from distance and mixed provenance during sedimentation.

The Shahganj formation is forms litho-stratigraphic unit overlying boulder conglomerate and occupies large area in the central part of the valley. It is exposed in the bluff section of Narmada around Narayanpur, Sardarnagar, Hathnora, Shahganj and Hoshangabad. This formation is equivalent to the Shivpur formation described by Khan (1984) from the down stream of Hoshangabad. It consists of sediments of palaeo-domain of Narmada. It is represented by clay, silt, sand accompanied by inconsistent pebbly bed containing quartzite, gneiss, basalt, chart, agate. The average measured thickness of this formation is about 15m. These sediments constitution three units each characterized by distinct lithology, rock classics and diagnostic sediment characters.

The Hoshangabad is the younger formation of the Narmada and is represented by the flood plain facies. It forms a distinct morphostratigraphic unit; above the present day flood plain of Narmada. It is crescent-shaped and is confined within the meander looped of the Narmada. This formation comprises three units. The basal unit predominantly consists of rock gravels of quartzite, gneiss, basalt and agate imbricated in the grayish sand and silt. The middle unit is represented by unconsolidated imbricated matrix of sand and silt supported by bimodal sorted gravel of quartzite, basalt, agate and chart. The upper unit contains yellowish and brownish silt and clay with occasional calcium concretion. It is capped by black soil. The measured thickness of the units is 6.5m, 5.00m and 3.5m respectively. The sediments of fluvial domain of Narmada identified between an elevations of 268 to 350 m above m.s.l. and were deposited in channel and flood plain environments during upper Pleistocene time

The sequence of Quaternary events and the history of sedimentation of Narmada indicate that the upper 70m to 90m of the Narmada alluvium was deposited in a single aggradation episode with minor pauses when dissection of the alluvium produced two terraces (NT₃-NT₂). The sediments of this aggradation episode constitute three lithostratigraphic units viz. Boulder conglomerate, Sohagpur and Shahganj formation. The sediments of the alluvial phase are underlain by a boulder bed of glacio-fluvial origin. Thus, the fossiliferous boulder conglomerate, the basal unit of alluvium marks a disconformity between the lower glacial-boulder layer and upper fluvial sediments. The fossiliferous basal boulder conglomerate is being of middle Pleistocene age (Khan 1992)

The sediments of present domain of Narmada is represented by sediments of active flood plain, point bar and sand bar facies of present domain and consist of unconsolidated imbricated, stratified, polygonal sorted rock-gravel supported by very coarse to very fine-sand and is named as Janwasa formation, after the village Janwasa where it is best developed. The measured thickness in the valley is about 5m. The Quaternary deposits of Narmada are known for fossil assemblage (Biswas 1997) these deposits and identified two biostratigraphic units Lower assemblage and Upper assemblage. The lower assemblage comprises *Stegodon namadicus*, *Equus namadicus*, *Elephas namadicus* (broad molar) and *Homo erectus*. The upper assemblage comprises *Elephas namadicus* (narrow molar); *Hippopotamus namadicus* and *Equus onager khur*. Fossils of the lower assemblage are found in the Surajkund Formation. Fossils of the upper assemblage occur in the Baneta, Hirdepur and Bauras Formations.

Fossils are not reported from the stratotype section of the Dhansi Formation as well as from its other two localities (Burhi-Narmada and Shakkar river section).

The Quaternary deposit of the Narmada valley is known since the last century for their rich content of mammalian fossil. Some of the early records are by Splibury (1844), Falconer and Cautly (1847), Theobald (1860) and Pricep (1883). A systematic description of these mammalian fossils was given by Lydekker (1874-80) and Pilgrim (1926, 1939). During the last fifteen years, some additional collection of fossil mammals has been made with precise stratigraphic record (Biswas and Dassarma, 1981, 1984; Tripathi and Basu, 1983; Biswas, 1988). A new dimension has recently been added by the discovery of a skull of *Homo erectus* (Sonakia, 1984).

The present biostratigraphic work is based on the field work carried out in parts of Hoshangabad, Sehore, and Raisen, Narsinghpur and Jabalpur district of M.P. and subsequent detailed taxonomic studies in the laboratory.

A large number of fossil mammals have been collected so far, but they represented only a few genera and species. Systematic description for some of these has been published recently. Mention may be made of skull of *Stegodon namadicus* (Biswas and Dassarma, 1981) skull of *Elephas cf namadicus* (Tripathi and Basu, 1983), skull and mandibles of *Hippotamus namadicus* (Biswas and Dassarma, 1984), isolated teeth of *Equus namadicus*, skull, mandible and isolated teeth of *Equus onager khur* (Biswas, 1988) skull of *Homo erectus* (Sonakia, 1984; Lumely and Sonakia, 1985) a mandible of *Cuon alpinus* (Biswas and Basu, in press), and a skull of *Hystrix* (Biswas et al, 1988).

During the progress of the present multi-disciplinary project, a number of fossil vertebrates have also been collected. These included skulls, mandibles, horncores and other post-cranial bones of *Stegodon namadicus*, *Elephas namadicus*, *Bos namadicus*, *Bubalus palaiendicus*, *Boselaphus namadicus*, *Antelope cervicapra*, *Hippopotamus namadicus*, *Sus namadicus*, *Equus*, *Boselaphus namadicus*, *Antelope cervicapra*, *Hippopotamus namadicus*, *Sus namadicus*, *Equus namadicus*, *Equus onager khur*, *onager khur*, *Cervus cf unicolor* and *C. duvauceli*. Amongst these some.

No remarkable distinction has been made between the fossil of Dhansi / Nasrullaha, Shivepuri Formation Khan (1992), Tiwari (1997). Fossil of Dhansi Nasrullahaganj formation, are, however, very fragmentary. These formations have yielded species like *stegodon namadicus*, *Equus namadicus*, *Hippopotamus namadicus*, *Cuon alpinus tripathii* and *Homo erectus*, *stegodon namadicus* shows affinity with the middle Pleistocene *S. orientalis* of SE Asia (Biswas and Dassarma, 1981). The *Equus namadicus* originated from the lower Pleistocene, giving way to extant *Equus onager Khur* during the late Pleistocene (Biswas, 1988). *Sus namadicus*, *Cuon alpinus tripathi* and *Homo erectus* also show Middle Pleistocene affinity Pilgrim (1929); Biswas and Basu, in press and Sonakia (1984). The present study indicated that Surajkund / Shivepur formation is Middle Pleistocene in age whereas the underlying the Dhansi/--- Nasrullahaganj formation probably lowers Pleistocene to Middle Pleistocene in age.

The faunal contact of Baneta Sahaganj Hirdepur/ Demawar Shahganj / Hoshnagabad formations more or less similar. Khan (1992) Other than the species of *Bos*, *Bubalus*, *Boselaphus*, *Antelope*, *Cervus* and *Elephas*, characteristic forms of these formations include *Hippopotamus palaeindicus*, *Hystrix crassidens* and *Equus onager Khur*. *Hippopotamus palaeindicus* succeeded Middle Pleistocene *H. namadicus* and itself became extinct probably during late Pleistocene or Lower Holocene. There is no evidence for *Hippopotamus* in the archaeology sites or cave painting in India. Prolific development of *Hippopotamus palaeindicus* has been reported from the adjacent Late Pleistocene Son Valley deposit (Dassarma and Biswas, 1997). *Hystrix crassidens* is also an extinct form and is reported from late Pleistocene to lower Holocene deposits of *Bankura* (Dassarma et al 1982) and kurnool caves (Lydekker, 1886). *Equus onager khur* is an extant form. A significant feature of Deonawar Baneta formation is the total absence of *Stegodon* which is more or less prolific in the underlying Shivpur / Surajkund and Formation. All these criteria taken together point to late Pleistocene to Lower Holocene Age for three Formations.

The mammalian genera listed above are more or less specialized and sensitive to climate condition. They are mainly warmth loving. None of the above mentioned fauna is indicative of cold climate. However, gigantism observed in some Narmada Middle Pleistocene forms. May be indicative of the Pleistocene's freezing climate. This may be also due to the fact that Narmada fauna mostly represent intermediate stage between the Siwalika fauna and the extant fauna.

The analysis of the available data reveals the dominance of grazers over browsers. This may be partly due to chance of better preservation of grasses. Grass-feeders roaming on flat terrains by the side of the river, definitely had a better chance of preservation as fossils than the browsers roaming in the jungles or hills slopes.

Some forms like *Bubalus*, *Hippopotamus*, *Stegodon*, *Cervus*, *Sus* etc. suggest a prevailing, tropical, humid climate. At the same time some fauna like *Equus* point towards aridity. Admixture of the fauna like *Equus* point towards aridity. Admixture of the fauna depicting humidity and aridity is due to oscillation of climate from humid to arid in warm temperature.

4.5.00 The Narmada Homo erectus and associated Human culture

The Quaternary deposits of Narmada valley in the area Hominid Fossil locality Hathnora consists of 70 to 90 m of Narmada alluvium was deposited in single aggradational episode with minor pauses when dissection of the alluvium produced two terraces (NT1-NT2) The Sediments of gradational episode constitutes three lithostratigraphic units viz Boulder conglomerate, Sohagpur Formation. A Boulder bed of glacio-fluvial origin underlies the sediments of the alluvium phase. Thus the fossiliferous boulder conglomerates, the basal unit of the alluvium marks disconformities between the lower glacio – fluvial boulder layer and the upper fluvial sediments. The fossiliferous boulder conglomerate is considered to be middle Pleistocene age (Khan & Sonakia 1992). The Boulder Conglomerate (Hathnora Formation) is thick wedge of semi consolidated

fossiliferous sediments and concealed under thick deposits of clay and silt of the older alluvium in valley. It may contain the richest vertebrate fossil assemblage and has yielded only known Hominid fossil from the Indian sub-continent (Sonakia 1984). The Boulder bed which yielded Hominid fossil for the first time reported to be of glacial & fluvial origin Khan & Sonakia (1992). The fossiliferous horizons is concealed wedge, it is exposed only at some places in valley by lateral dissection by superimposed drainage, lateral river cutting, dissection of quaternary blanket, and other emotional activities. The Concealed and hidden disposition of fossiliferous horizon and its mysterious intricacy in tectonic zone and due to failure to understand basin configuration has disappointed the geoscientists to add further to any new find of Human skull, in spite of keen interest and intensive and extensive work.

However early human settlements have left widespread signatures on Indian river valley systems in the form of extensively large Paleolithic industry sites, whereas the hominid fossil evidences are very rare of scanty. The skull cop of *Homo erectus*, the 'Narmada Man' is the only example in the long stretch of Quaternary alluvial deposits occurring at varying topographic levels in the Narmada river valley of Central India. Some of the terraces are pretty rich in Paleolithic and Neolithic cultural materials, bearing testimony to the hunting and gathering abilities of early man. Elsewhere in the world, a century of discoveries precedes the discovery of the Indian fossil in 1982. The pioneering discoveries succeeded, with a couple of few finds from java in Indonesia and from China. Later, when the search involved the continents of Africa and Europe there was a spurt in the rate of discoveries and as a result, a large number of human fossils were discovered, thereby making the study of human evolution more logical and factual. It has to be mentioned here that initially discoveries of human fossils from Germany in Europe and from java in Asia, were very much neglected. The Indian fossil mentioned by Theobald (1881) was lost due to sheer curatorial negligence (Kennedy, 1980), whereas the Garman Neandethal specimen was initially misinterpreted. It took time to describe the Narmada specimen in detail and it also invited controversies regarding its taxonomic. (De Lumley and Sonakia, (1985); Kennedy at al, (1991). Some of the observations made later during detailed studies of the Narmada specimen are listed in the description given below.

The Narmada hominid specimen was discovered way back in 1982, during the course of systematic collection of faunal material in the Quaternary alluvial deposits of the Narmada Valley to erect the mammalian biostratigraphy. The Specimen lay exposed with its left Lateral cross-section leveled along with the eroded, gravelly terrace deposits on the right bank of the Narmada river near the village Hathnora in Shahganj block of Sehore district, about 80 km South South East of Bhopal.

Better preservation of human fossils at the beginning of burial as a ritual, has always been a matter of very rare chance. This leaves much of it to reconstruction of missing parts. Like many other controversies, the Narmada hominid has preserved many vital parts, thus leaving little scope for baseless speculations. Barring parts of facial lets parital and left temporal; all other bones are intact in the skull. The brownish color has developed as a result of intense mineralization. The skull is heavy, massive and to some

extent robust. Contrary to the progressive brachycephalic trend, the Narmada skull is long and low. It has been classed in the dolicocephalic primitive group. The skull is wider near the base, whereas with progressive evolution the human skull grew wider and bulbous in the parietal region. In posterior view, a peculiar angularity, casting a pentagonal form is observed, the apex of this angularity is formed due to sagittal prominence and while descending laterally due to a wide angularity at the level of superior temporal line, from where the skull wall actually falls apart. There is a slight acute angle, when the temporal bone curves down in basal part of the skull. The mid sagittal curve is regular, gentle and smooth. The glabellar region is detached but the adjacent supr-orbital torus is quite massive and prominent, which project interiorly, shielding the eye sockets with almost circular portals a feature very much common to East African predecessors.

A sharp and prominent superior temporal line, developed as a remarkable positive relief, gradually curves down posterior to form a very strong, knot-like, angular torus. The mastoid and supra mastoid crests bulge more prominently than in some of the *Homo erectus* crania. The partially abraded mastoid process is typically small as observed in the primitive crania. Its shorter length is further confirmed by the posterior lengthening of incisures parietalis. Placed in a slightly higher position clutched between two secondary branches of the root of zygomatic process the external auditory meatus shows archaic features in tympana area. There is more or less a deep fissure between the petrosal crest of tympanic plate and the mastoid process. This sort of primitive trait is very much present in almost all the Indonesian and *Sinanthrope* skulls.

In *Homo erectus* occipital, the inion-endinion distance is quite a complex figure. It is as a result of interaction two morphologies, the endocrinal proportions and the nuchal musculature (stringer 1984). The inion is placed at a moderately higher level and coincides with opisthocranium, thereby leaving significant occipital plenum below, bounded by a definitely strong but abraded occipital torus. The sagittal suture is placed in a raised and prominent parietal keel which is flanked continuously on either side by a pair of shallow parasagittal depression.

There are certain basic similarities between the Narmada skull and the better established *Homo erectus* crania viz., the Sagiran 17 *Sinanthropes Pakinensis* (Particularly skull XI) studied by Weidenreich (1943) many of the so called primitive features are restricted to the Asian clan of *Homo erectus*, whereas others are restricted to the African continent and are not reflected in Asian counterparts.

The gravel that yielded the Narmada *Homo erectus* carries heavy hand axes, cleavers and scrapers. While taking cross-traverses between Vindhya and Satpura mountains vast industry sites of the same palaeolithic implements have been located close to the mountain ranges. These sites appear to have been inhabited by Narmada hominid and its contemporaries for generations together. The excavation out by Deccan College, Pune (personal communication from V.N.Mishra) have yielded the lower Palaeolithic assemblage, in an excavation near Samnapur 10 km North of Narsimhpur. A similar excavation was planned 4 km north of the Hathnora hominid site near villages Nimton

and Rampura where the cultivated land is practically strewn with Paleolithic implements. The excavation was carried out in three pits measuring 1 m x 1m x .60 m. In all the pits each dig was approximately 10 cm. Thus nearly 6 digs were made. Debit age included a very large proportion of the material excavated. The size of debit age gradually decreased towards bottom. The debit age material comprises mainly Vindhyan quartzites which are found locally. Thus excavation, if carried out in the Narmada Valley, would provide an excellent correlation of the Paleolithic and Neolithic cultures which prevailed in Central India

The Quaternary deposited of Narmada Valley consist deposit of sediments of two domain viz. deposits of interglacial domain (Boulder conglomerate-Hathnora formation) and fluvial deposit of paleo-domain of Narmada. The boulder conglomerate is fossiliferous horizon of Narmada and has yielded skull cap of *Homo erectus* Narmadensis at (1984). It is marker horizon and represents interglacial phase in the history of Quaternary sedimentation in Narmada Valley around Hathnora upstream of Hathnora, Gurwara, and Sardapur. The occurrence of ash bed reported from Quaternary sediments of Narmada (1991). The Quaternary deposit of Narmada is associated with two horizons of ash beds of middle and upper Pleistocene age. These ash beds are designated as NAB-I and NAB-II in ascending antiquity in the valley. The Ash bed NAB-I is associated with upper gritty units of boulder conglomerate (Hathnora formation) and is identified at an elevation of about 290 m above m.s.l. The Ash bed NAB-II is associated with upper units of clay silt deposits of paleo-domain of Narmada (Shahganj formation) and is identified at an elevation of 310 m above m.s.l. The study of grain morphology of glass matrix, their relation with other minerals shape, size, texture of lithic fragments and association of other ashy sediments of pyroclastic origin suggest that sediments were brought from distant source by Aeolian agencies, during the different phases of sedimentation in Narmada Valley. These ash beds are used as tool in correlation of different quaternary deposits in the valley.

The Quaternary geological history of Narmada is characterized by eruption of voluminous Deccan basaltic lava flows in the southern margin represented by Satpura and Vindhyan sedimentation represented by cuesta on the northern margin which have sharp cut across the deep valley floor of pre Cambrian basement. The rift linear trench is intruded by the dolerite and other mafic and siliceous dykes and sills along lineaments in different phases of tectonic deformation. The mafic and siliceous volcanic activity is both isolated and simultaneous and their disposition is mostly asymmetric across the rift zone. The Quaternary sedimentation incepting from glacial activity, followed by fluvio-glacial, lacustrine and fluvial phase within the rising and sinking environment, block faulting and segmental and linear displacement and dislocation, uplifting and isolated domal up-lift, Neogene rifting and Quaternary sedimentation and rift-bound Pliocene-Pleistocene rifting and volcanic activity specifically during glacial and fluvio-glacial phase are major component of the Quaternary period and tectonic processes of the Narmada Rift System. The Quaternary sedimentation was triggered by tectonic activities / up lift and climatic changes. The provenance for these sediments is the weathering products of eroding pre-Cambrian, meta-sediments, sedimentary and volcanic rocks along the watershed upland, rift

escarpments and shoulders; faulted and uplifted blocks, volcanic fissure zones, and plateaus within and outside the rift. The Narmada Rift System, bounded by adjacent plateaus rising 300–700 m above the rift floor, consists of number symmetrical and symmetrical faulted blocks, escarpment, rock cut terraces, rock floors and segments of micro half grabben. Although rift-related basins started to form during the late Oligocene to early Miocene times, the Narmada Rifts were fully defined by middle to late Miocene time.

The Narmada rift system basins provided a unique setting for dynamic ecosystems that were characterized by Rift-related subsidence and coeval sedimentation also created an ideal loci of Quaternary sedimentation and environment for the accumulation of sediments volcanic fabrics sediments, burial, diagenesis, and preservation of organic remains. Because rifts formed after widespread Quaternary sedimentation occurred and voluminous sediments in the rift basins were accumulated by glacial activity consequential upon the lowering of temperature and climatic changes in the region.

The Miocene -Pliocene–Pleistocene lake deposit of Katni on the eastern rift shoulder was created by faulting, topographic control, or isostatic depression similar to that of other Rift system.

Abundant faunal remains in rift settings, including hominids Sonakia (1984), suggest that the Narmada Rift System created productive ecosystems during Pliocene–Pleistocene time. The volcanic rocks within the fossiliferous sediments provide temporal information for calibrating and sequencing hominid and other faunal evolution. The detailed geological sedimentological geochemical of interbedded tephra ,Quartz grain morphology of sediments of quaternary strata and palo-sole of and geochronological studies of from the different localities for establishing accurate biostratigraphic and lithostratigraphic data, sedimentation rates, and paleoenvironmental and tectonic histories of different sdimt coloumns in area along the rift system. Interbedded volcanic rocks allow determination of the time of rifting, the beginning of sedimentation, sedimentation rates, and the oscillation from glacial, fluvio-glacial lacustrine to fluvial environments. The cyclic environmental transitions recorded in the sedimentary sequences of the rift basins are caused by tectonic activities (uplift and subsidence), changes in relief, and climatic variations. The climatic changes in uplift ,topographic and landscape features, coupled with block faulting , rinsing and sinking platform , created basins for the accumulations of thick lacustrine and fluvial sediments sequences with terrestrial and aquatic fossils. The sequential change in the sediment fancies from finely bedded lacustrine deposits to fluvial sediments are commonly noted in the sedimentary sequences and reflect environmental and tectonic changes that can be temporally determined. Moreover, regional correlation based on the chemistry and geochronology of interbedded tephra has made it possible to establish accurate stratigraphic relations that are useful for paleoenvironment reconstruction and evolutionary studies of fossil remains in the Narmada rift valley Khan et.al. (2012). Regional tephra correlation is being used increasingly to link sites together, and has

732 already established that similar tephra layers are known from other parts of rift
733 valley, as well as from other basin and peninsular India Achariya, (2002), Biswas,
734 Khan (1992) Khan et.al. (2012) Tiwari (1996). There is a great potential for
735 further correlation of tephra in the Rift System and marine sediments in the
736 Arabian Sea. The Arabian Sea has a continuous record of deposition that extends to
737 at least 7 million years. The Quaternary sediments interbedded with tephra with
738 within the age range of the ODP Ocean Drilling Program 721/722 stratigraphic
739 sections of the Arabian Sea are also present within the rift floor and the western rift
740 margin of the region. The chemical and chronological correlations of ash beds
741 within the rift sequences of have been made with ashes described in marine
742 sections. Detailed correlations based on orbitally calibrated time scales of pale
743 magnetic stratigraphy Rao (1996) within Quaternary sediments of rift deposits will
744 provide ties to establish global climate changes based on the terrestrial and marine
745 sediments of the rift system.

746
747 Moreover, because of tephra layers in sedimentary basins of different geologic
748 periods, processes such as faulting, rifting, sedimentation and diagenesis, impact of
749 climatic changes, age of fossils, nature and acquisition of archeological implements,
750 and the origin, distribution, and functional significance of early hominid artifact
751 assemblages can be deciphered. However, evidence of the effects of tectonics on
752 fauna and flora are distinct and its signatures on dislocation and concealing of
753 fossiliferous horizons are uncontrolled and ill defined in the ecosystem in the valley
754 during the Pliocene–Pleistocene periods. The boulder conglomerate which yielded
755 the skull cap of *Homo erectus* in Narmada rift from Hathnora Sonakia (1984)
756 remained only discovery of hominid fossil in last two and half decade due
757 inconsistency concealed nature of fossiliferous horizon due faulting, dislocation and
758 subsidence of Quaternary blanket of Narmada rift system as such researcher and
759 scientist failed to add any further knowledge to hominid discovery any further .

760
761 The Narmada Rift System consists of symmetrical basins that have been evolved in
762 different stages of tectonism. The 100 -120 km-wide and 1300 km long rift
763 bounded by Satpura in south and Vindhyan in north constitutes conspicuous ENE-
764 WSW to E-W rift basin zone is filled with Pliocene–Pleistocene sediments, whereas
765 some of them contain Miocene sedimentary deposits. Most of the sedimentary
766 sequences contain faunal and floral remains including hominid species. Most of the
767 basin-fill sediments were derived from topographically elevated rocks that are
768 present within and outside the rift basins. Lava flows and tephra are interbedded
769 with the fossiliferous sediments clastic sediments derived from the provinces of
770 sediment mostly from crystalline basement volcanic, sedimentary, meta basic and
771 sedimentary rocks, aided in the cementation and preservation of organic remains by
772 providing secondary minerals released during alteration in a burial environment.
773 Quick burial minimized the effect of preburial taphonomic processes. Moreover,
774 chemical constituents released by the alter have provided critical temporal and
775 spatial information without which the study of hominid evolution and pale
776 environmental reconstruction in the Rift System would have been impossible.

777

Moreover, because of tephra layers in sedimentary basins of different geologic periods, processes such as faulting, rifting, sedimentation and diagenesis, impact of climatic changes, age of fossils, nature and acquisition of archeological implements, and the origin, distribution, and functional significance of early hominid artifact assemblages can be deciphered. However, evidence of the effects of tectonics on fauna and flora are distinct and its signatures of dislocation and concealing of fossiliferous horizons are uncontrolled and ill defined and the ecosystem in the rift system during the Pliocene–Pleistocene periods is not clear. The boulder conglomerate which yielded the skull cap of *Homo erectus* in Narmada rift from Hathnora remained only discovery of hominid fossil in last two and half decade due to inconsistency of concealed nature of fossiliferous horizon in Narmada valley. Historical or modern analogs illustrate the potential of the regional and sometimes global effects of such major silica eruptions in the geologic past of sedimentation, sedimentation rates, and the oscillation from lacustrine to fluvial environments. The cyclic environmental transitions recorded in the sedimentary sequences of the rift basins are caused by tectonic activities (uplift and subsidence), changes in relief, and climatic variations. Changes in topographic features, coupled with volcanic damming, created basins for the accumulations of thick glacial, fluvio-glacial lacustrine and fluvial sequences with terrestrial and aquatic fossils. Changes from finely bedded lacustrine deposits to fluvial sediments are commonly noted in the sedimentary sequences and reflect environmental and tectonic changes that can be temporally determined. Moreover, regional correlation based on the chemistry and geochronology of interbedded tephra has made it possible to establish accurate stratigraphic relations that are useful for pale environment reconstruction and evolutionary studies of fossil remains in the rift valleys across India sub continent. Regional tephra correlation is being used increasingly to link sites together, and has already established that similar tephra layers from known area in Rift system.

The critical study of the area around Hathnora (22°52'77°52') between Vindhya and Satpura mountains vast industry sites of the same palaeo lithic implements have been located along the edge and pediment slope of the Vindhyan and basaltic upland. These sites appear to have been inhabited by Narmada hominid and its contemporaries for generations together as they left over stone industry sites and unfinished material haphazardly indicate sudden dissemination and migration of hominines in search of shelter and safe places after volcanic eruption. The reporting of remains of *H. sapiens* in the area of Bhembetaka about 20 Kms north of hominid locality of Hathnora and relicts of rocker shelter and ancient human signature documents the dispersal of hominines after the eruption of volcanic matrix and its fall at Wakankar, V.S., 2002. The association of skull cap Sonakia (1984) and Sankhyan (2007) described two new hominine fossils from Netankheri up stream of Hathnora in the Central Narmada valley. They include a partial left humeral diaphysis and a distal shaft fragment of the left femur associated with Hathnora formation. Khan & Sonakia 1992 and analysis of quartz grain morphology of sediment columns of Hathnora and quartz grain morphology of paleosol across the quaternary strata of bore hole sampling across 556 m of rock basin revealed that *Homo erectus* of Narmada partly sustained in the glacial and

fluvio-glacial environment which is documented and witnessed by Hathnora formation which marks the end hostile climate and environment of sedimentation in Narmada valley Khan (2013). The study of quartz grain and their micro structures of paleo soil identified in concealed blanket of quaternary deposits display relatively heterogeneity in sediment characteristics throughout across the Quaternary column of Hathnora section in Central Narmada valley. The quartz grain of paleo soil and sediments display digonestic characteristics of glacial fluvio-glacial and fluvial under concealed environment of sedimentation at different depth. The significant breaks in grain morphology of quartz grain surface texture and associated elements and granular matrix in the sequence of quaternary strata is recorded at to, 150, 350 and 350 and beyond which linked to long evolution of glacial fluvio glacial and fluvial environment of sedimentation in time and space in increasing antiquity in the valley from the base Khan (2013). The occurrence of Bolder conglomerate and occurrence of haphazard and disserted sites of Palaeolithic sites indicates that area suffered dual hostility first the end of glacial activity with the stratigraphic column at the level of 83 m where the Homo erectus survived Sonakia (1984) both by glacial activity and volcanic eruption, whereas in the second hostility evolving human activity by volcanic eruption and dense smoke and dust and darkness. And consequent upon H Sapiens and associated were forced to migrated to safer places. In view of the recorded observation the concealed quaternary strata to the level of about 350 m below ground level further needs attention of geoscientist for the search of human remains in Narmada valley.

The study of assemblage of glass matrix of Ash bed, grain morphology of glass their relation with other minerals shape, size, texture of litho fragments of pyroclastic origin suggest that sediments were brought from distant source by Aeolian agencies in the form of thick cloud containing volcanic dust, rock matrix and different gases which remained in atmosphere for very long time and settled down across the Indian sub continent during the different phases of river sedimentation. Further study of Ash bed material and silica revealed diagnostic morphological characters of glass shards which are typical of silica volcanism (Heiken, 1972, 1974) and show close similarity with those reported from the Quaternary tephra beds of the Narmada, Son, Purna and Kukdi basins (Basu et. al., 1987; Khan et.al. 1991 Basu and Biswas, 1991; Singaraju and Shivaji, (1991) Mukhopadhyay, (1992). It is significant to note that the occurrences and association of two marked horizons at different levels further reveal that the cyclic eruption and settling of volcanic matrix has taken place with pause in the valley.

In Narmada valley the association of Ash bed NAB-I with Hathnora formation at the depth of 78 m in Quaternary column and occurrences of skull cap of Homo erectus at the depth of 83 m in decreasing antiquity from the top assumed that Toba eruption have taken place later than existence of Homo erectus which appeared and resided in the valley for long time before the fall of Toba ash. The association of Ash is NAB-II at the depth of 72 m with the younger deposit revealed the second cyclic fall of Toba ash which certainly have had influenced on hominines and had collective and cumulative impact on Homo erectus (Sonakia1984) Homo sapiens (Thobold 1860, 81), in Narmada valley and Indian sub-continent.

Using phytogeographic data, Oppenheimer (2003) argues that Homo. Sapiens occupied India before ~74 ka and may have undergone “mass extinction” as a result of the Toba eruption. The argument of Oppenheimer (2003) is in strong conformity with the present observation of authors. As sediment & Ash bed sequence of Quaternary column of Narmada (325 m) and occurrences of fossil of skull cap of Homo erectus (Sonakia 1984) at 83 m & human cranium Homo sapiens (Thebold 1960, 1981) transported have existed prior to fall of Toba ash and they are among the few who in spite of mass extinction caused by mega dislocation in ecology and environment related with volcanic eruption survived in Narmada Valley. It is further documented by the rarest occurrences of these fossils in subcontinent which also confirm the intensive impact of volcanic ash fall on these hominines and their consequential mass extinction.

The Toba eruption of 74 ka was distinctly and clearly a mega event of very great magnitude and intensity, far greater than any known historical eruption, suggesting it had very devastating impact and repercussions. It has changed the global climate environment and ecology.

The Quaternary deposit of Narmada Valley consists of sediments of three domains viz. deposits glacial, fluvio-glacial (interglacial) and fluvial. The boulder conglomerate is fossiliferous horizon of Narmada and has yielded skull cap of Homo erectus Narmadenesis Sonakia (1984). The Ash bed is found associated with Hathnora formation Khan et al. (1991), in the area around Hathnora, upstream of Hathnora, Gurwara, Sardapur two horizons of Ash bed are identified in Boulder conglomerate (Hathnora formation). These are designated as NAB-I and NAB-II in ascending antiquity in the valley. The Ash bed NAB-I is associated with lower litho units of boulder conglomerate which is quite persistent and well preserved. The NAB-I (L1 to L3) contains three micro layers and NAB-II (L4 to L5) two micro layers in increasing antiquity. The lower unit of boulder conglomerate is associated with NAB-I, whereas NAB-II is associated with younger deposits. The study of grain morphology of glass matrix, their relation with other minerals shape, size, and texture of lithic fragments and association of other ashy sediments of pyroclastic origin suggest that sediments were brought from distant source by Aeolian agencies which fell and settled down during the different phases of sedimentation in valley. The Ash bed NAB-I is embedded within the reddish brown sand silt and in the upper pebbly gritty unit of boulder conglomerate (Hathnora formation).

The Ash bed NAB-II is associated with yellow, reddish, brown, clay and silt of fluvial deposits Khan et al. (1992). It is pinkish grey, light brown in colour, fine grained, porous and non-plastic in nature. The average thickness is about 25 cms

The study of grain morphology of glass matrix, their inter-relation with other minerals shape, size, texture of lithic fragments and association of other ash sediments of pyroclastic origin, indicate that they are the product of highly explosive silica volcanism. Pumice shards tend to develop from relatively high viscosity rhyolitic

916 magmas with temperature less than 850 C whereas cusped shards are most likely to
917 develop from low viscosity rhyolitic magma with temperature more than 850 C suggest
918 that sediments of pyroclastic origin and were brought from distant source by Aeolian
919 agencies, after extrusive volcanic activity in middle and upper Pleistocene time.

920
921 There are quarries and questions that necessitate consideration of two separate aspects of
922 the eruption; first, the consequences of possible rapid global climatic deterioration and
923 second, the direct effects of the ash-fall on hominines and their environments in Narmada
924 valley and in Indian subcontinent.

925
926 The association of Ash bed NAB-I with Hathnora formation at the depth of 78 m in
927 Quaternary column of Narmada and occurrences skull cap of *Homo erectus* of Narmada
928 at the depth of 83 m assumed that Toba eruption might have taken place later than that of
929 existence of *Homo erectus* which appeared and resided in the valley for long time before
930 the fall of Toba ash. The association of Ash bed NAB-II at the depth of 72 m with the
931 younger deposits indicates the second cyclic of fall of Toba ash which might have
932 influenced the younger Hominines i.e *Homo erectus* & *Homo sapiens*, in Narmada
933 valley.

934
935
936 The occurrences of these skull caps with short range of their occurrences in the
937 stratigraphic column of Narmada with the Ash beds horizon NAB-I and NAB-II and
938 specially with the Hathnora formation one at the top at an average elevation of about 268-
939 273 m above the mean sea level and other with younger deposits had revealed the close
940 association with volcanic activity with their existence. The Toba Ash fall is also in very
941 close range with the sequence of sedimentation and occurrences with both the skull caps,
942 which certainly has its impact on the middle and late Pleistocene Hominines in
943 Narmada valley and Indian subcontinent.

944
945 The oldest fossil from India is represented by the Narmada hominine dated to not less
946 than 236 ka (Cameron et al., 2004), or to some time in between 150 and 250 ka
947 (Kennedy, 2001:167). Modern human remains have been discovered in an undated Late
948 Paleolithic context at Bhimbetka rock shelter III-A-28 (Wakankar, 2002:5) which is
949 situated about 20 km north of Hominid locality Hathnora and from three cave sites in Sri
950 Lanka, dating from 27.7 ka (Kennedy 1999, 2001). Using phytogeographic data,
951 Oppenheimer (2003) argues that *H. sapiens* occupied India before ~74 ka and may have
952 undergone "mass extinction" as a result of the Toba eruption. The later argument is in
953 conformity with the observation of authors as it is well illustrated by close association of
954 Ash bed and *Homo erectus* of in sediment sequence of Quaternary column of Narmada.

955
956 The study of grain morphology of glass matrix, their relation with other minerals shape,
957 size, and texture of fragments and sediments of pyroclastic origin suggest that sediments
958 were brought from distant source in the form of thick cloud containing dust matrix and
959 volcanic ash which was highly explosive and siliceous in nature and remained in
960 atmosphere for quite long time. The height of the eruption column appears to be
961 considerable. It is postulated that the tephra preserved as disconnected bodies within the

river valley sediments represent rapidly settled ash falls from a volcanic ash cloud which formed a canopy over a large part of river basins for longer time of Peninsular India where sedimentation was on in different river basins including Narmada valley. The discontinuity of Ash bed in Narmada valley and Indian subcontinent is attributed to be associated with column of volcanic eruption, quantum of volcanic matrix, wind direction, moisture density of air and rate of fall of matrix on oscillating platforms of sedimentation in different basin. It is significant to note that the occurrences and association of two marked horizons at different levels further reveal that the cyclic eruption and settling of volcanic matrix was with pause in the valley which perhaps related with pause in volcanic eruption

The volcanic eruption and consequential ash fall has created severe dislocation in ecology and environment and adversely affected hominines in Narmada valley and Indian subcontinent. It is witnessed by association of Ash bed NAB-I with Hathnora formation at the depth of 78 m in Quaternary column and occurrences skull cap of Homo erectus at the depth of 83 m in decreasing antiquity from the top assumed that Toba eruption have taken place later than existence of Homo erectus which appeared and resided in the valley for long time before the fall of Toba ash. The association of Ash is NAB-II at the depth of 72 m with the younger deposit revealed the second cyclic fall of Toba ash which have had influenced collective and cumulative the Homo erectus (Sonakia1984) Homo sapiens (Thobold 1860, 81), in Narmada valley and Indian sub-continent.

The study of cyclic Toba ash fall and using phytogeographic data, Oppenheimer (2003) argues that Homo.Sapiens occupied India before ~74 ka and may have undergone “mass extinction” as a result of the Toba eruption. The argument of Oppenheimer (2003) is in strong conformity with the present observation of authors. As sediment & Ash bed sequence of Quaternary column of Narmada (325m) and occurrences of fossil of skull cap of Homo erectus (Sonakia1984) at 83 m & human cranium Homo sapiens (Thebold 1960,1981) (transported) have rarest occurrences of human fossils in Narmada valley and subcontinent which also confirm the intensive impact of volcanic ash fall on these hominines and their consequential mass extinction caused by mega dislocation in ecology and environment by volcanic eruption.

Toba Eruption and Its Impact on Environment

In Narmada valley and Indian sub continent the Environment and climate during late pleistocene has been significantly affected after Toba eruption, according to Rampino et al. (1988) the size of Toba at 74 ka could have induced a volcanic winter, similar to predicted nuclear winter scenarios, as modeled by Turco et al. (1983, 1990). The injection of vast amounts of gaseous aerosols and volcanic dust and matrix into the atmosphere, which follow large volcanic eruptions, is predicted to have detrimental and decisive consequences for changes of global climate. Past historical eruptions, such as Tambora in 1815 (Stothers, 1984) and Pinatubo in 1991 (Mc Cormick et al., 1995), have provided evidence of post-eruption climatic deterioration and climatic. With the eruption


of Toba having been far larger than both of these historical eruptions, its consequences are therefore assumed to have been far more devastating. The imprints of divesting impacts in the area of study and specific on human population are matter of scientific concern and yet to be investigated. The studies conducted of Ash bed associated with Quaternary deposits of Narmada Valley consist of sediments of two domain viz. deposits of interglacial domain (Boulder conglomerate-Hathnora formation) and fluvial deposit of paleo-domain of Narmada. The boulder conglomerate is fossiliferous horizon of Narmada and has yielded skull cap of *Homo erectus* Sonakia (1984). (Khan & Sonakia 1991). It is marker horizon and represents interglacial phase in the history of Quaternary sedimentation in Narmada Valley. The occurrence of ash bed reported from Quaternary sediments of Narmada are associated with two horizons of ash beds of middle and upper Pleistocene age (Khan et al. 1991). These ash beds are designated as NAB-I and NAB-II consisting of five layers designated as (L1 to L5) in ascending antiquity in the valley. The Ash bed NAB-I (L1- to L-3) is associated with upper gritty units of boulder conglomerate (Hathnora formation) and is identified at an elevation of about 290m. Above m.s.l. The Ash bed NAB-II (L-4 to L5) is associated with upper units of clay silt deposits of paleo-domain of Narmada (Shahganj formation) and is identified at an elevation of 310m. above msl. The study of grain morphology of glass matrix, their relation with other minerals shape, size, texture of lithic fragments and association of other ashy sediments of pyroclastic origin suggest that sediments were brought from distant source by Aeolian agencies, during the different phases of sedimentation in Narmada Valley. It is observed that in stratigraphic column two horizons of Ash bed occurred at the vertical distance of about 20 m. which indicate that there were two phases of settling volcanic ash in Narmada valley and Indian subcontinent as a whole with significant time break. The element of time break is in terms of sedimentation perceptible appears to be related with global climatic changes induced by the super volcanic eruption. These ash beds are used as tool in correlation of different quaternary deposits and archeological sites in the valley. The occurrences of two horizons of Ash bed and their deposition by settling of volcanic dust with time gap suggest induced and defused atmospheric conditions in Narmada valley for a very long time after volcanic eruption.

The atmosphere contaminated by volcanic ash volcanic matrix and huge amount of gases and dust particles had definitely affected the atmosphere and climate had have led to temporarily darkened skies in the Narmada valley and Indian subcontinent for very long time. Ash, however, it has had short stay in the atmosphere depending of particles load, wind direction and velocity, moisture density, and volcanic column of eruption and its shrinkage, land barriers, temperature association of gases and longevity of sulphuric acid aerosols. The association of sulphuric acid predicted to have had the most damaging repercussions for Indian subcontinents and global climate. Dense concentrations of H₂SO₄ aerosols in the stratosphere had been the main causative factor to prevent radiation reaching the Earth from the Sun. (Rampino et al., 1988). This process results in cooling of both the atmosphere and the surface of the Earth.

Toba in Indian Subcontinent with special reference to Narmada Valley

The occurrence of ash bed reported from Quaternary sediments of Narmada are associated with two horizons of ash beds of middle and upper Pleistocene age (Khan et al. 1991). These ash beds are designated as NAB-I and NAB-II consisting of five layers designated as (L1 to L5) in ascending antiquity in the valley. The Ash bed NAB-I (L1- to L-3) is associated with upper gritty units of boulder conglomerate (Hethnora formation) and is identified at an elevation of about 290m. Above m.s.l. The Ash bed NAB-II (L-4 to L5) is associated with upper units of clay silt deposits of palco-domain of Narmada (Shahganjg formation) and is identified at an elevation of 310 m. above m.s.l. The study of grain morphology of glass matrix, their relation with other minerals shape, size, texture of lithic fragments and association of other ashy sediments of pyroclastic origin suggest that sediments were brought from distant source by Aeolian agencies, during the different phases of sedimentation in Narmada Valley. It is observed that in stratigraphic column two horizons of Ash bed occurred at the vertical distance of about 20 m. which indicate that there were two phases of settling volcanic ash in Narmada valley and Indian subcontinent as a whole with significant time break. The element of time break is in terms of sedimentation perceptible appears to be related with global climatic changes induced by the super volcanic eruption. These ash beds are used as tool in correlation of different quaternary deposits and archeological sites in the valley. The occurrences of two horizons of Ash bed and their deposition by settling of volcanic dust with time gap suggest induced and defused atmospheric conditions for a very long time after volcanic eruption

Having reviewed the theories surrounding the effects of the Toba eruption on both global climate and human populations in general, it is now possible to turn our attention to evidence provided in the specific context of the Indian subcontinent. By way of background, it is important to recognize that ca. 800 km³ of fine-grained volcanic ash was injected into the stratosphere during the 74 ka Toba eruption. The ash was carried by prevailing winds in all directions, apart from to the south-east. As a result, large areas of southern and southeastern Asia were covered in tephra, from the Arabian Sea in the west to the South China Sea in the East (Figure 1). YTT has been documented in marine cores from the northeast Indian Ocean and Bay of Bengal (Ninkovich et al., 1978a, 1978b; Dehn et al., 1991) and more recently from the Central Indian Ocean Basin (Pattan et al., 1999, 2002), Arabian Sea (Schulz et al.



1090 1998, 2002; Pattan et al., 2001) and South China Sea (Bühring et al., 2000; Song et al.,
1091 2000). The thickness of the YTT bed in these deep-sea cores ranges from ~2 to ~40 cm
1092 and does not appear to reduce significantly with increasing distance from the Toba
1093 caldera. However, marine ash layers, as with terrestrial tephra beds, can experience
1094 reworking and bioturbation and thus some may not preserve a precise record of air-fall
1095 thickness.

1096
1097 Terrestrial YTT deposits have been identified in India (Rose and Chesner, 1987; Khan
1098 et.al.1991) Acharyya and Basu, 1993; Shane et al., 1995; Westgate et al., 1998), Malaysia
1099 (Ninkovich et al., 1978a; Rose and Chesner, 1987; Shane et al., 1995) and possibly in
1100 Bangladesh (Acharyya and Basu, 1993; Acharyya et al.,2000). Of these, Indian YTT
1101 occurrences are far more numerous and extensive, having been documented in river
1102 valleys throughout the subcontinent (Table 1). Well-described Toba ash localities include
1103 those from the Middle Son and Central Narmada valleys in Madhya Pradesh, the Kukdi
1104 river in Maharashtra, and the Indravati, Nagavali, Vansadhara,

1105
1106

1107 The documentation of occurrences of YTT deposits in India has been relatively recent.
 1108 Williams and Royce (1982) reported the first discovery of volcanic ash in India,
 1109 encountered during explorations in the Middle Son valley in 1980. This was followed by
 1110 further discoveries of tephra deposits in the Middle Son valley, as well as in the central
 1111 Narmada valley (Basu et al., 1987, Khan et.al 1991), the Kukdi river valley (Korissettar et
 1112 al., 1988), and in the Vansadhara, Nagavali, Mahanadi, Brahmani and Indravati river
 1113 basins of Orissa (DevdasandMeshram,1991).Around the same time a tephra layer was
 1114 recorded
 1115

Table 1. A summary of Indian tephra localities, indicating the nature of the tephra deposits, those that have been identified as YTT and dated by absolute means, and those that have been found in association with Paleolithic artifacts

<i>River Valley</i>	<i>Tephra Locality</i>	<i>Tephra Thickness (m)</i>	<i>Extension of Tephra (km)</i>	<i>Geochemically Characterized as YTT?</i>	<i>Absolute Dates</i>	<i>Archaeology-tephra Associations?</i>
Middle Son	Ghogara	1.5	30	Yes ^{a,b,c,d}	No	Yes (Middle and Upper Paleolithic)
	Ramnagar	0.5				
	Nakjhar	0.7				
	Khutiali	1.5				
Central Narmada	Pawlaghat	0.24–0.4	20	Yes ^{b,c,d}	121 ± 22 ka ^d	Yes (Middle and Upper Paleolithic)
	Guruwara	0.8				
	Devakachar	<1				
	Hirapur	<1				
Kukdi	10 sections near Bori	0.2–1	3.5	Yes ^{b,c,d}	1.4 Ma ^a ; 0.54 Ma, 0.64 Ma, 0.023 Ma ^f ; 0.67 Ma ^e .	Yes (Lower Paleolithic)
Vansadhara	Goguparu	1.25	2	Yes ^b	No	Yes (Lower Paleolithic, but stratigraphic association between ash and artifacts is unclear)
	Kareni	3–5				
Nagavali	Rayagada	1.5–6	17	Not analysed	No	Not reported
Mahanadi	Kumbia	2.1	45	Yes ^d	No	Not reported
	Krushnamohanpur	2				
	Pitamchul	3				
	Sonepur	1.5–2.5				
Brahmani	Samal	2.5	2.5	Not analysed	No	Not reported
Indravati	Damkelar	0.2–0.3	0.5	Not analysed	No	Not reported
Sagileru	(not specified)	0.3–2	90	Not analysed	No	Not reported
Barakar	Opposite Barakar town	1.83	0.1	Yes ^b	No	Not reported
Purna	e.g. "Gandhigram Place"	Not specified	Not specified	Yes ^{c,d}	84 ± 16 ka ^d	Yes Paleolithic (Acheulean?) ^f
Karha	Morgaon	0.17	0.54	Not analysed	No	Yes Lower Paleolithic (Acheulean)
(Andhra Pradesh?)	"Karnool area" ^d	Not specified	Not specified	Yes	No	Not reported

aRose and Chesner (1987);

Khan et.al (1991)* In Narsighpur- Sardapur Sardarpur-Hathnora & Guruwara and Pawlaghat section

bAcharyya and Basu (1993);

cShane et al. (1995);
dWestgate et al. (1998);
eKorisettar et al. (1989);
fHorn et al. (1993);
gMishra et al. (1995)

from the Barakar river basin in West Bengal- Bihar (Basu and Biswas, 1990), and along the Sagileru river valley in Andhra Pradesh where an extensive tephra bed is traceable for 90 km (Anonymous, 1991; Acharyya and Basu, 1993).

None of these tephra deposits were identified as YTT until 1987 when Rose and Chesner compared the Middle Son tephra with YTT samples from Lake Toba and marine cores. They concluded that the Son tephra was evidence of an extensive co-ignimbrite ashfall (Rose and Chesner, 1987). Further, the true extent of YTT distribution in India was not realized until the work of Khan et al. (1991) who has identified well defined two level of Ash bed for the first time in vertical column of Quaternary deposits at different level and well defined different micro layers in the womb of Narmada. These ash beds are designated as NAB-I and NAB-II consisting of five layers designated as (L1 to L5) in ascending antiquity in the valley. The Ash bed NAB-I consist of three layers (L1- to L-3) is associated with upper gritty units of boulder conglomerate (Hethnora formation) and is identified at an elevation of about 290 m above m.s.l. The Ash bed NAB-II (L-4 to L5) is associated with upper units of clay silt deposits of palco-domain of Narmada (Shahganj formation) and is identified at an elevation of 310 m. above m.s.l. These layers have distinct and diagnostic chemical composition and associated quartz grains have distinct imprints of morphological elements. (Table No-----). Acharyya and Basu (1993). They geochemically characterized tephras from five Indian river valleys (Kukdi, Narmada, Son, Vansadhara, Barakar), identifying all of them as YTT. Their work was later confirmed and built upon by two separate studies. First, by Shane et al. (1995) who analyzed tephras from the Kukdi, Narmada, Purna and Son valleys, Second, by Westgate et al. (1998) who confirmed the presence of YTT in the Vansadhara, Mahanadi, Middle Son, Central Narmada, Purna, and Kukdi river valleys, as well as in the Karnool area. They also obtained fission-track dates for ash samples from the Purna and Narmada basins, giving dates of 84 ± 16 ka and 121 ± 22 ka respectively (Westgate et al., 1998). It appears that samples of volcanic ash from the Indravati, Brahmani, Nagavali and Sagileru river valleys have not been geochemically characterized, although it is assumed by Acharyya and Basu (1993) that these are all deposits of YTT.

Impacts of volcanic Ash-Fall on Late Pleistocene Hominines in the Indian Subcontinent

The Toba eruption of 74 ka was clearly an mega event of great magnitude, far greater than any known historical eruption (Table 2; Figure 3), suggesting it had devastating impact and repercussions as it has change global climate and environment. There are Questions and quarries from many quarters, however, as to the scale of these repercussions is concerned these were of size able magnitude and has influenced late

pleistocene Hominess in Narmada valley and Indian subcontinent. The effects of Toba may or may not have been far-reaching such that the eruption can be said to have had an impact on a global scale. Perhaps impacts were on a more regional scale, for example, at the level of Southeast Asia or the Indian subcontinent as a whole. Alternatively, the hazardous effects of Toba may have been localized, whereby individual habitats or ecosystems were affected, yet other areas in a region remained unscathed.

These questions necessitate consideration of two separate aspects of the eruption; first, the consequences of possible rapid global climatic deterioration and second, the direct effects of the ash-fall on hominines and their environments in India. An additional issue that needs to be addressed is the identification of those hominines that inhabited India ~74 ka and thus those who may have been affected by Toba. In this regard, numerous questions need to be asked. How many hominines species occupied India at the time of the eruption, one species or more? Was *H. sapiens* present in the subcontinent? ~74 ka, or did initial colonization occur at a later date?

Unfortunately, hominine fossil evidence from the Indian subcontinent is limited and cannot currently provide answers to these questions. Only a few specimens are known, none of which belong to the time-frame of the Toba

S. C. JONES


Table 2. Eruptive volumes (DRE) of a number of well-known volcanic eruptions

Volcanic Eruption	Date of Eruption	Location	Dense Rock Equivalent (DRE) in km ³	Reference
Fish Canyon Tuff	27.8 Ma	La Garita caldera, Colorado (USA)	4500	Mason et al., 2004
YTT	74 ka	Toba caldera, Sumatra	2800	Rose and Chesner, 1987
Huckleberry Ridge Tuff	2 Ma	Yellowstone, Wyoming (USA)	2200	Mason et al., 2004
OTT ^a	788 ka	Toba caldera, Sumatra	800–1000	Lee et al., 2004
Oruanui	26.5 ka	Taupo, New Zealand	530	Mason et al., 2004
OTT ^b	840 ka	Toba caldera, Sumatra	500	Diehl et al., 1987; Chesner, 1998
Bishop Tuff	700 ka	Long Valley, California (USA)	450	Mason et al., 2004
Campanian Ignimbrite	33.5 ka	Campi Flegrei, Bay of Naples, Italy	80	Rosi et al., 1996
MTT	501 ka	Toba caldera, Sumatra	60	Chesner, 1998
Tambora	1815 AD	Lesser Sunda Islands, Indonesia	50	Self et al., 1996
Krakatau	1883 AD	Sunda Strait, Indonesia	10	Self et al., 1996
Pinatubo	1991 AD	Luzon, Philippines	5	Self et al., 1996
Vesuvius	79 AD	Bay of Naples, Italy	1.5	Gurioli et al., 2005
Mount St Helens	1980 AD	Washington (USA)	0.55	Carey and Sigurdsson, 1982

aThe background behind the estimates obtained by Lee et al. (2004) is discussed in the text. Their DRE estimate combines the estimate of 500km³ for terrestrial OTT, with an estimate of 310–590km³ for distal air-fall ash. The latter estimate is calculated using evidence from layer D in the Indian Ocean deep-sea core, ODP Site 758, and 3 marine cores in the South China Sea. Lee et al. (2004) believe that layer D and the South China Sea ash are OTT. bOTT date and volume estimates prior to the work of Lee et al. (2004). The date of 840 ka was obtained by Diehl et al. (1987), using ⁴⁰Ar/³⁹Ar.

eruption. The oldest fossil from India is represented by the Narmada hominine (see Acharya, this volume), dated to not less than 236 ka (Cameron et al., 2004), or to some time in between 150 and 250 ka (Kennedy, 2001:167). Modern human remains have been discovered in an undated Late Paleolithic context at Bhimbetka rock shelter III-A-28 (Wakankar, 2002:5) and from three cave sites in Sri Lanka, dating from 27.7 ka (Kennedy 1999, 2001).

Investigations of modern human genetic variation have been providing some important insights into the timing of modern human dispersals into the Indian subcontinent (see Endicott et al., this volume), yet it does not provide information on other hominine species that may have inhabited India at the time of the Toba event. A synthesis of modern genetic data illustrates that the initial colonization of India by *H. sapiens* took place after the Toba eruption via a southern dispersal route from Africa, through southern Arabia and into India. Various dates have been obtained for the colonization of India; ~50 to 60 ka



(Quintana-Murci et al., 1999), 45 to 50 ka (Underhill et al., 2001), 60 ka (Kivisild et al., 2003), 60 to 70 ka (Palanichamy et al., 2004), 65 ka (Macauley et al., 2005), and 50 to 70 ka (Thangaraj et al., 2005). It should be noted how close some of these quoted dates are to the age of the Toba eruption. Evidence for pre-Toba migrations to India is provided by Oppenheimer (2003) and possibly by Forster and Matsumura (2005), the latter giving a timeframe of between 85 and 55 ka for dispersals to India. Using phytogeographic data, Oppenheimer (2003) argues that *H. sapiens* occupied India before ~74 ka and may have undergone “mass extinction” as a result of the Toba eruption (Oppenheimer, 2003).

A comprehensive review of the Late Pleistocene archaeological record of the Indian




Figure 3. Eruptive volumes (Dense Rock Equivalent (DRE) in km³) of a number of well-known volcanic eruptions

Subcontinent has recently been provided by James and Petraglia (2005), with particular focus on the colonization of the subcontinent by *H. sapiens*. They argue that two distinct lithic industries characterize the Late Pleistocene in South Asia; an earlier Middle Paleolithic, beginning ~150 ka (Misra, 1995), and a later technological complex (“Late Paleolithic”), beginning ~45 ka. They find no evidence of a symbolic (or Upper Paleolithic) revolution in the Indian record that can be said to be associated with the appearance of the first modern humans. Instead, they tentatively date the colonization of India by *H. sapiens* to ~70-50 ka and indicate that these early colonizers may have used Middle Paleolithic technologies. In this scenario, distinguishing assemblages created by *H. sapiens* from those made by other hominine species, who also produced Middle Paleolithic artifacts, is considered problematic (James and Petraglia, 2005). Therefore, in the broadest sense it is possible to predict that hominine in India were practicing Middle Paleolithic technologies at the time of the Toba eruption, however, it is not possible to discern which hominine species manufactured them.

Archaeological evidence from the Indian subcontinent can only provide evidence of the impacts of the Toba eruption on hominines if Paleolithic remains are preserved in either extremely well-dated contexts or in direct association with deposits of the Youngest Toba Tuff. Unfortunately, the former are currently a rarity in India; however, a number of sites exist in the Indian subcontinent where Toba tephra is preserved together with Paleolithic assemblages.

Quaternary deposits and Paleolithic site and Stone Age tools.

In several river valleys preserve tephra deposits that are directly associated with stone artifacts that pre-date and post-date the eruption. While Indian YTT deposits are relatively well-described in the geological literature, the archaeological significance of any associations between ash and Paleolithic remains has yet to be explored. There is currently very little published information which describes these associations.

The Central India Narmada River has conspicuously has straight, course as it is controlled by E-W lineament. It is bound by Vindhyaachal in the north and Satpura range to the south; the area in between these two upland is found to be ideal for a study of Quaternary sedimentation as witnessed by the presence of multicyclic sequence of Quaternary terraces in the valley. These are well developed in the Central Narmada valley stretch over 300 km, from Jabalpur in the east Harda in the west. The width of the valley varies from 20 to 80 km. The Quaternary deposits are 400m thick, however, only the upper 50 m has been exposed in river section. The boulder conglomerate exposed at Hathnora ($22^{\circ}52'77''52'$) has yielded skull of early man (*Homo erectus* skull) of undisputed Middle Pleistocene hominine remains Sonakia (1984)

The Quaternary deposits of Narmada valley in the area Hominid Fossil locality Hathnora consists of 70 to 90 m of Narmada alluvium was deposited in single aggradational episode with minor pauses when dissection of the alluvium produced two terraces (NT1-NT2) The Sediments of gradational episode constitutes three lithostratigraphic units viz Boulder conglomerate, Sohagpur Formation. A Boulder bed of glacio-fluvial origin underlies the sediments of the alluvium phase. Thus the fossiliferous boulder conglomerates, the basal unit of the alluvium marks disconformities between the lower glacio – fluvial boulder layer and the upper fluvial sediments. The fossiliferous boulder conglomerate is considered to be middle Pleistocene age (Khan & Sonakia 1992). The Boulder Conglomerate (Hathnora Formation) is thick wedge of semi consolidated fossiliferous sediments and concealed under thick deposits of clay and silt of the older alluvium in valley. It may contain the richest vertebrate fossil assemblage and has yielded only known Hominid fossil from the Indian sub-continent (Sonakia 1984). The Boulder bed which yielded Hominid fossil for the first time reported to be of glacial & fluvial origin Khan & Sonakia (1992). The fossiliferous horizons is concealed wedge,

it is exposed only at some places in valley by lateral dissection by superimposed drainage, lateral river cutting, dissection of quaternary blanket, and other erosional activities. The concealed and hidden disposition of fossiliferous horizon and its mysterious intricacy in tectonic zone and due to failure to understand basin configuration has disappointed the geoscientists to add further to any new find of Human skull, in spite of keen interest and intensive and extensive work. This formation is associated with various stone tools and stone implements like scraper, cutter, axe, blade with unfinished tools and scrap material. The area around homonid fossil possesses sites of Stone Age. The gravel that yielded the Narmada Homo erectus carries heavy hand axes, cleavers and scrapers. The critical three dimensional study of the area around Hathnora (22°52'77°52') between Vindhyaachal and Satpura mountains vast industry sites of the same palaeolithic implements have been located along the edge and pediment slope of the Vindhyan and basaltic upland. These sites appear to have been inhabited by Narmada hominid and its contemporaries for generations together.

The Narmada Homo erectus and associated Multi Layered Volcanic Ash.

The early human settlements have left widespread signatures on Indian River valley systems in the form of extensively large Paleolithic industry sites and with association of Stone Age tools, whereas the hominid fossil evidences are very rare and scanty. The skull cop of *Homo erectus*, the 'Narmada Man' (Fig 20) is the only example in the long stretch of Quaternary alluvial deposits occurring at varying topographic levels in the Narmada river valley of Central India. Some of the terraces are pretty rich in Paleolithic and Neolithic cultural materials, bearing testimony to the hunting and gathering abilities of early man. Elsewhere in the world, a century of discoveries precedes the discovery of the Indian fossil in 1982. The pioneering discoveries succeeded, with a couple of few finds from java in Indonesia and from China. Later, when the search involved the continents of Africa and Europe there was a spurt in the rate of discoveries and as a result, a large number of human fossils were discovered, thereby making the study of human evolution more logical and factual. It has to be mentioned here that initially discoveries of human fossils from Germany in Europe and from java in Asia, were very much neglected. The Indian fossil mentioned by Theobald (1881) was lost due to sheer curatorial negligence (Kennedy, 1980), whereas the Garman Neandethal specimen was initially misinterpreted.

It took time to describe the Narmada specimen in detail and it also invited controversies regarding its taxonomic. (De Lumly and Sonakia, 1985; Kennedy et al, 1991).

Quaternary Tectonics Evolution, Sedimentation, and Paleo- anthropological
Records of Impact of Ash fall on Pleistocene Hominines in Narmada Rift System
(M.P.) Central India

1314

1315 The Main Narmada rift is both symmetrical and asymmetrical in different segments
1316 along its length of about 1300 km. Several paleo- anthropological localities,
1317 archeological sites ranging in age from the Pliocene-Pleistocene times were discovered
1318 within these basins. The discovery of Human Skull *Homo erectus* form boulder
1319 conglomerate bed of Hathnora formation Khan (1992) by Sonakia (1984) De Lumley,.
1320 and Sonakia, (1985): in Sehore district M.P.India was first fossil skull of man from
1321 Indian sub-continent. It is correlated with *Homo-eructs* of China on Quaternary
1322 Platform is found to be the oldest homo-erectus in Asia Khan et.al (2013)

1323

1324 The known Pliocene-Pleistocene paleo-anthropological localities have given us
1325 information about ancestors who were habitants and sparsely concentrated in the
1326 Narmada rift valley. This is not a coincidence, because the volcanic and tectonic
1327 activities that were responsible for the formation of the rift basins and formed the loci
1328 of Quaternary sedimentation & created ideal environments for the proliferation of life
1329 and the preservation of faunal and floral remains. The Quaternary volcanic eruption, ash
1330 fall, repeated tectonic dislocation and were responsible for the quick burial and
1331 preservation of fossils during diagenesis. The assemblages of sediments and
1332 granulometric parameters, diagenetic processes involving silicification, calcification,
1333 feldspathization, clay formation, and pedogenesis all played vital roles in fossil
1334 preservation in the sediments. The various rock fabrics, ash bed, paleo- sole inter
1335 bedded with the fossiliferous sediments also provide temporal information about
1336 geologic processes, faunal evolution, paleo -environment, and early hominid behavior
1337 and lithic technology.

1338

1339 The skull cap of Narmada Man *Homo erectus* was found in Narmada Valley near village
1340 Hathnora (22 ° 52" N; 77 ° 52" E) in fossiliferous boulder conglomerate, in district
1341 Sehore, M.P., India. The skull cap is completely fossilized undistorted, renal vault nearly
1342 complete except few left Supra-orbital and statures are nicely preserved. The various
1343 morphological features and robust form of skull and excessive thickness of the bones
1344 indicate that it belongs to adult male individual (Sonakia, 1984). The discovery of skull
1345 cap of *Homo erectus* in fossiliferous boulder conglomerate in association of other
1346 mammalian fossil is recorded in stratigraphic column of Quaternary deposits at the depth
1347 of 83 m , where estimated total thickness of deposits is about (325 m).This blanket
1348 consist of sediments of three domain viz. glacial, fluvio-glacial and fluvial, which were
1349 deposited in distinct environment during Pleistocene to Holocene time (Khan & Sonakia
1350 (1992), (Khan et.al. in press).The statistical analysis of sediments from these different
1351 domain in vertical column has been conducted to ascertain the environment of
1352 sedimentation and trace the breaks in climate (Khan & Aziz 2015)). An attempt has been
1353 made for the first time Khan et.al (2013) to correlate the various stratigraphic columns
1354 of associated hominid fossils of Narmada valley (325 m) India and that of
1355 Luochuan sequence,(90-120 m) Chenjiawoe (50m) and Congwanling sequence (36 m
1356) of China on unified Quaternary platform tied up and developed at mean sea level.
1357 The study revealed that the depth of occurrence of Narmada skull cap on unified
1358 Quaternary platform is about (83 m) as compared to with that of Chenjiawo and
1359 Gongwangling of China which occur at very shallow depth of 38 and 26 m

respectively. The estimated age of Narmada Man based on these parameters is about 1.38 m.y. (+), which is greater than *Homo erectus* of Chenjiawo 0.65 m.y. and Gongwangling 1.15 m.y. of China An Zhisheng and Ho Chuan Kun (1989). On the merits of correlation of stratigraphic columns of Quaternary of Narmada, accumulation of sediment, rate of sedimentation, palaeo- environments, lithostratigraphy and biostratigraphic position of boulder conglomerate in unified Quaternary Platform, author consider it as one of the earliest and oldest *Homo erectus* in Asia. Khan et.al (2013).

Khan (1984) carried out detailed geological, geomorphological and sedimentological studies and identified three prominent terraces in the Central sector of the Narmada. (Khan Sonakia 1992), studied Quaternary deposit in vertical column in relevance to occurrence of Human Skull and reported glacial and fluvio- glacial deposit in Narmada valley for the first time. The boulder bed which yielded Hominid fossil for the first time reported to be of glacial - fluvial origin Khan & Sonakia (1992). Beside occurrences of association of ash beds with fossiliferous boulder conglomerate Khan & Rahate (1991) indicates some distant volcanic source. .

Khan and Sonakia (1992) reported for the first time glacial and interglacial deposit in the Narmada valley, Central India which is represented by arid and humid cycles. Magneto- stratigraphic studies are being carried out by Rao, et.al (1997). The lithostratigraphy of Narmada valley described by Khan (1984), Khan & Benarjee (1984), Khan & Rahate (1990-91), Khan & Sonakia (1992), Khan *et al* (1991), Rahate & Khan (1985), Khan (1991), Khan & Sonakia (1992), Yadav & Khan (1996).

In Narmada Rift system presence of the Katni Formation with angiosperm flora suggests that sedimentation continued during Mio-Pliocene in localized lakes. The relative disposition of such lakes and subsequent deformation and structural dislocation on oscillating valley platform clubbed with rifting and faulting during Quaternary period has shifted the site of the lakes towards the present alluvium-covered area between Harda and Jabalpur as presumed: where as the present study of various aspects of Quaternary blanket in SONATA lineament ZONE revealed that quaternary sedimentation was a sequential and continuous process in rift valley system from Mio-Pliocene Pleistocene time, has deposited complete sequence of glacial, fluvio-glacial lacustrine and fluvial deposits with changing environments and climate in time chronology. The present disposition of quaternary blankets in Son Narmada Tapti and Purna basin is due to post deposition Quaternary tectonics which is solely responsible for dislocation, faulting and shifting of different blocks and distorting quaternary blanket ecology in rift system. The occurrence of Boulder bed and Boulder Conglomerate in Son Narmada Tapti and Purna with similar rock assemblages and suites of rock fabrics, heavy mineral assemblages, quartz grain morphology in critical and crucial sections across the SONATA LINEAMENT ZONE strongly support tearing and rifting of quaternary blanket during late Pleistocene time. The presence of thick boulder bed in Harda inliers area, such as at Chandgarh and north east of Barwaha, boulder bed in confluence are of Tapti and waghur around Khadgaon in Tapti valley Khan et.al (1984) supports this assumption.

The present work was initiated in 1982 and continued over a decade, during the tenure tremendous data on various aspects was of Quaternary geology, sedimentology, pedology, archaeology, tectonic, neotectonic Ash bed and other related studies were carried out and accordingly acquired and accrued tremendous data from the ground by systematic field work, by working on official assignment as part of government duty. The virgin and previously unknown areas of the rift basins were studied and inventory of paleoantological and paleoanthropological Ash bed resources were made. The study indicated the potential of the Mio-Pliocene Pleistocene time and late and early Pleistocene Quaternary sediments of the Narmada Rift System for paleoantological paleoanthropological Ash bed association its impact & research.

There is a strong link with association of Ash bed and ash fall environment and association of faunal and floral remains especially in the Pliocene–Pleistocene sedimentary strata. The time-stratigraphic data obtained from tephra interbedded with fossiliferous Quaternary sedimentary deposits provided an important framework for the study of hominid origins, evolution, adaptations, environment, climatic and impact of ash fall in time and space.

The paleoantological information from these localities is remained closely associated with Quaternary sedimentary deposits boulder conglomerate and boulder bed and Ash bed often related to the trench Quaternary sedimentation, formation and development of rift and linear basin caused by repeated uplift, and the development of rift basins that began in the middle to late Pliocene and Pleistocene period. The unfortunate part of these deposits is that due repeated tectonic dislocation and faulting they are dislocated and distorted and at present are concealed under the thick pile of sediments of present and paleo domain of Narmada of late Pleistocene and Holocene time. These deposits do not provide adequate opportunity to researcher to study the human remain as postulated, except in limited section where they are exposed.

In Narmada valley the most of the hominid remains and associated artifacts in the would have been found associated with Miocene Pliocene–Pleistocene sediments in increasing antiquity unfortunately same are not exposed due rift system and tectonic setting. In the rift system the type development of Quaternary blanket is confined between Jabalpur–Harda section, which posses the complete sequence of all three domain in increasing antiquity from the bottom of the rift trench Boulder bed (glacial), Boulder conglomerate (fluvio-glacial) sediments of paleo-domain of Narmada (fluvial). The intense tectonic activities within the basins of the Narmada Rift System during the Neogene and Quaternary periods have destroyed fossil record except the fossiliferous horizons exposed in river sections. The erosional-sedimentary cycle has persisted in the rift valley environment for millions of years as a result of the interplay between depositional and erosional forces driven by tectonic processes; there are numerous gaps in the fossil record, particularly in the important time period between Mio-Pliocene Pleistocene times.

It is pertinent to the understand the origin of Hominid during the late Miocene, but it is difficult to disclose mysteries of human evolution in Narmada due to concealed nature of these deposits in rift system, however the complementary part of Tapti-Purna Quaternary blanket may be potential and possessive of human remain and should be studied to trace further the imprints of fossil man taking in to account of SONATA LINEAMANT ZONE as single ecosystem for evolution of man in Indian subcontinent.

Moreover, because of tephra layers in sedimentary basins of different geologic periods, processes such as faulting, rifting, sedimentation and diagenesis, impact of climatic changes, age of fossils, nature and acquisition of archeological implements, and the origin, distribution, and functional significance of early hominid artifact assemblages can be deciphered. However, evidence of the effects of tectonics on fauna and flora are distinct and its signatures on dislocation and concealing of fossiliferous horizons are uncontrolled and ill defined in the ecosystem in the valley during the Pliocene–Pleistocene periods. The boulder conglomerate which yielded the skull cap of *Homo erectus* in Narmada rift from Hathnora Sonakia (1984) remained only discovery of hominid fossil in last two and half decade due to concealed and hidden nature of Mio-Pliocene Pleistocene deposits in rift system and inconsistency in exposure of fossiliferous horizon due faulting, dislocation and subsidence of Quaternary blanket of Narmada rift system.

The intense tectonic activities within the basins of the Narmada Rift System during the Neogene and Quaternary periods have destroyed fossil record except the fossiliferous horizons exposed in river sections. The erosional-sedimentary cycle has persisted in the rift valley environment for millions of years as a result of the interplay between depositional and erosional forces driven by tectonic processes; there are numerous gaps in the fossil record, particularly in the important time period between Mio-Pliocene Pleistocene times.

It is relevant to the understand the origin of Hominid during the late Miocene, but it is difficult to disclose mysteries of human evolution in Narmada due to concealed nature of these deposits in rift system, however the complementary part of Tapti-Purna Quaternary blanket may be potential and possessive of human remain and should be studied to trace further the imprints of fossil man taking in to account of SONATA LINEAMANT ZONE as single ecosystem for evolution of man in Indian subcontinent. On the merits of correlation of stratigraphic columns of Quaternary of Narmada Rift System, accumulation of sediment, rate of sedimentation, palaeo- environments, lithostratigraphy and biostratigraphic position of boulder conglomerate on unified Quaternary Platform, author consider it as one of the earliest and oldest *Homo erectus* in Asia. Khan et.al (2013).

The boulder conglomerate of fluvio-glacial origin which yielded human skull from Hathnora formation Sonakia (1982) Khan & Sonakia (1992) is exposed impermissibly in scarp section of Narmada at few places only. The type section of Boulder Bed and Boulder Conglomerate which are potential sediments of human remains of Pliocene

Pleistocene time are hidden and concealed under sediments of present and paleo domain of Narmada in the valley. The quaternary sequence and depth of occurrence of skull cap, rate of sedimentation of Hominid locality Hathnora (22°52'77°52') (325 m) India, has been correlated with the Quaternary of Luochuan (90-120 m) Chenjiawo (50m) and Congwanling sequence (36 m) of China on unified coded model of Quaternary platform tied up and developed at mean sea level to ascertain the depth of occurrence and of age of skull cap of Narmada *Homo erectus* Sonakia (1984), at the rate of sedimentation (83 m) Khan et.al (1213) which indicates that the human skull *Homo erectus* of Narmada is oldest in Asia which lived survived and sustain prior volcanic eruption and ash fall..

The central Narmada valley possess rich Palaeolithic occurrences about 300 Lower Palaeolithic occurrences have been reported in different contexts. The present work deals with these assemblages and associated raw material sources in central Narmada valley. The Central Narmada valley is rich in various rock types which were used as raw material by various hominine populations. The Narmada River cut across Indian subcontinent negating its course along SONATA lineament zone dividing in to dividing the peninsular India into two geological provenances. Whereas, in the south, there are primarily the Gondwana Supergroup, Deccan Trap and the Mahakoshal Group. The Central part is occupied by thick sequence of Quaternary deposit. The main raw material types in the north of Narmada are quartzite and sandstone (Vindhyan Supergroup), whereas, in the south of Narmada, the main raw materials are quartzite (Gondwana Supergroup) and chert (Deccan Trap). Acheulean sites are mostly found along the foothills of Vindhyan as well as along the banks of the Narmada River and its tributaries. It is attempted that to correlate these material sources with occurrence of Paleolithic sites to have a better understanding of past human activity and environment in which the hominine resided. The area become more interesting after the recovery of *Homo erectus* from boulder conglomerate from Hathnora by Sonakia (1982)

In Central Narmada valley and around Hominid Locality Hatnora 330 occurrences have been reported, which signifies its importance in palaeo-anthropological which have imprints of human activity. The sites are located along northern bank on pediment surface of Vindhyan. The Narmada and Sonad Tapti Rivers are three of the most important geographic features that have shaped the course of Acheulean adaptations in this region of the subcontinent, The central sector of Narmada which embodies steeped sequence of alluvial which possess good quality of raw material.

Vindhyan Supper Group Of Rocks

The Vindhyan Super group is one of the largest and thickest Proterozoic sedimentary successions in the world, spanning an exposed area of 60,000 sq km and extending invisibly under the Gangetic Alluvium and Deccan Traps. It mainly consists of non-

1537 metamorphosed sandstone, shale, and limestone ranging in age from 1.72 to 0.75 billion
1538 years:

1539 The Vindhyan basin is a classic example of Proterozoic intracontinental basin that
1540 developed in the central part of the Indian shield along with several other basins such as
1541 Cuddapah, Chattisgarh, etc. The strata are exposed in three major sectors: Son valley,
1542 Bundelkhand and Rajasthan. Substantially thick Vindhyan rocks have also been
1543 recognized under the Gangetic alluvium.

1544 The Vindhyan Basin, containing more than 5000 m thick sequence of sandstones, shales
1545 and limestones, occupies an area of about 1,62,000 sq.km of which about 80,000 Sq.km
1546 extends into the Ganga valley in the north and northeast beneath the Tertiary sediment of
1547 the Himalayan foredeep. In the southwest, the Vindhyan rocks are covered by Deccan
1548 volcanics.

1549 The basin is bounded by the Son-Narmada Geofracture in the south, the Great Boundary
1550 Fault in the west, the Monghyr-Saharsa Ridge in the east, and the Bundelkhand Massif
1551 and Indo-Gangetic Plains in the north. Bundelkhand Massif divides this basin into two
1552 parts – the Son Valley on the southeastern side and the Chambal Valley where exposures
1553 occur from Agra (Uttar Pradesh) to Chittorgarh (Rajasthan).

1554 The southern margin of the Vindhyan basin is marked by a major ENE–WSW trending
1555 lineament termed Narmada–Son lineament south of which occurs the Satpura orogen. A
1556 NE–SW trending major lineament known as Great Boundary Fault separating the
1557 Aravalli–Delhi orogen from the Vindhyan, also marks the western margin of the
1558 Vindhyan basin in the Rajasthan sector.

1559 The Vindhyan strata define a broad, regional syncline trending ENE–WSW. The axis of
1560 the syncline is slightly curved and plunges gently towards west. The average dip of the
1561 southern limb is greater than that of the northern limb rendering the axial plane to dip
1562 southerly.

1563 The Vindhyan Basin was formed as a result of a large crustal downwarp in the northern
1564 part of the Indian Platform, after the Delhi orogeny. The initial transgression of the sea
1565 from the north is inferred to have taken place in the eastern part of the basin over the
1566 Bijawars. The shallow sea appears to have established lagoonal conditions near the
1567 coastal part during the subsequent regressive phase. Initially, the Son-Narmada
1568 Lineament was dormant, but at the onset of Vindhyan sedimentation later, the fault
1569 system along this downwarp became active with the formation of the southern limit of
1570 Vindhyan deposition.

1571 After the deposition of Kajrahat Limestone, the Son-Narmada Lineament again became
1572 active resulting in emission of volcanic material, which was deposited as the Jardepahar
1573 Porcellanite. In the subsequent regression, the shore line shifted towards northwest. The
1574 Fawn Limestone was deposited over the shelf in a tidal flat environment. This was

1575 followed by shallowing of the basin as is evident from the overlying Glauconitic
1576 Sandstone. The fresh marine transgression resulted in the deposition of marine shales
1577 followed by Rohtas Limestone.

1578 The entire Vindhyan succession, maximum thickness estimated to be around 3 km, and
1579 comprising mainly sandstone, shale and limestone is assigned as the Vindhyan
1580 Supergroup. The Supergroup is divisible into four groups:

- 1581 1. Semri Group,
- 1582 2. Kaimur Group,
- 1583 3. Rewa Group and
- 1584 4. Bhander Group,

1585 from bottom to top. Each group is again subdivided into several formations. The Semri
1586 Group in the Son valley rests unconformably on a variety of pre-Vindhyan rocks such as
1587 granites and metamorphics. In the Bundelkhand area, the group overlies the Bundelkhand
1588 Granite Gneisses and Bijawar Group of metamorphics, whereas in the southern Son
1589 valley, Mahakoshal is the basement in most places; however, in some localities (e.g.,
1590 Deoland, M.P.) the basement is granite. The Semri succession of the Bundelkhand area
1591 has two detached outcrops around Chitrakut and Bijawar respectively, and is only a few
1592 tens of meters thick.

1593 The Vindhyan supper Group of rocks in eastern extremity around Bhopal inliers and
1594 southern arm of these rocks on the obsequent slope embodies Bhimbetka shelter mostly
1595 the rock cavities of intermeddled Quartzite lime stone sandstone and shale which have
1596 been which were exposed to post erosinal and depositional activities to million and
1597 million years resulting circular hollow and cylindrical cavities of lime stone and shale
1598 interbedded strata both along and across bedding planes which have created natural
1599 shelters in Vindhyan rocks, These were occupied by migrants Hominess in event of
1600 volcanic eruption and dense atmospheric contamination and fall of ash in subcontinents..
1601 The cluster of these rock shelter was discovered in 1957-58 by Dr V,SWakankar.

1602 The Rock Shelters of Bhimbetka are located about 45 km south east of Bhopal on the
1603 road to Hoshangabad. The site spread over 10 km in length and about 3 km in width has
1604 more than 700 rock shelters, of which over 400 have paintings. The earliest human
1605 activities are known from the numerous stone tools including hand axes, cleavers and
1606 also the pebble tools.
1607

1608 The latter pertinently was found in primary contexts. The continuity of human evolution
1609 from the Lower Palaeolithic Period is noticed by the smaller size of stone tools in the
1610 following Middle Palaeolithic Period besides new tools like scrapers. During the Upper
1611 Palaeolithic Period newer tool types like: blades, borers and bruins had also emerged.
1612 However, it is in the Mesolithiic Period that there is a clear change in the materials and


tool typology. Earlier, the tools were largely made of quartzite and sandstone, whereas the tools being made in the Mesolithic Period were most often of chalcedony. The stone tools of this period include blades, triangles, trapezes, crescents besides quern and muller.

The Mesolithic culture at Bhimbetka continued much longer as understood by the presence of Chalcolithic potteries in otherwise Mesolithic contexts. By the Early Historic times it appears that interaction with the surrounding cultures became more pronounced. This is evidenced by the presence of rock-cut beds in a rock shelter on the top portion of an inselberg like outcrop not far from the later built temple at this site. In all respect it resembles the Sallekhana spots observed in South India.

The cave also has a small inscription of the Maurya/Sunga period. Within the general area of Bhimbetka Group of rock shelters small stupas have been found at Bhimbetka, near Bhoranwali, at Bineka, at Lakhajuar and midway between Lakhajuar and Bhimbetka. At Bineka besides the stupa enclosure wall and other structures were found. There are a large number of Shanka Lipi inscriptions in the Bhimbetka cluster of rock shelters.

However it is the rock paintings of Bhimbetka that steals the limelight. of particular interest to the tourist are the Auditorium Rock Shelter, Zoo Rock and Boar Rock in Bhimbetka Cluster. The earliest endeavor here other than for mere run of the mill activities for survival is the engravings of small cup like depressions at the end of the Auditorium Rock Shelter, which is dated to nearly 100000 years. Near the end of this tunnel there is a cluster of painting depicting a hunter, deer, tiger cattle and stylized peacock. Further ahead on the same path one comes across the Zoo Rock Shelter, which qualifies as the most densely painted rock shelter, paintings spanning from the Mesolithic to the Mediaeval. The paintings here include those of A Mesolithic boar painted in dark red, animals like: elephant, rhinoceros, boar, barasingha, spotted deer and cattle and snake, etc. Later paintings include battle scenes painted in red and an elephant painted in white. The Boar Rock, which is the last among the rock shelters

accessible for tourist has a depiction of a mythical boar with horns that is many more times larger than the human being chased by it.



CONCLUSION

The study of the area in Central Narmada valley around Hathnora (22°52'77°52') between Vindhyachal and Satpura mountains vast industry sites of the same palaeo lithic implements have been located along the edge and pediment slope of the Vindhyan and

basaltic upland. These sites appear to have been inhabited by Narmada hominid and its contemporaries for generations together the left over stone industry sites and unfinished material haphazardly indicate sudden dissemination and migration of hominines in search of shelter and safe places after volcanic eruption as also documented by the relicts of ruins Paleolithic and Mesolithic industrial site and scattered occurrence of stone implements around Hominid locality of Hathnora

The area possesses rich Palaeolithic occurrences about and 300 Lower Palaeolithic occurrences have been reported in different contexts. The present work deals with these assemblages and associated raw material sources in valley .These rock types were used as raw material by various hominines populations. The area appears to be thickly inhabited and influence by human activity which was adversely affected by severe Toba volcanic eruption and hominines were forced to be migrated in struggle of survival to the safe places. The nearest safe place was Bhimbetka about twenty kilometers in the north located in consequent slope of Vindhyan rocks of Bhopal inliers which possess about 700 rock shelters of different size and dimension. The study of human activity in the area around Bhimbetka indicates the hominines populations lived around 100000 years.

The 74,000 year-old super eruption of the Toba volcano, located in northern Sumatra, is recognized as one of Earth's largest known eruptions and was certainly the largest of the Quaternary period (Smith and Bailey, 1968). It is presumed that it have led to both global climatic and environmental deterioration and had a impact on decimation of modern human populations (Rampino et al., 1988; Rampino and Self, 1992, 1993a) (Rampino and Self, 1993b; Ambrose, 1998, 2003a, 2003b; Rampino and Ambrose, 2000). However, the severity of Toba's impact on climate and hominines has been contested and debated by scientist (Oppenheimer, 2002; Gathorne-Hardy and Harcourt-Smith, 2003). Geological, paleontological and archaeological evidence from the Indian subcontinent provides an excellent opportunity to address these issues. The intensity and scale of the Toba super eruption was multidimensional and its column of eruption was so vast that it forma canopy of volcanic matrix led to the deposition of a blanket of volcanic ash across the continents and river valleys over India, Malaysia, the Indian Ocean, and the Arabian and South China Seas. Resulting persistent terrestrial tephra deposits have been documented in a number of river valleys throughout India. (Khan et.al 1991 Acharyya and Basu, 1993; Shane et al., 1995; Westgate et al., 1998).

The occurrence of Bolder conglomerate and occurrence of haphazard and disserted sites of Palaeolithic sites indicates that area suffered dual hostility first the end of glacial activity with the stratigraphic column at the level of 83 m where the Homo erects survived Sonakia (1984) both by glacial activity and volcanic eruption, whereas in the second hostility evolving human activity by volcanic eruption and dense smoke and dust and darkness. And consequent upon H Sapiens and associated were forced to migrate to

safer places. In view of the recorded observation the concealed quaternary strata to the level of about 350 m below ground level further needs attention of geoscientist for the search of human remains in Narmada valley.

The argument of Oppenheimer (2003) is in strong conformity with the present observation of authors. As sediment & Ash bed sequence of Quaternary column of Narmada (325 m) and occurrences of fossil of skull cap of *Homo erectus* (Sonakia 1984) at 83 m & Human cranium *Homo sapiens* (Thebold 1960, 1981) transported have existed prior to fall of Toba ash and they are among the few who inspite of mass extinction caused by mega dislocation in ecology and environment related with volcanic eruption survived in Narmada Valley. It is further documented by the rarest occurrences of these fossils in subcontinent which also confirm the intensive impact of volcanic ash fall on these hominines and their consequential mass extinction.

The Toba eruption of 74 ka was distinctly and clearly a mega event of very great magnitude and intensity, far greater than any known historical eruption, suggesting it had very devastating impact and repercussions. It has change the global climate environment and ecology.

The association of Ash bed NAB-I with Hathnora formation at the depth of 78 m in Quaternary column of Narmada and occurrences skull cape of *Homo erectus* of Narmada at the depth of 83 m assumed that Toba eruption might have taken place later than that of existence of *Homo erectus* which appeared and resided in the valley for long time before the fall of Toba ash. The association of Ash bed NAB-II at the depth of 72 m with the younger deposits indicates the second cyclic of fall of Toba ash which might have influenced the younger Hominines i.e. *Homo erectus* & *Homo sapiens*, in Narmada valley.

The occurrences of these skull caps with short range of their occurrences in the stratigraphic column of Narmada with the Ash beds horizon NAB-I and NAB-II and specially with the Hathnora formation one at the top at an average elevation of about 268-273 m above the mean sea level and other with younger deposits had revealed the close association with volcanic activity with their existence. The Toba Ash fall is also in very close range with the sequence of sedimentation and occurrences with both the skull caps, which certainly has its impact on the middle and late Pleistocene Hominines in Narmada valley and Indian subcontinent.

The Hominines residing in Central Narmada valley after eruption of Topa and sudden decline in climatic ecology rapidly migrated to the nearest rock shelter of Bhimbetka in the north leaving behind hapazrdly their tool making sites, raw material unfinished tool and other belonging behind. The Bhimbetka posses about 700 rock shelters. of different size and dimentions and shape within the periphery about eight kilometers. The relict of

human activity, tools, rock paintings, are recorded in these shelter where they resided about 100000 years.

. It is reported that remains of *H. sapiens* in the area of Bhembetaka north of hominid locality of Hathnora and relicts of rocker shelter and ancient human signature indicate the dispersal of Hominines after the eruption of volcanic matrix and its fall Wacahlakar, V.S., 2002. The association of skull cap Sonakia, A (1984) and Sankhyan (2007) described two new hominine fossils from Netankheri up stream of Hathnora in the Central Narmada valley. They include a partial left humeral diaphysis and a distal shaft fragment of the left femur associated with Hathnora formation Khan & Sonakia 1992. The analysis of quartz grain morphology of sediment columns of Hathnora and quartz grain morphology of paleo-soil across the quaternary strata of bore hole sampling across 556 m of rock basin revealed that *Homo erectus* of Narmada partly sustain in the glacial and fluvio-glacial environment as witnessed by Hathnora formation (Boulder & gravel bed) which marks the end of hostile climate and environment of sedimentation in Narmada valley Khan (2013).

The study of quartz grain and their micro structures of paleo-soil identified in concealed blanket of quaternary deposits display relatively heterogeneity in Sediment characteristics throughout across the Quaternary column of Hathnora section in central Narmada valley. The quartz grain of paleo-soil and sediments display diagnostic characteristics of glacial fluvio-glacial and fluvial concealed environment of sedimentation at different depth. The significant breaks in grain morphology of quartz grain surface texture and associated elements and granular matrix in the sequence of quaternary strata is recorded at 150, 350 and 350 and beyond which linked to long evolution of glacial fluvio-glacial and fluvial environment of sedimentation in time and space in increasing antiquity in the valley from the base Khan (2013). In view of the recorded observation the concealed quaternary strata to the level of about 350 m below ground level further needs attention of geoscientist for the search of human remains Ash bed in Narmada valley.

The atmosphere contaminated by volcanic ash volcanic matrix and huge amount of gases and dust particles had definitely affected the atmosphere and climate had have led to temporarily darkened skies in the Narmada valley and Indian subcontinent for very long time.

The study of grain morphology of glass matrix, their relation with other minerals shape, size, texture of lithic fragments and association of other ashy sediments of pyroclastic origin suggest that sediments were brought from distant source by Aeolian agencies during the different phases of sedimentation in Narmada Valley.

References

- Acharyya, S.K., Basu, P.K., 1993. Toba ash on the Indian subcontinent and its implications for correlation of Late Pleistocene alluvium. *Quaternary Research* 40, 10–19.
- Acharyya, S.K., Basu, P.K., 1994. Reply to comments by S. Mishra and S.N. Rajaguru and by G.L. Badam and S.N. Rajaguru on “Toba ash on the Indian subcontinent and its implication for the correlation of Late Pleistocene alluvium”. *Quaternary Research* 41, 400–402.
- Acharyya, S.K., Lahiri, S., Raymahashay, B.C., Bhowmik, A., 2000. Arsenic toxicity of groundwater in parts of the Bengal basin in India and Bangladesh: the role of Quaternary stratigraphy and Holocene sea-level fluctuation. *Environmental Geology* 39 (10), 1127–1137.
- Acharyya, S.K., 2003. Recent findings of the Acheulian of Isampur excavations and its dating. *Current Science* 84 (2), 127–128.
- Aldiss, D.T., Ghazali, S.A., 1984. The regional geology and evolution of the Toba volcanotectonic depression, Indonesia. *Journal of the Geological Society* 141 (3), 487–500.
- Ambrose, S.H., 1998. Late Pleistocene human populations bottlenecks, volcanic winter, and differentiation of modern humans. *Journal of Human Evolution* 34, 623–651.
- Ambrose, S.H., 2002. Small things remembered: origins of early microlithic industries in Sub-Saharan Africa. In: Elston, R., Kuhn, S. (Eds.), *Thinking Small: Global Perspectives on Microlithic Technologies*. Archaeological Papers of the American Anthropological Association 12, pp. 9–29.
- Ambrose, S.H., 2003a. Population bottleneck. In: Robinson, R. (Ed.), *Genetics*, volume 3. Macmillan, New York, pp. 167–171.
- Ambrose, S.H., 2003b. Did the super-eruption of Toba cause a human population bottleneck? Reply to Gathorne-Hardy and Harcourt-Smith. *Journal of Human Evolution* 45 (3), 231–237.
- Anonymous, 1990. Late Quaternary ash bed and Acheulian implements from Orissa – the implication. *News Geological Survey of India, Eastern Region* 10 (1), 7.
- Anonymous, 1991. Volcanic ash in the Quaternary Formations Sagileru valley. *News Geological Survey of India, Southern Region* 9 (2), 9.
- Athreya, S., 2007. Was *Homo heidelbergensis* in South Asia? A test using the Narmada fossil from Central India. In: Petraglia, M.D., Allchin, B. (Eds.), *The Evolution and History of Human Populations in South Asia: Inter-disciplinary Studies in Archaeology, Biological Anthropology,*

1851 Linguistics and Genetics. Springer, Netherlands, pp. 137–170.

1852 Badam, G.L., Misra, V.D., Pal, J.N., Pandey, J.N., 1989. A preliminary study of

1853 Pleistocene fossils from the Middle Son Valley, Madhya Pradesh. *Man and*

1854 *Environment* 13, 41–47.

1855

1856 Basu, P.K., Biswas, S., Acharyya, S.K., 1987. Late Quaternary ash beds from Son and

1857 Narmada Basins, Madhya Pradesh. *Indian Minerals* 41 (2), 66–72.

1858 Basu, P.K., Biswas, S., 1990. Quaternary ash beds from Eastern India. *Records of the*

1859 *Geological Survey of India* 123 (2), 12.

1860

1861 Biswas, S., Dassarma, D.C., 1986. Stratigraphy of the Quaternary alluvial deposits of

1862 the Central Narmada Valley, Madhya Pradesh. *Proceedings of the XI Indian*

1863 *Colloquium on Micropalaeontology and Stratigraphy, Bulletin Geological Mining*

1864 *Metallurgical Society of India* 54, 18–27.

1865

1866 Blumenschine, R.J., Chattopadhyaya, U.C., 1983. A preliminary report on the terminal

1867 Pleistocene fauna of the Middle Son valley. In: Sharma, G.R., Clark, J.D. (Eds.),

1868 *Palaeoenvironments and Prehistory in the Middle Son Valley*. Abinash Prakashan,

1869 Allahabad, pp. 281–284.

1870

1871 Brandon-Jones, D., 1998. Pre-glacial Bornean primate impoverishment and Wallace's

1872 line. In: Hall, R., Holloway, J.D. (Eds.), *Biogeography and Geological Evolution of SE*

1873 *Asia*. Backbuys Publishers, Leiden, pp. 393–404.

1874

1875 Bühring, C., Sarnthein, M., Leg 184 Shipboard Scientific Party, 2000. Toba ash layers

1876 in the South China Sea: evidence of contrasting wind directions during eruption ca. 74

1877 ka. *Geology* 20 (3), 275–278.

1878

1879 Cameron, D., Patnaik, R., Sahni, A., 2004. The phylogenetic significance of the Middle

1880 Pleistocene Narmada hominin cranium from central India. *International Journal of*

1881 *Osteoarchaeology* 14 (6), 419–447.

1882 Carey, S.N., Sigurdsson, H., 1982. Influence of particle aggregation on deposition of

1883 distal tephra from the May 18, 1980, eruption of Mount St. Helens volcano. *Journal of*

1884 *Geophysical Research* 87 (B8), 7061–7072.

1885 Chesner, C.A., Rose, W.I., Deino, A., Drake, R., Westgate, J.A., 1991. Eruptive history

1886 of Earth's largest Quaternary caldera (Toba, Indonesia) clarified. *Geology* 19, 200–203.

1887 Chesner, C.A., 1998. Petrogenesis of the Toba Tuffs, Sumatra, Indonesia. *Journal of*

1888 *Petrology* 29 (3), 397–438.

1889 Clark, J.D., Williams, M.A.J., 1987. Paleoenvironments and prehistory in North Central

1890 India:

1891 a preliminary report. In: Jacobsen, J. (Ed.), *Studies in the Archaeology of India and*

1892 *Pakistan*. Aris and Phillips Ltd, Warminster, pp. 19–41.

1893 Dassarma, D.C., Biswas, S., 1977. Newly discovered late Quaternary vertebrates from

1894 the terraced alluvial fills of the Son valley. *Indian Journal of Earth Sciences* 4 (2), 122–

1895 136.

1896 Dehn, J., Farrell, J.W., Schmincke, H.-U., 1991. Neogene tephrochronology from Site
 1897 758 on Northern Ninetyeast Ridge: Indonesian arc volcanism of the past 5 Ma.
 1898 Proceedings of the Ocean Drilling Program, Scientific Results 21, 273–295.
 1899 Deo, S.G., Ghate, S., Rajaguru, S.N., Karmalkar, N., Kale, M., 2004. Discovery of an
 1900 Acheulian site in association with tephra at Rajbag, Taluka Supa, District Pune,
 1901 Maharashtra. *Man and Environment* 29 (1), 108.
 1902 Devdas, V., Meshram, S.N., 1991. Search for Quaternary ash bed in the Quaternary
 1903 Basins of Orissa. *Records of the Geological Survey of India* 124 (3), 40–42.
 1904 Diehl, J.F., Onstott, T.C., Chesner, C.A., Knight, M.D., 1987. No short reversals of
 1905 Brunhes age recorded in the Toba tuffs, north Sumatra, Indonesia. *Geophysical*
 1906 *Research Letters* 14, 753–756.
 1907 Endicott, P., Metspalu, M., Kivisild, T., 2007. Genetic evidence on modern human
 1908 dispersals in South Asia: Y chromosome and mitochondrial DNA perspectives. In:
 1909 Petraglia, M.D., Allchin, B. (Eds.), *The Evolution and History of Human Populations in*
 1910 *South Asia: Inter-disciplinary Studies in Archaeology, Biological Anthropology,*
 1911 *Linguistics and Genetics*. Springer, Netherlands, pp. 229–244.
 1912 Field, J., Petraglia, M., Lahr, M., 2006. The southern dispersal hypothesis and the South
 1913 Asian archaeological record: examination of dispersal routes through GIS analysis.
 1914 *Journal of Anthropological Archaeology*, in press.
 1915 Forster, P., Matsumura, S., 2005. Did early humans go North or South? *Science* 308
 1916 (5724), 965–966. Gagneux, P., Wills, C., Gerloff, U., Tautz, D., Morin, P.A., Boesch,
 1917 C., Fruth, B., Hohmann, G., Ryder, O.A., Woodruff, D.S., 1999. Mitochondrial
 1918 sequences show diverse evolutionary histories of African hominoids. *Proceedings of the*
 1919 *National Academy of Sciences* 96, 5077–5082.
 1920 Gathorne-Hardy, F., Harcourt-Smith, W., 2003. The super-eruption of Toba, did it
 1921 cause a human bottleneck? *Journal of Human Evolution* 45 (3), 227–230.
 1922 Goldberg, T.L., Ruvoilo, M., 1997. The geographic apportionment of mitochondrial
 1923 genetic diversity in East African chimpanzees, *Pan troglodytes schweinfurthii*.
 1924 *Molecular Biology and Evolution* 14 (9), 976–984.
 1925 Gurioli, L., Houghton, B.F., Cashman, K.V., Cioni, R., 2005. Complex changes in
 1926 eruption dynamics during the 79 AD eruption of Vesuvius. *Bulletin of Volcanology* 67,
 1927 144–159.
 1928 Harpending, H.C., Sherry, S.T., Rogers, A.R., Stoneking, M., 1993. The genetic
 1929 structure of ancient human populations. *Current Anthropology* 34 (4), 483–496.
 1930 Harpending, H.C., Batzer, M.A., Gurven, M., Jorden, L.B., Rogers, A.R., Sherry, S.T.,
 1931 1998. Genetic traces of ancient demography. *Proceedings of the National Academy of*
 1932 *Sciences* 95 (4), 1961–1967.
 1933 Harpending, H., Rogers, A., 2000. Genetic perspectives on human origins and
 1934 differentiation. *Annual Review of Genomics and Human Genetics* 1, 361–385.
 1935 Horn, P., Müller-Sohnius, D., Storzer, D., Zöller, L., 1993. K-Ar-, fission-track-, and
 1936 thermoluminescence ages of Quaternary volcanic tuffs and their bearing on Acheulian
 1937 artifacts from Bori, Kukdi Valley, Pune District, India. *Zeitschrift der Deutschen*
 1938 *geologischen Gesellschaft* 144, 326–329.
 1939 Huang, C-Y., Zhao, M., Wang, C-C., Wei, G., 2001. Cooling of the South China Sea by
 1940 the Toba eruption and correlation with other climate proxies ~71,000 years ago.
 1941 *Geophysical Research Letters* 28 (20), 3915–3918.

1942 James, H.V.A., Petraglia, M.D., 2005. Modern human origins and the evolution of
 1943 behavior in the later Pleistocene record of South Asia. *Current Anthropology* 46, S3–
 1944 S27. Jones, S.C., Pal, J.N., 2005. The Middle Son valley and the Toba supervolcanic
 1945 eruption of 74 kyr BP: Youngest Toba Tuff deposits and Palaeolithic associations.
 1946 *Journal of inter-disciplinary Studies in History and Archaeology* 2 (1), in press. Kale,
 1947 V.S., Ganjoo, R.K., Rajaguru, S.N., Ota, S.B., 1986. A discovery of an Acheulian site at
 1948 Bori, Dist. Pune (Maharashtra). *Bulletin of the Deccan College Postgraduate and*
 1949 *Research Institute* 45, 47–55. Kale, V.S., Patil, D.N., Powar, N.J., Rajaguru, S.N., 1993.
 1950 Discovery of a volcanic ash bed in the alluvial sediments at Morgaon, Maharashtra.
 1951 *Man and Environment* 18 (2), 141–143. Kale, V.S., Joshi, V.U., Hire, P.S., 2004.
 1952 Palaeohydrological reconstructions based on analysis of a palaeochannel and Toba-ash
 1953 associated alluvial sediments in the Deccan Trap region, India. *Journal Geological*
 1954 *Society of India* 64, 481–489.
 1955 Kennedy, K.A.R., 1999. Paleoanthropology of South Asia. *Evolutionary Anthropology*
 1956 8 (5), 165–185.
 1957 Kennedy, K.A.R., 2000. God-apes and fossil men. *Paleoanthropology in South Asia*.
 1958 The University of Michigan Press, Ann Arbor.
 1959 Kennedy, K.A.R., 2001. Middle and Late Pleistocene hominids of South Asia. In:
 1960 Tobias,
 1961
 1962 Khan, A.A. (1984): Geological and Geomorphological studies around Tapti-Vagher
 1963 confluence district Jalgaon, Maharashtra. *Geol, Surv. India Rec.* V.113 pt 6 pp 99-109.
 1964 Khan, A.A. & Banerjee, S.N. (1984): Geology and Geomorphological studies in the
 1965 parts of Narmada Basin, Sehore district of M.P. Un Pub. Report. *Geol. Surv. India*.
 1966 Khan, A.A., and Rahate, D.N. (1990-91 & 1991-92): Geological and Geomorphological
 1967 studies in parts of Narmada Basin) parts Hoshangabad and Narsingpur district, M.P.
 1968 *Geol. Surv. Of India Unpublished Progress Report*.
 1969 Khan A.A and A. Sonakia (1992): Quaternary deposits of Narmada with special
 1970 reference to the Hominid Fossils, *Jour. Geol. Soci. Of India Vol. 39 No2*, pp 147-154
 1971 Khan, A.A, Rahate, D.N. (1991): Volcanic Ash from Quaternary deposits of Narmada
 1972 Valley Central India. *Proceed, of 78th session of Indian Sci. Cong. Association*.
 1973 (Abstract) pt. III pp 28-29
 1974 Rahate. D.N.Khan A.A and Banerjee S.N. (1985): Geomorphological and geological
 1975 studies of Quaternary sediments in collaboration with project CRUMANSONATA in
 1976 parts of Narmada valley, Sehore, Hoshangabad and Dewas districts, M.P. progress
 1977 report (Unpublished) *Geo. Surv. Ind.*
 1978 Khan, A. A; Rahate; D.N, Fahim, M; & Banerjee, S.N.(1991) : Evaluation of
 1979 Quaternary terrace of lower Narmada valley , Districts Sehore and Hoshangabad,
 1980 Madhya Pradesh
 1981 Khan; A.A. (1991): Geological studies of Harda – Barwaha basin in parts of Dewas,
 1982 Sehore, Hoshangabad and Khandwa districts with the Aid of Satellite imagery and
 1983 Remote Sensing Techniques, *Geol. Surv. Ind, Rec. Vol; 126 pt-6*
 1984 Khan, A.A; D.N, Fahim, M. and Banarjee, S.N. (1992): Evaluation of Geology and
 1985 Geomorphology in Central Narmada Valley (Districts Sehore and Hoshangabad,
 1986 Madhya Pradesh) Scientific Publishers, Jodhpur.

1987 Khatri, A.P (1966): Pleistocene mammalian fossils of the Narmada river valley and
1988 their horizons Asian Perspective 9.pp113-133
1989 N.L. Yadova and Khan , A.A. (1995): evaluation of Geology and structure of Eastern
1990 Harda- Barwah Basin, Madhya Pradesh using Remote Sensing Techniques with Field
1991 Checks. Jour.Ind.Soci. Remote Sensing (Photonirvachak)
1992 Khan, A.A. (1990): Geomorphology of Narmada Valley Of Jabalpur_ Handia Section
1993 Unpublished G.S.I Note.
1994 Khan, A.A. & Vatsa, U.S. (1987-88): Geoenviromental Appraisal of the area around
1995 Jabalpur City, Madhya Pradesh G.S.I... Rec. Vol 122 pt-pp 163-64.
1996 Khan.A.A (.2018) Quaternary deposits of Narmada valley in relation to
1997 Geomorphology and Neotectonic of the area with special reference to Hominid
1998 occurrence D.sc. Unpublished thesis submitted to Barkatullah University Bhopal, M.P.
1999 India.
2000
2001 P.V., Raath, M.A., Moggi-Cecchi, J., Doyle, G.A. (Eds.), Humanity from African
2002 Naissance to Coming Millennia – Colloquia in Human Biology and
2003 Palaeoanthropology. Florence University Press, Florence, pp. 167–174.
2004 Kerr, R.A., 1996. Volcano-Ice Age link discounted. Science 272, 817.
2005
2006 Kivisild, T., Rootsi, S., Metspalu, M., Mastana, S., Kaldma, K., Parik, J., Metspalu, E.,
2007 Adojaan, M., Tolk, H-V., Stepanov, V., Gölge, M., Usanga, E., Papiha, S.S.,
2008 Cinnioğlu, C., King, R., Cavalli- Sforza, L., Underhill, P.A., Villems, R., 2003. The
2009 genetic heritage of the earliest settlers persists both in Indian tribal and caste
2010 populations. American Journal of Human Genetics 72, 313–332.
2011 Korisettar, R., Mishra, S., Rajaguru, S.N., Gogte, V.D., Ganjoo, R.K., Venkatesan,
2012 T.R., Tandon, S.K., Somayajulu, B.L.K., Kale, V.S., 1988. Age of the Bori volcanic ash
2013 and Lower Palaeolithic culture of the Kukdi Valley, Maharashtra. Bulletin of the
2014 Deccan College Postgraduate and Research Institute 48, 135–138.
2015 Korisettar, R., Venkatesan, T.R., Mishra, S., Rajaguru, S.N., Somayajulu, B.L.K.,
2016 Tandon, S.K., Gogte, V.D., Ganjoo, R.K., Kale, V.S., 1989. Discovery of a tephra bed
2017 in the Quaternary alluvial sediments of Pune District (Maharashtra), Peninsular India.
2018 Current Science 58 (10), 564–567.
2019 Korisettar, R., 1994. Quaternary alluvial stratigraphy and sedimentation in the Upland
2020 Deccan Region, Western India. Man and Environment 19 (1–2), 29–41. Lahr, M.M.,
2021 Foley, R., 1994. Multiple dispersals and modern human origins. Evolutionary
2022 Anthropology 3, 48–60.
2023 Lee, M-Y., Chen, C-H., Wei, K-Y., Iizuka, Y, Carey, S., 2004. First Toba supereruption
2024 revival. Geology 32, 61–64.
2025 Macaulay, V., Hill, C., Achilli, A., Rengo, C., Clarke, D., Meehan, W., Blackburn, J.,
2026 Semino, O., Scozzari, R., Cruciani, F., Taha, A., Shaari, N.K., Raja, J.M., Ismail, P.,
2027 Zainuddin, Z., Goodwin, W., Bulbeck, D., Bandelt, H-J., Oppenheimer, S., Torroni, A.,
2028 Richards, M., 2005. Single, rapid coastal settlement of Asia revealed by analysis of
2029 complete mitochondrial genomes. Science 308 (5724), 1034–1036.
2030 Mason, B.G., Pyle, D.M., Oppenheimer, C., 2004. The size and frequency of the largest
2031 explosive eruptions on Earth. Bulletin of Volcanology 66, 735–748.

2032 McCormick, M. P., Thomason, L. W., Trepte, C. R., 1995. Atmospheric effects of the
2033 Mt Pinatubo eruption. *Nature* 373, 399–404.

2034 Mishra, S., Ghate, S., 1990. Ranjani: A Middle Palaeolithic site from Maharashtra. *Man*
2035 *and Environment* 15 (2), 23–27.

2036 Mishra, S., Rajaguru, S.N., 1994. Comment on “Toba ash on the Indian subcontinent
2037 and its implication for the correlation of Late Pleistocene alluvium”. *Quaternary*
2038 *Research* 41, 396–397.

2039 Mishra, S., Venkatesan, T.R., Rajaguru, S.N., Somayajulu, B.L.K., 1995. Earliest
2040 Acheulian industry from Peninsular India. *Current Anthropology*
2041 36 (5), 847–851.

2042 Mishra, S., Rajaguru, S.N., 1996. Comment on “New geochemical evidence for the
2043 Youngest Toba Tuff in India”. *Quaternary Research* 46, 340–341.

2044 Misra, V.N., 1995. Geoarchaeology of the Thar Desert, North West India. In: Wadia, S.,
2045 Korisettar, R., Kale, V.S. (Eds.), *Quaternary Environments and Geoarchaeology of*
2046 *India*. Geological Society of India: Bangalore, pp. 210–230.

2047 Newhall, C.G., Self, S., 1982. The Volcanic Explosivity Index (VEI): an estimate of the
2048 explosive magnitude for historical volcanism. *Journal of Geophysical Research* 87 (C2),
2049 1231–1238.

2050 Ninkovich, D. Shackleton, N.J., Abdel-Monem, A.A., Obradovich, J.D., Izett, G.,
2051 1978a. K-Ar age of the late Pleistocene eruption of Toba, north Sumatra. *Nature* 276,
2052 574–577.

2053 Ninkovich, D., Sparks, R.S.J., Ledbetter, M.T., 1978b. The exceptional magnitude and
2054 intensity of the Toba eruption, Sumatra: an example of the use of deep-sea tephra layers
2055 as a geological tool. *Bulletin Volcanologique* 41, 286–298.

2056 Oppenheimer, C., 2002. Limited global change due to the largest known Quaternary
2057 eruption, Toba ≈ 74 kyr BP? *Quaternary Science Reviews* 21, 1593–1609.

2058 Oppenheimer, S., 2003. *Out of Eden: The Peopling of the World*. Constable, London.

2059 Palanichamy, M.G., Sun, C., Agrawal, S., Bandelt, H-J., Kong, Q-P., Khan, F., Wang,
2060 C-Y., Chaudhuri, T.K., Palla, V., Zhang, Y-P., 2004. Phylogeny of Mitochondrial DNA
2061 macrohaplogroup N in India, based on complete sequencing: implications for the
2062 peopling of South Asia. *American Journal of Human Genetics* 75, 966–978.

2063 Pattan, J.N., Pearce, N.J.G., Banakar, V.K., Parthiban, G., 2002. Origin of ash in the
2064 Central Indian Ocean Basin and its implication for the volume estimate of the 74,000
2065 year BP Youngest Toba eruption. *Current Science* 83 (7), 889–893.

2066 Pattan, J.N., Shane, P., Pearce, N.J.G., Banakar, V.K., Parthiban, G., 2001. An
2067 occurrence of ~ 74 ka Youngest Toba Tephra from the Western Continental Margin of
2068 India. *Current Science* 80 (10), 1322–1326.

2069 Pattan, J.N., Shane, P., Banakar, V.K., 1999. New occurrence of Youngest Toba Tuff in
2070 abyssal sediments of the Central Indian Basin. *Marine Geology* 155, 243–248.

2071 Petraglia, M.D., 1998. The Lower Palaeolithic of India and its bearing on the Asian
2072 record. In: Petraglia, M.D., Korisettar, R. (Eds.), *Early Human Behaviour in Global*
2073 *Context: The Rise and Diversity of the Lower Palaeolithic Record*. Routledge, London
2074 and New York, pp. 343–390.

2075 Poddar, B.C., Verma, K.K., Tewari, M.P., Rahate, D.N., Khan, A.A., Sonakia, A.,
2076 Biswas, S., Nandi, A., Krishna, S.G., Dubey, U.S., Bhai, H.Y., Fahim, M., Sitaramaiah,

2077 Y., 1991. Narmada Valley Project. Records of Geological Survey of India 124 (6), 245–
2078 247.

2079 Quintana-Murci, L., Semino, O., Bandelt, H-J., Passarino, G., McElreavey, K.,
2080 Santachiara-Benerecetti, A.S., 1999. Genetic evidence of an early exit of *Homo*
2081 *sapiens sapiens* from Africa through eastern Africa. Nature Genetics 23, 437–441.

2082 Rajaguru, S.N., Kale, V.S., Badam, G.L., 1993. Quaternary fluvial systems in Upland
2083 Maharashtra. Current Science 64, 817–822. Rajaguru, S.N., Deo, S.G., Mishra, S.,
2084 Ghate, S., Naik, S., Shirvalkar, P., 2004. Geoarchaeological significance of the detrital
2085 laterite discovery in the Karha Basin, Pune District, Maharashtra. Man and Environment
2086 29 (1), 1–6.

2087 Rampino, M. R., Stothers, R.B., Self, S., 1988. Volcanic winters. Annual Review of
2088 Earth and
2089 Planetary Sciences 16, 73–99.

2090 Rampino, M. R., Self, S., 1992. Volcanic winter and accelerated glaciation following
2091 the Toba supereruption. Nature 359, 50–52.

2092 Rampino, M.R., Self, S., 1993a. Climate-volcanism feedback and the Toba eruption of
2093 ~74,000 years ago. Quaternary Research 40, 269–280.

2094 Rampino, M.R., Self, S., 1993b. Bottleneck in human evolution and the Toba eruption.
2095 Science 262 (5142), 1955.

2096 Rampino, M.R., Ambrose, S.H., 2000. Volcanic winter in the Garden of Eden: The
2097 Toba supereruption and the late Pleistocene human population crash. In: McCoy, F.W.,
2098 Heiken, G. (Eds.), Volcanic Hazards and Disasters in Human Antiquity. Geological
2099 Society of America Special Paper 345, pp. 71–82.

2100 Rampino, M.R., 2002. Supereruptions as a threat to civilisations on Earth-like planets.
2101 Icarus 156, 562–569.

2102 Rogers, A.R., Harpending, H., 1992. Population growth makes waves in the distribution
2103 of pairwise genetic differences. Molecular Biology and Evolution 9, 552–569.

2104 Rogers, A.R., Jorde, L.B., 1995. Genetic evidence on modern human origins. Human
2105 Biology 67 (1), 1–36.

2106 Rose, W.I., Chesner, C.A., 1987. Dispersal of ash in the great Toba eruption, 75 ka.
2107 Geology 15, 913–917.

2108 Rose, W.I., Chesner, C.A., 1990. Worldwide dispersal of ash and gases from earth's
2109 largest known eruption: Toba, Sumatra, 75 ka. Palaeogeography, Paleoclimatology,
2110 Palaeoecology 89, 269–275.

2111 Rosi, M., Vezzoli, L., Aleotti, P., Censi, M. De., 1996. Interaction between caldera
2112 collapse and eruptive dynamics during the Campanian Ignimbrite eruption, Phlegraean
2113 Fields, Italy. Bulletin of Volcanology 57 (7), 541–554.

2114 Schulz, H., von Rad, U., Erlenkeuser, H., 1998. Correlation between Arabian Sea and
2115 Greenland climate oscillations of the past 110,000 years. Nature 393, 54–57.

2116 Schulz, H., Emeis, K-C., Erlenkeuser, H., von Rad, U., Rolf, C., 2002. The Toba
2117 volcanic event and interstadial/stadial climates at the Marine Isotopic Stage 5 to 4
2118 transition in the Northern Indian Ocean. Quaternary Research 57, 22–31.

2119 Self, S., Zhao, J-X., Holasek, R.E., Torres, R.C., King A.J., 1996. The atmospheric
2120 impact of the 1991 Mount Pinatubo eruption. In: Newhall, C.G., Punongbayan, R.S.
2121 (Eds.), Fire and Mud: Eruptions and Lahars of Mount Pinatubo, Philippines. University
2122 of Washington Press, Seattle, pp. 1089–1115.

2123 Shane, P., Westgate, J., Williams, M., Korisettar, R., 1995. New geochemical evidence
 2124 for the youngest Toba tuff in India. *Quaternary Research* 44, 200–204.
 2125 Shane, P., Westgate, J., Williams, M., Korisettar, R., 1996. Reply to Comments by S.
 2126 Mishra and S.N. Rajaguru on “New Geochemical Evidence for the Youngest Toba Tuff
 2127 in India”. *Quaternary Research* 46, 342–343.
 2128 Sharma, G.R., 1980. History to prehistory. *Archaeology of the Ganga Valley and the*
 2129 *Vindhyas*. Department of Ancient History, Culture and Archaeology, University of
 2130 Allahabad, Allahabad.
 2131 Sharma, G.R., Clark, J.D., 1983. Introduction. In: Sharma, G.R., Clark, J.D. (Eds.),
 2132 *Palaeoenvironments and Prehistory in the Middle Son Valley*.
 2133 Abinash Prakashan, Allahabad, pp. 1–8.
 2134 Sherry, S.T., Rogers, A.R., Harpending, H., Soodyall, H., Jenkins, T., Stoneking, M.,
 2135 1994. Mismatch distributions of mtDNA reveal recent human population expansions.
 2136 *Human Biology* 66 (5), 761–775.
 2137 Smith, R.L., Bailey, R.A., 1968. Resurgent cauldrons. *Geological Society of America*
 2138 *Memoir* 116, 613–662.
 2139 Sonakia, A., 1984. The skull-cap of early man and associated mammalian fauna from
 2140 Narmada valley alluvium, Hoshangabad area, Madhya Pradesh (India). *Records of the*
 2141 *Geological Survey of India* 113 (6), 159–172.
 2142 Sonakia, A., Biswas, S., 1998. Antiquity of the Narmada *Homo erectus*, the early man
 2143 of India. *Current Science* 75 (4), 391–392.
 2144 Song, S.-R., Chen, C.-H., Lee, M.-Y., Yang, T.F., Iizuka, Y., Wei, K.-Y., 2000. Newly
 2145 discovered eastern dispersal of the youngest Toba Tuff. *Marine Geology* 167, 303–312.
 2146 Stothers, R.B., 1984. The great Tambora eruption in 1815 and its aftermath. *Science*
 2147 224, 1191–1198.
 2148 Supekar, A.G., 1968. Pleistocene stratigraphy and prehistoric archaeology of the central
 2149 Narmada basin. Ph.D. Dissertation, Deccan College Postgraduate and Research
 2150 Institute, Poona. Thangaraj, K., Chaubey, G., Kivisild, T., Reddy, A.G., Singh, V.K.,
 2151 Rasalkar, A.A., Singh, L., 2005.
 2152 Reconstructing the origin of Andaman islanders. *Science* 308 (5724), 996.
 2153 Turco R.P., Toon O.B., Ackerman T.P., Pollack J.B., Sagan C., 1983. Nuclear winter:
 2154 global consequences of multiple nuclear explosions. *Science* 222, 1283–1292.
 2155 Turco R.P., Toon O.B., Ackerman T.P., Pollack J.B., Sagan C., 1990. Climate and
 2156 smoke: an appraisal of nuclear winter. *Science* 247, 166–176.
 2157 Underhill, P.A., Passarino, G., Lin, A.A., Shen, P., Lahr, M.M., Foley, R.A., Oefner,
 2158 P.J., Cavalli-Sforza, L.L., 2001. The phylogeography of Y chromosome binary
 2159 haplotypes and the origins of modern human populations. *Annals of Human Genetics*
 2160 65, 43–62.
 2161 van Bemmelen, R.W., 1939. The volcano-tectonic origin of Lake Toba (North
 2162 Sumatra). *De Ingenieur Nederlandsch-Indie* 6, 126–140.
 2163 Wakankar, V.S., 2002. Burial systems in Bhimbetka. In: Kennedy, K.A.R., Lukacs,
 2164 J.R., Misra, V.N. (Eds.), *The Biological Anthropology of Human*
 2165 *Skeletal Remains from Bhimbetka, Central India*. Indian Society for Prehistoric and
 2166 *Quaternary Studies*, Pune, pp. 1–5.
 2167 Watson, E., Forster, P., Richards, M., Bandelt, H.-J., 1997. Mitochondrial footprints of
 2168 human expansions in Africa. *American Journal of Human Genetics* 61, 691–704.

Westgate, J.A., Shane P.A.R., Pearce, N.J.G., Perkins, W.T., Korisettar, R., Chesner, C.A., Williams, M.A.J., Acharyya, S.K., 1998. All Toba tephra occurrences across peninsular India belong to the 75,000 yr B.P. eruption. *Quaternary Research* 50, 107–112.

Williams, M.A.J., 2004. Environmental impacts of extreme events: the Toba mega-eruption, volcanic winter and the near demise of humans. *Allahabad Journal of interdisciplinary Studies in History and Archaeology* 1 (1), 118–119.

Williams, M.A.J., Clarke, M.F., 1984. Late Quaternary Environments in north-central India. *Nature* 308, 633–635.

Williams, M.A.J., Clarke, M.F., 1995. Quaternary geology and prehistoric environments in the Son and Belan valleys, north central India. In: Wadia, S., Korisettar, R., Kale, V.S. (Eds.), *Quaternary Environments and Geoarchaeology of India*. Geological Society of India, Bangalore, pp. 282–308.


Williams, M.A.J., Royce, K., 1982. Quaternary geology of the Middle Son Valley, north central India: implications for prehistoric archaeology. *Palaeogeography, Palaeoclimatology, Palaeoecology* 38, 139–162.

Williams, M.A.J., Royce, K., 1983. Alluvial history of the Middle Son Valley, north central India. In: Sharma, G.R., Clark, J.D. (Eds.), *Palaeoenvironments and Prehistory in the Middle Son Valley*. Abinash Prakashan, Allahabad, pp. 9–21.

Zielinski, G.A., Mayewski, P.A., Meeker, L.D., Whitlow, S., Twickler, M., 1996a. A 110,000-year record of explosive volcanism from the GISP2 (Greenland) ice core. *Quaternary Research* 45, 109–118.


Zielinski, G.A., Mayewski, P.A., Meeker, L.D., Whitlow, S., Twickler, M., Taylor, K., 1996b. Potential atmospheric impact of the Toba mega-eruption ~71,000 years ago. *Geophysical Research Letters* 23, 837–840.

Zielinski, G.A., 2000. Use of paleo-records in determining variability within the volcanismclimate system. *Quaternary Science Reviews* 19, 417–438.




COMPOSITE LITHOSTRATIGRAPHIC SECTION OF QUATERNARY DEPOSITS OF NARMADA VALLEY


SEHORE AND HOSHANGABAD DISTRICTS, MADHYA PRADESH, INDIA




2219
2220




2221
2222
2223
2224



2225
2226
2227




2228
2229
2230




2231
2232
2233

Plate/Figure No.AF-4



2234
2235
2236


Plate/Figure No.AF-5



2237
2238


2239
2240

Plate/Figure No.AF-6




2241
2242
2243


Plate/Figure No.AF-7





2244
2245



2247
2248
2249




2250
2251




2257
2258

Plate/Figure No.AF-12





2259
2260
2261

Plate/Figure No.AF-13



2262
2263



UNDER PEER REVIEW IN IJAR